

TABLE X.—VAGINAL DELIVERIES IN CASES PREVIOUSLY DELIVERED BY SECTION.

*Number of Cases, 40. Maternal Mortality, 0. Gross Fœtal Loss, 1 { Stillbirths, 1.
Neonatal Deaths, 0.*

BOOKED CASES—continued.

Reg. No.	Age	Para	Weight of Infant	Method of Delivery	Variety of previous Section	Indication for previous Section	Result		REMARKS
							M.	C.	
5676	43	10th	lb. oz. 6 6	Spontaneous	Lower segment Cesarean section	Disproportion	L.	L.	First 5 normal deliveries of infants weighing 7½ lb., then 3 abortions: 9th Cesarean section during labour—infant 9½ lb. P.O.M. at 38 weeks with this pregnancy for unstable lie and contracted pelvis. Spontaneous delivery. Pelvimetry: Gynaecoid flat. T.C. 9.6 cms. T. 12.4 cms. All characteristics average.
12785	35	4th	6 12	Spontaneous	Lower segment Cesarean section	Placenta prævia	L.	L.	Lower segment Cesarean section on 3rd pregnancy for placenta prævia.
14327	42	10th	7 8	Spontaneous	Lower segment Cesarean section	Obstructed labour due to disproportion	L.	L.	Eight previous vaginal deliveries: 9th obstructed labour—Cesarean section in this hospital—infant 8 lb. 6 oz. Pelvimetry: Small gynaecoid. Spines prominent. Remaining characteristics average. T.C. 10.4 cms. T. 12.8 cms. This pregnancy 9 hours in labour. No disproportion.
14395	49	13th	6 14	Spontaneous	Lower segment Cesarean section	Unsuccessful pregnancies	L.	L.	Two previous Cesarean sections for unsuccessful pregnancies, 9 abortions and 1 N.N.D. following spontaneous delivery. Pelvimetry: Gynaecoid ———. T.C. 11.2 cms. T. 13.0 cms. This pregnancy spontaneous delivery following 5 hours in labour.

TABLE X.—VAGINAL DELIVERIES IN CASES PREVIOUSLY DELIVERED BY SECTION—*continued*.

Number of Cases, 40. *Maternal Mortality*, 0. *Gross Fœtal Loss*, 1 { *Stillbirths*, 1.
Neonatal Deaths, 0.

BOOKED CASES—*continued*.

Reg. No.	Age	Para	Weight of Infant	Method of Delivery	Variety of previous Section	Indication for previous Section	Result		REMARKS
							M.	C.	
16809	34	8th	lb. oz. 9 12	Spontaneous	Lower segment Casarean section	Placenta prævia	L.	L.	Pelvimetry: Gynæcoid with anthropoid tendency. All characteristics average. T.C., 12·8 cms. T. 12·8 cms. Fourteen hours labour.
17304	35	4th	7 1	Spontaneous	Lower segment Casarean section	Placenta prævia	L.	L.	Lower segment Casarean section for placenta prævia. Second spontaneous delivery since.
17472	28	8th	4 8	Leg brought down	Lower segment Casarean section	Placenta prævia	L.	S.B.	See Placenta Prævia—Table No. XV, page III.
19243	39	3rd	7 12	Spontaneous	Lower segment Casarean section	Disproportion and breech	L.	L.	First prolapse of cord with breech, S.B.; 2nd Casarean section for breech. Pelvimetry. Small gynæcoid. All characteristics average. T.C. 10·4 cms. T. 12·4 cms. Nineteen hours in labour.
22275	33	4th	7 8	Forceps	Lower segment Casarean section	Disproportion	L.	L.	First S/B, forceps delivery, infant 6 lb. 11 oz. (S.B. not related to delivery as F.H. disappeared in 1st stage). Second—ectopic pregnancy. Third—lower segment Casarean section for contracted pelvis, infant 7 lb. This pregnancy easy forceps delivery of living infant, 7 lb. 8 oz., after 13 hours in labour. Pelvimetry: Android flat pelvis. T.C. 10·4 cms. T. 12·8 cms.

TABLE X.—VAGINAL DELIVERIES IN CASES PREVIOUSLY DELIVERED BY SECTION—continued.

Number of Cases, 40. Maternal Mortality, 0. Gross Fetal Loss, 1. $\left\{ \begin{array}{l} \text{Stillbirths, 1.} \\ \text{Neonatal Deaths, 0.} \end{array} \right.$

BOOKED CASES—continued.

Reg. No.	Age	Para	Weight of Infant	Method of Delivery	Variety of previous Section	Indication for previous Section	Result		REMARKS
							M.	C.	
24747	35	3rd	lb. oz. 6 12	Spontaneous	Lower segment Caesarean section (2)	Disproportion	L.	L.	See Symphysiotomy—Table No. VIII, page 69.
24952	31	4th	8 4	Forceps	Lower segment Caesarean section	Pre-eclampsia	L.	L.	Previous history: 1st—Caesarean section for pre-eclampsic toxæmia, living; 2nd—abortion; 3rd—eclampsia, 1 fit intra-partum, forceps delivery, S/B. All in this hospital. This pregnancy: admitted at 39 weeks, B.P. 130/105, trace albumin. Spontaneous onset of labour 4 days later. Steady rise in blood pressure and increase in albuminuria. Heavily sedated with paraldehyde, morphia, pethidine and mag. sulphate. When fully dilated B.P. 200/120, albumin+2—in labour 14 hours, Low forceps.
25184	31	5th	7 4	Spontaneous	Lower segment Caesarean section	Placenta prævia	L.	L.	Twelve hours labour.
27660	34	5th	9 12	Spontaneous	Lower segment Caesarean section	Pre-eclampsia	L.	L.	Pelvmetry: Gynæcoid. Sacrum well curved. Remaining characteristics average. T.C 12.0 cms. T. 13.6 cms. Seven and a half hours labour. Fourth—vaginal delivery following section.

TABLE X.—VAGINAL DELIVERIES IN CASES PREVIOUSLY DELIVERED BY SECTION—*continued*.

*Number of Cases, 40. Maternal Mortality, 0. Gross Fœtal Loss, 1 { Stillbirths, 1.
Neonatal Deaths, 0.*

BOOKED CASES—*continued*.

Reg. No.	Age	Para	Weight of Infant	Method of Delivery	Variety of previous Section	Indication for previous Section	Result		REMARKS
							M.	C.	
27749	37	4th	lb. oz. 7 4	Spontaneous	Lower segment Cæsarean section	Placenta prævia	L.	L.	First—normal delivery; 2nd—abortion; 3rd—Cæsarean section for placenta prævia. Duration of labour 53 hours.
30308	32	3rd	6 0	Forceps	Lower segment Cæsarean section	Disproportion	L.	L.	First—prolapsed cord, perforation and extrac-tion; 2nd—L.S.C.S. for fœtal and maternal distress following 9 hours' trial of labour—infant 7 lb. 7 oz. Admitted at term. Pel-vimetry: Small gynaecoid. Sub-pubic—male type. Remaining characteristics average. T.C. 9.6 cms. T. 11.6 cms. Head free and not entering well. Trial of labour decided upon with symphysiotomy early in labour if progress not satisfactory. Spontaneous onset of labour with good advance. Examined when os 2 fingers. No disproportion. When os full fœtal distress, head in mid-strait. Easy forceps.
31106	29	2nd	8 2	Spontaneous	Lower segment Cæsarean section	Eclampsia	L.	L.	Duration of labour 7 hours.

TABLE X.—VAGINAL DELIVERIES IN CASES PREVIOUSLY DELIVERED BY SECTION—*continued*.

Number of Cases, 40. Maternal Mortality, 0. Gross Fatal Loss, 1 { Stillbirths, 1.
Neonatal Deaths, 0.

BOOKED CASES—*continued*.

Reg. No.	Age	Para	Weight of Infant	Method of Delivery	Variety of previous Section	Indication for previous Section	Result		REMARKS
							M.	C.	
31177	34	6th	lb. oz. 7 12	Breech	Lower segment Casarean section	Placenta prævia	L.	L.	Pelvimetry: Gynecoid. Prominent spines. Sacrum well curved. Sub-pubic good. T.C. 11.2 cms. T. 13.2 cms. Duration of labour 4 hours.
31754	29	3rd	9 2	Spontaneous	Lower segment Casarean section	Accidental hæmorrhage and pre-eclampsia	L.	L.	Second vaginal delivery. Pelvimetry: Gynecoid. All characteristics average. T.C. 11.2 cms. T. 12.8 cms. Duration of labour 6 hours.
32118	38	5th	6 15	Spontaneous	Lower segment Casarean section	Disproportion	L.	L.	First—forceps, S/B.; 2nd—breech, S/B.; 3rd—lower segment Casarean section for obstetrical history. All outside our services. Fourth—normal delivery of living infant in this hospital. Pelvimetry—gynecoid. Spines prominent. Remaining characteristics average. T.C. 12.0 cms. T. 13.2 cms. Duration of labour 10 hours.
33093	23	3rd	6 10	Spontaneous	Lower segment Casarean section (elective)	Pre-eclampsia	L.	L.	Pelvimetry: Anthropoid. All characteristics average. T.C. 13.6 cms. T. 12.4 cms. Second—vaginal delivery.
33190	35	3rd	7 4	Spontaneous	Lower segment Casarean section	Disproportion	L.	L.	See Symphysiotomy—Table VIII, page 69.

TABLE X.—VAGINAL DELIVERIES IN CASES PREVIOUSLY DELIVERED BY SECTION—*continued*.

*Number of Cases, 40. Maternal Mortality, 0. Gross Fœtal Loss, 1 { Stillbirths, 1.
Neonatal Deaths, 0.*

BOOKED CASES—*continued*.

Reg. No.	Age	Para	Weight of Infant	Method of Delivery	Variety of previous Section	Indication for previous Section	Result		REMARKS
							M.	C.	
34298	31	4th	lb. oz. 10 15	Forceps	Lower segment Cæsarean section	Placenta prævia	L.	L.	Pelvimetry: Large gynæcoid. All characteristics average. T.C. 13.6 cms. T. 13.6 cms. Duration of labour 5 hours.
34830	25	2nd	6 6	Spontaneous	Lower segment Cæsarean section	Disproportion	L.	L.	See Sympysiotomy—Table VIII, page 69.
34788	32	2nd	8 2	Spontaneous	Lower segment Cæsarean section	Pre-eclamptic toxæmia and inertia	L.	L.	Antenatal care from 38th week for mild pre-eclampsia. X-ray pelvimetry: Gynæcoid. Outlet somewhat diminished by forward inclination of sacrum. T.C. 11.4 cms. T. 13.0 cms. No disproportion. Spontaneous onset of labour at term with spontaneous delivery following easy labour of 7 hours' duration.
35580	32	7th	7 10	Spontaneous	Lower segment Cæsarean section	Placenta prævia	L.	L.	—

TABLE X.—VAGINAL DELIVERIES IN CASES PREVIOUSLY DELIVERED BY SECTION—*continued*.

Number of Cases, 40. *Maternal Mortality*, 0. *Gross Fœtal Loss*, 1 { *Stillbirths*, 1.
Neonatal Deaths, 0.

BOOKED CASES—*continued*.

Reg. No.	Age	Para	Weight of Infant	Method of Delivery	Variety of previous Section	Indication for previous Section	Result		REMARKS
							M.	C.	
35611	25	2nd	lb. oz. 9 11	Forceps	Lower segment Cesarean section	Toxæmia and free head at term	L.	L.	Spontaneous onset of labour at term (43 weeks?). Head free at onset of labour, but good progress made. When 3 hours fully dilated, no advance. P.O.P. Easy manual rotation and forceps.
36047	43	5th	7 8	Forceps	Not known	Uterine inertia	L.	L.	Three abortions. One Cesarean section for uterine inertia outside our service. When head showing and in the 2nd stage 35 minutes, fetal distress developed.
38715	26	3rd	9 10	Spontaneous	Lower segment Cesarean section	Pre-eclampsia	L.	L.	First delivery lower segment Cesarean section for toxæmia; 2nd vaginal delivery. Duration of this labour 7 hours.
40937	34	2nd	8 1	Spontaneous	Unknown	Inertia	L.	L.	Duration of labour, 10 hours.
NON-BOOKED CASES.									
25531	28	5th	7 13	Spontaneous	Lower segment section (2)	Disproportion.	L.	L.	First—forceps delivery, S.B. following failed forceps outside our service; 2nd—L.S.C.S., elective at term; 3rd—elective section at term; 4th—symphysiotomy, spontaneous delivery; 5th—spontaneous after 5 hours' labour.

TABLE X.—VAGINAL DELIVERIES IN CASES PREVIOUSLY DELIVERED BY SECTION—*continued*.

*Number of Cases, 40. Maternal Mortality, 0. Gross Fetal Loss, 1 { Stillbirths, 1.
Neonatal Deaths, 0.*

NON-BOOKED CASES—*continued*.

Reg. No.	Age	Para	Weight of Infant	Method of Delivery	Variety of previous Section	Indication for previous Section	Result		REMARKS
							M.	C.	
26338	24	4th	lb. oz. 6 4	Spontaneous	Lower segment Casarean section	Contracted pelvis	L.	L.	First—lower segment Casarean section following trial of labour of 36 hours; 2nd—elective symphysiotomy, duration of labour 8 hours; 3rd—normal delivery, duration of labour 3 hours 15 minutes. No locomotor or urinary disability.
31236	33	2nd	6 8	Forceps	Lower segment Casarean section	Cardiac disease	L.	L.	See Rheumatic Heart Disease—Table No. 1 (a), page 21.
31406	31	2nd	7 3	Spontaneous	Lower segment section	Cardiac disease	L.	L.	See Rheumatic Heart Disease, Table No. 1 (a), page 21. Duration of labour 6 hours.
31768	29	2nd	5 12	Forceps	Lower segment Casarean section	Cardiac disease	L.	L.	See Rheumatic Heart Disease Table No. 1 (a), page 21.
32653	38	2nd	6 0	Spontaneous	Lower segments Casarean section	Inertia	L.	L.	Spontaneous delivery without difficulty following 2 hours in labour. Previous Casarean section for inertia—100 hours in labour.
38317	33	4th	6 6	Unassisted breech	Lower segment Casarean section	Placenta prævia	L.	L.	First miscarriage at 18 weeks; 2nd L.S.C.S. for placenta prævia; 3rd normal delivery of premature infant. N.N.D.

TABLE X.—VAGINAL DELIVERIES IN CASES PREVIOUSLY DELIVERED BY SECTION—*continued*.

Number of Cases, 40. Maternal Mortality, 0. Gross Fetal Loss, 1 { Stillbirths, 1. Neonatal Deaths, 0.

NON-BOOKED CASES—*continued*.

Reg. No.	Age	Para	Weight of Infant	Method of Delivery	Variety of previous Section	Indication for previous Section	Result		REMARKS
							M.	C.	
38788	25	4th	lb. oz. 7 12	Forceps	Two classical Casarean sections	Unknown	L.	L.	First abortion; 2nd Classical section at 26 weeks following appendicectomy; 3rd repeat classical section. All outside our service. Pelvimetry; Gynaecoid. T.C. 11.2 cms. T. 12.8 cms. Easy forceps delivery for foetal distress after 15 hours' labour.
39222	29	2nd	6 14	Forceps	Lower segment Casarean section	Disproportions	L.	L.	Previous section for disproportion outside our service—hydrocephalus, S.B. Pelvimetry: Gynaecoid flat. T.C. 11.2 cms. T. 12.4 cms. Twenty-four hours in labour with this pregnancy—easy delivery.
40427	29	2nd	6 13	Forceps	Classical Casarean section	Concealed accidental haemorrhage	L.	L.	First pregnancy, very severe concealed accidental haemorrhage with classical section. No toxæmia this pregnancy. Pelvimetry: Antherpoid. Prominent spines. Long sacrum T.C. 13.4 cms. T. 12.8 cms. Accidental haemorrhage when os full and head on pelvis floor—immediate forceps. Duration of labour 36 hours.
40554	26	2nd	8 12	Spontaneous	Classical Casarean section	Inertia	L.	L.	Antenatal care for pyelitis. Previous classical section outside our service for inertia. Infant stillborn. Pelvimetry. T.C. 13.2 cms. T. 14.0 cms. Head engaged, 5 hours in labour. P.P.H. No evidence of rupture. Uterus plugged.