

ANTIAPOPTOTIC PROTEINS

The Bcl-2 and Inhibitor of Apoptosis Protein Families

Quinn L. Deveraux, PhD, Sharon L. Schendel, PhD,
and John C. Reed, MD, PhD

Apoptosis, or programmed cell death, ensures that the genesis of new cells by way of cell division is controlled appropriately and offset by cell loss.¹⁰³ Cell death is a natural accompaniment of the physiology of self-renewing tissues in which terminally differentiated cells turn over to be replaced by new cells, such as in the skin, intestine, immune system, mammary gland, and uterus. Developmental organization requires removal of some cells for achieving the final desired structures, such as digits and canals, and ensuring proper cytoarchitectures of most organs, such as the kidney, heart, and brain. Moreover, elimination of cells compromised by viral infection, oxidation, hypoxia, or DNA damage is important for maintaining healthy tissues.³²

Apoptotic defects are the primary lesion in some types of cancer and leukemia, allowing malignant cells to survive longer than their intended life span and endowing these cells

with a selective survival advantage relative to their normal counterparts. Therefore, cell death can contribute to neoplastic expansion in the absence of increased cellular division.⁸⁴ Excessive apoptosis, however, has been implicated in spinal muscular atrophy (SMA), autoimmune disease, acquired immunodeficiency syndrome (AIDS), Alzheimer's disease, myocardial infarction, and stroke.³² Thus, appropriate regulation of apoptosis is critical to development and maintenance of normal tissues.

CELL DEATH PATHWAYS

The biochemical basis for most of the morphologic changes associated with apoptosis, such as membrane blebbing, chromatin condensation, and DNA fragmentation, can be traced to the actions of a family of cysteine proteases called the *caspases*. Once activated, the caspases can cleave cytoskeletal and nuclear matrix-associated proteins that are required for cellular integrity, such as lamins, inhibitors of DNA degradation enzymes, such as inhibitor of caspase-activated deoxyribonuclease (ICAD), and DNA repair enzymes,

This work is supported by a postdoctoral fellowship the Leukemia and Lymphoma Society (QLD) and by a postdoctoral fellowship DAMD-17-981-8167 from United States Army Medical Research and Materiel Command (SLS).

From the The Burnham Institute Program on Apoptosis and Cell Death Research, La Jolla, California

such as DNA-protein kinase (PK) and poly (adenoside diphosphate [ADP]-ribose) polymerase (PARP), to name only a few of the known caspase substrates. Caspase-mediated cleavage of these and other cellular proteins facilitates the ordered dismantling of the cell and the irreversible destruction of its genome.

Most caspases are synthesized initially as inactive precursors (zymogens) that undergo proteolytic processing to produce subunits that form the active heterotetrameric protease. In mammalian cells, activation of the caspase zymogens has been reported to occur through at least three independent mechanisms: (1) cleavage by upstream active caspases; (2) cleavage by granzyme B, an aspartate-specific serine protease found in the granules of cytolytic T-cells; and (3) autoprocessing of zymogens with assistance from other caspase-interacting proteins, which can occur in either a cis- or transmanner (reviewed in references 90 and 105).

Cellular and genotoxic stresses, such as those inflicted by chemotherapeutic drugs and radiation, can induce the expression of proapoptotic members of the Bcl-2 family. At least one of these proteins, Bax, has been shown to promote the release of cytochrome c from mitochondria (reviewed in references 37 and 86). Once released, cytochrome c catalyzes the formation of the *apoptosome* a deoxyadenosine (dATP)- or adenosine triphosphate (ATP)-dependent complex consisting of the apoptosis protease activating factor 1 (Apaf-1) protein and procaspase-9.^{57, 59, 86, 136} Apoptosome formation results in activation of bound procaspase-9, which then can directly cleave and activate procaspase-3, resulting in additional caspase activation and apoptosis (Fig. 1). In addition to fostering procaspase-3 activation, caspase-9 may have other functions, because in some cells, it can translocate to the nucleus following apoptotic stimulus.⁵³

Another prototypical mechanism for triggering autoprocessing and activation of caspases requires the recruitment of procaspase-8 to plasma membrane receptor complexes, such as Fas, which is a member of the tumor necrosis factor (TNF) family of cell death receptors. Procaspase-8 possesses approximately 1% the activity of the processed fully active protease. When brought into close ap-

position by oligomerization around Fas receptor complexes, these zymogens transprocess each other, yielding autonomous, active caspase-8.^{47, 67, 104} Once activated, caspase-8 then can directly activate procaspase-3 and other downstream caspases (see Fig. 1).

Bcl-2 FAMILY PROTEINS

In certain apoptotic pathways, Bcl-2 family proteins govern a cell's decision to heed or ignore death signals. The progenitor of this family is the Bcl-2 protein, first identified at a chromosomal breakpoint in human B-cell lymphomas.¹¹⁷ The family subsequently has expanded and now includes at least 18 members with representatives from mammalian species, viruses, and *Caenorhabditis elegans*.

The family can be divided into two groups: (1) antiapoptotic, which includes Bcl-2 and Bcl-x_L; and (2) proapoptotic, which includes Bax and Bid (Fig. 2). As in all families, some members are of close relation, whereas other members can claim only distant relation. In this respect, all family members share pockets of sequence similarity, denoted BH1, -2, -3, and -4.¹⁶ The BH3 domain is common to almost all family members, and some family members, including Bid, Bad, and Hrk, have the BH3 domain as their only link to the family. All of these so-called "BH3-only" proteins are proapoptotic. The BH4 domain is unique to antiapoptotic proteins and is found at the extreme amino terminus of these proteins.

Deletion mutagenesis has suggested that these regions of sequence similarity are important in regulating protein-protein interactions between the family members to form either homo- or heterodimers.^{99, 129, 133} The fate of cell seems to lie with the relative amounts of the pro- and antiapoptotic proteins and the identity of the predominating protein complexes.⁵²

FORM FOLLOWING FUNCTION?

Despite the important role the Bcl-2 protein family plays in cell death pathways, the exact biochemical mechanism by which the Bcl-2




Figure 1. Apoptotic pathways. Several members of the tumor necrosis factor (TNF) family of death receptors (Fas) recruit caspase-8 to their cytosolic domains on ligand binding, resulting in proteolytic activation of this proximal caspase. Once activated, caspase-8 can induce directly or indirectly the activation of caspase-3. A separate pathway for caspase activation is dependent on mitochondria and involves release of cytochrome c (cyto c) from this organelle into the cytosol—an event often mediated by proapoptotic Bcl-2 family members, such as Bax. On entering the cytosol, cyto c catalyzes the formation an Apaf-1/caspase-9 complex (apoptosome). The apoptosome activates caspase-3, initiating a caspase cascade that executes the apoptotic program. In some cells, Fas-induced cell death depends on mitochondria. In these cells, activated caspase-8 cleaves Bid, which then targets the mitochondria and promotes cyto-c release. Inhibitor of apoptosis proteins (IAPs) can block each of these apoptotic pathways by directly inhibiting distinct caspases.

family of proteins modulates apoptosis remains unclear. A possibility for the biochemical function of the Bcl-2 protein family was suggested following the determination of the 3-dimensional structure of Bcl-x_L.⁷⁴

Bcl-x_L is a bundle of 7 helices arranged in three layers. The outer two layers of amphipathic helices enclose between them two central helices. These two helices are long (each approximately 20 residues) and have a pronounced hydrophobicity.

The Bcl-x_L structure bears a strong resemblance to the previously determined structures of the membrane translocation domains of the bacterial toxins diphtheria and colicins A and E1.⁷⁴ The diphtheria toxin membrane translocation domain forms a channel in the endosomal membrane, through which the ADP-ribosylating subunit passes,⁶⁰ while the colicins kill sensitive *Escherichia coli* strains by way of the formation of a highly conductive ion channel that depolarizes the target cell's plasma membrane, resulting in cell death.

Although the colicin and diphtheria toxins attack very different organisms, the structures of these toxins use a similar *cloak-and-dagger* strategy in which the hydrophobic *dagger* is hidden within a *cloak* of amphipathic helices that allows these proteins to exist in a soluble state, but under certain conditions the hydrophobic dagger is *unsheathed* allowing the protein to insert into membranes.

The coordinates of the three-dimensional Bcl-x_L structure can be used as a scaffold on which to build models for the other Bcl-2 family members. Despite their opposing functions and their isolated regions of sequence similarity, models for Bcl-2 and Bax could be built using the Bcl-x_L structure for a guide.⁹³ Both proteins share a similar silhouette, in that they are α -helical bundles having at their core two long central hydrophobic helices. The three-dimensional structure for Bid has also been determined, and it also shows the same characteristics.^{16, 70} The structural similarity between these Bcl-2 family members




Figure 2. Bcl-2 protein family homology domain organization. Two subgroups are indicated: antiapoptotic (A), which discourages cell death, and proapoptotic (B) which promotes cell death. The relative locations of the homology domains (BH1, -III 2, -III 3, and -III 4) and the C-terminal transmembrane domain (TM) are indicated. For the pore-forming proteins, the putative pore-forming helices fall between the BH1 and BH2 domains, with the exception of Bid.

and the pore-forming domains of bacterial toxins suggests that the Bcl-2 protein family may possess pore-forming potential.

In addition to similarity to pore-forming proteins, the structure of Bcl-x_L reveals other details about how this protein is regulated. For example, a long loop lacking defined secondary structure intervenes between the first and second helices of Bcl-x_L. This loop sequence is a feature of only the antiapoptotic family members, and, although it is dispensable to their protective action, this region may represent a regulatory domain, because it is vulnerable to protein digestion¹⁰⁶ and possesses several phosphorylation sites.¹⁴ Thus, posttranslational modifications or conformational changes occurring in this domain may act as a means for modulating the protective effects of Bcl-2 and Bcl-x_L.⁷¹

The BH1, -2, and -3 domains cluster together on one side of the molecule, forming a hydrophobic cleft. This structural feature, along with results from site-directed mutagenesis studies, suggests that this patch may participate in hydrophobic protein-protein interactions between Bcl-2 family members. A peptide corresponding to the BH3 domain of the proapoptotic family member, Bak, was able to nestle into the cleft, which is just wide enough to accommodate the α -helix of dimerizing partners.⁹¹

IN VITRO CHANNEL FORMATION

In order for the Bcl-2 family proteins to form pores they must have α -helices that are of sufficient length to completely span a

membrane bilayer, and these helices must be largely lacking in charged residues. Each residue of an α -helix donates 1.5 Å to the overall helix length. If a typical lipid bilayer has a hydrophobic cross-section of approximately 30 Å,⁷³ then it follows that the helix must contain at least 20 residues. Bcl-x_L has two α -helices that satisfy this requirement: the two central helices. Although two α -helices are insufficient to enclose a channel lumen, the tendency of the Bcl-2 protein family to form dimers suggests that two or more molecules could coalesce to form a channel (reviewed in reference 93).

In vitro channel-forming ability has been demonstrated by several family members: Bcl-2, Bcl-x_L, Bax, and, discussed later, Bid.^{4, 72, 94–95, 97} Bcl-2, Bcl-x_L, and Bax each form channels in large unilamellar liposomes and in planar bilayer systems, from which information about channel characteristics, such as conductance and ion selectivity, can be gained. Each protein displayed a population of channels with varied conductance states, ranging from 20 pS to nearly 2000 pS.⁹³ The colicin E1 channel also produces 20 pS channels in planar lipid bilayers,¹² and this channel is predicted to be composed of four transmembrane α -helices, two hydrophobic and two amphipathic.²⁰ By analogy, the 20 pS channel formed by the Bcl-2 protein family members also could consist of a four-helix bundle, but in contrast to colicin, which has a monomeric channel, it is likely that two molecules must donate the their central fifth and sixth α -helices to form a conductive channel. Indeed, the channel-forming activity appears to lie within these helices, because their removal abolishes the channel activity for Bcl-2 and Bax.^{68, 94} In the case of Bax, oligomerization appears to be a crucial event for channel formation as the insertion-competent state of the protein elutes in gel filtration experiments as an oligomer of 160 kDa suggesting that at least six subunits of Bax may participate in channel formation.⁵ This oligomerization state may also account for the larger conductances observed for Bax channels of up to 2000 pS.⁴

A PORE ALTERNATIVE

That several Bcl-2 family members form channels in vitro while claiming allegiance to

opposing family branches, that is, anti- and proapoptotic, at first appears to be a paradox. Their similar structures may confer on most family members the ability to induce ion conductance in vitro, but this general trait may acquire a different purpose in vivo. It is possible that ion conductance, while detectable in vitro, may not be relevant to the Bcl-2 protein family function. The ability of these proteins to exist either in a soluble, globular form, or a membrane-inserted form may represent a means by which these proteins take on alternative conformations and in doing so, expose parts of the proteins that normally may be tucked out of reach. Such a situation could apply to regulation of protein-protein interactions. Bcl-2 and Bcl-x_L have displayed an affinity for a variety of proteins, including the protein kinase Raf-1, the protein phosphatase calcineurin, the *C. elegans* protein CED-4, and the Hsp70 modulating protein BAG-1 (Reviewed in reference 85). These protein-protein interactions are governed by the N-terminal BH4 domain¹³³ as mutations in this region abolish these interactions. The BH4 domain also appears important to formation of Bcl-2–Bax heterodimers as site-directed mutations in the Bcl-2 BH4 domain abolished Bcl-2–Bax interactions and the protective effect afforded by Bcl-2 against Bax and staurosporine-induced cell death.⁴¹ It is possible that in the uninserted soluble state, the BH4 domain is tucked against the protein and becomes accessible on insertion of the central hydrophobic helices.⁹⁴

A MATTER OF PLACEMENT: Bcl-2 FAMILY PROTEINS AND THE MITOCHONDRIA

The mitochondria, play significant roles in apoptosis regulation (Reviewed in references 37–38 and 119). Most Bcl-2 family proteins have at their C-terminus a stretch of approximately 20 hydrophobic residues, which seems to be critical to localize these proteins to mitochondria and to other cellular membranes, including the nuclear envelope and the endoplasmic reticulum (ER). Bcl-2 and Bcl-x_L appear to be permanently localized to the mitochondrial membrane,^{18, 54} while other Bcl-2 family proteins, largely the proapoptotic

members, such as Bax, are transient mitochondrial residents that change their cellular address from cytosolic to mitochondrial in response to several death signals.^{39, 125} The predicted hydrophobic central fifth and sixth α -helices of Bax seem to play a role in this change of address, because stripping these helices of charged residues (purported to line the aqueous lumen of a Bax channel) and substituting alanines resulted in a protein that was constitutively localized to mitochondria and hyperactive in its proapoptotic activity or "gain of function."⁷⁷

Although the Bcl-2 family proteins commonly are thought to inhabit only the outer mitochondrial membrane,¹¹⁹ immunoelectron microscopy revealed a nonuniform distribution of Bcl-2 in mitochondrial membranes, suggesting that this protein may be located preferentially at zones of adhesion, which join the outer and inner membranes,^{22, 54} a fact that could have importance in how these proteins might regulate the mitochondria's role in apoptosis.

As outlined previously, in some cell death pathways, escape of cytochrome c from the intermembrane space of mitochondria represents a key event in initiating the caspase activation cascade. Indeed, tissues from patients with end-stage human cardiomyopathy showed accumulation of cytosolic cytochrome c accompanied by caspase-3 activation.⁷⁶ Once liberated from the mitochondria, cytochrome c is free to participate in formation of the apoptosome.⁵⁷ In certain cells, other proteins that redistribute from the intermembrane space to the mitochondria include caspase-9 and caspase-3^{53, 109} and apoptosis inducing factor (AIF), which results in nuclear morphology changes.⁵¹ The mechanism by which these proteins pass into the cytoplasm remains unclear, although the Bcl-2 family proteins clearly regulate their escape.

The Bcl-2 protein family member Bax may provide a direct route for cytochrome c out of the mitochondria. Treatment of isolated mitochondria with recombinant Bax resulted in release of more than 30% of the total cytochrome c, suggesting that the Bax protein itself may be capable of forming a pore large enough to allow cytochrome c release.⁴⁸

Alternatively, mitochondrial swelling,

which eventually compromises outer membrane integrity, may result in cytochrome c leaking out into the cytosol. This swelling and subsequent rupture of the outer mitochondrial membrane could be induced directly through the channel activity of Bcl-2 family proteins,¹²⁰ or the Bcl-2 family could indirectly control mitochondrial volume by affecting the activity of the mitochondrial permeability transition pore (PTP).¹³⁵ The PTP pore allows passage of solutes with a molecular mass not exceeding 1500 Da. Although all of the components of PTP are not yet defined, the core participants appear to be the adenine nucleotide translocator (ANT) the voltage-dependent anion channel (VDAC).¹³⁵ ANT and VDAC are localized to the inner and outer mitochondrial membranes, respectively. A variety of parameters, including membrane potential, matrix pH, and oxidation state,⁸ affect the conductance state of the PTP. Opening of the PTP results in a rapid membrane depolarization.

Bcl-2 family proteins could regulate the cytochrome c release through interactions with proteins involved in the PTP. VDAC was reconstituted in liposomes and in the presence of recombinant proapoptotic proteins Bax and Bak the opening of VDAC was promoted, while Bcl-x_L appears to close the channel through direct binding. In cytochromec-loaded VDAC vesicles, Bax and Bak induced a loss of potential and cytochrome c release that could be inhibited by Bcl-x_L.¹⁰⁰ Although obtained from *in vitro* experiments, these results suggest that Bcl-2 family proteins may directly bind to VDAC and alter its activity, which should affect the activity of the PTP pore in mitochondria.

Another interaction that has been described is between Bax and ANT.¹¹ Again, ANT was reconstituted into lipid bilayers and its channel activity measured. On addition of Bax to these lipid bilayers, a composite channel is formed with an electrophysiological profile that differs from the channels formed by either Bax or ANT alone. This channel appears even under conditions where Bax has no detectable channel activity. In contrast, when reconstituted into lipid bilayers in the presence of Bcl-2, there is inhibition of channel formation.

The fact that ANT is inner membrane and that Bax is traditionally thought to have an outer mitochondrial localization poses some difficulty for thinking about this model. This can be remedied by the fact that the Bcl-2 family proteins do not appear to have a uniform mitochondrial distribution, but rather appear to cluster at adhesion sites where the outer and inner membrane are in contact.⁵⁴ An analogy can be drawn to the method of colicin action. In the case of colicins, many molecules may bind to the outer wall of the target *E. coli* cell, but very few access the inner membrane space, and only one colicin molecule seems to be necessary to deliver the lethal channel. Only those colicin molecules that bind to an outer membrane receptor, that is, associated with inner membrane-bound proteins and found at adhesion zones, seem to be capable of inserting to form their channel.⁹⁶ The same scenario also could exist for Bcl-2 family proteins. Most of the population may exist at the outer membrane surface, however, those molecules that are at contact sites, which themselves appear to be transient,⁹⁸ may be the active population in that they are in proper position to interact with PTP pore components.

CASPASE-8 Bid CLEAVAGE: A MITOCHONDRIAL LINK TO THE "Fas" TRACK

In response to Fas receptor ligation, procaspase-8 is recruited to the death receptor complex where local aggregation allows the processing of caspase-8 from the zymogen to active form within the death induced signaling complex (DISC), which includes in addition to procaspase-8 and Fas, Fas-associated death domain (FADD).⁷⁵ After activation at the DISC, caspase-8 is released and is available to activate downstream caspases, such as caspase-3 (see Fig. 1). There are two *tracks* a cell can follow with regards to DISC formation. Type 1 cells respond to Fas engagement by the activation of large amounts of caspase-8 by the DISC, whereas Type II cells have reduced DISC formation and consequently lower amounts of activated caspase-8. Examples of Type I and type II cells are lympho-

cytes and hepatocytes, respectively.⁹² The presence of cytosolic cytochrome c in compromised cardiac tissue and the expression of Bcl-2 in these cells suggests that cardiomyocytes might fall into the type II category. Type I cells cannot be rescued from cell death by Bcl-2 or Bcl-x_L overexpression, whereas type II cells can. This fact, along with a reduced $\Delta\psi_m$, suggests that type II cells may take a mitochondrial detour along their cell death pathway.

The amplification of Fas-mediated death signals by way of the mitochondria in type II cells suggested that there must be an intermediary substrate that caspase-8 cleaves with the cleavage product assisting in promoting cytochrome c release. This substrate was revealed by several groups to be the proapoptotic Bcl-2 protein family member, Bid.^{56, 64}

Bid is a 195 residue, 22 kDa protein that lacks the hydrophobic COOH-terminal domain, which confers a largely cytosolic localization.¹²¹ Bid interacts with Bcl-2, Bcl-x_L, and Bax by way of its BH3 domain and can annul the cytoprotective effects of Bcl-2 and Bcl-x_L.⁶⁴ The Bid amino acid sequence contains a putative caspase-8 cleavage site (57-L-Q-T-G-61) within its NH₂ terminus and Bid is indeed cleaved between residues 59 and 60 by caspase-8 *in vivo* and *in vitro*.^{56, 64} Following cleavage, the truncated Bid translocates to the mitochondria where it is a potent inducer of cytochrome c release, suggesting that the truncated Bid may play a role in increasing the permeability of the mitochondria membrane, allowing cytochrome c escape.

The three-dimensional structure of Bid shows a strong similarity to Bcl-x_L despite its modest sequence similarity to Bcl-x_L and other Bcl-2 family members. This structural similarity again implied that Bid might possess pore-forming capacity, and indeed BID does, but with a twist: Only the cleaved form of BID is able to form conductive channels *in vitro*.⁹⁵ The cleavage of Bid removes the amino-terminus, which results in an increased exposure of hydrophobic surface area, most notably of the central helix pair ($\alpha 4$ and $\alpha 5$) that are the putative pore-forming regions for Bid. This increase in exposed hydrophobic surface area may promote membrane insertion. Also, the cleaved form has

an increased accessibility of the BH3 domain that is involved in dimerization with other Bcl-2 family proteins,⁹¹ suggesting that the cleavage may promote protein-protein interactions that may modulate activity of other Bcl-2 family members involved in cytochrome c release, most notably, Bax.

Bid-null mice were generated and found to be resistant to treatment with anti-Fas antibody, which is normally lethal to wild-type mice.¹³⁰ In keeping with the two types of cell death for Fas-induced apoptosis, the Fas ligation in the Bid-null mice was reduced in hepatocytes, but the response of thymocytes and fibroblasts was largely unaffected.¹³⁰

Another study suggests that Bid works independently of caspase cleavage and in concert with Bax to promote cytochrome release. In this case, the BH3 domain of Bid would bind to the hydrophobic cleft of Bax, which would promote its translocation to the mitochondria and perhaps foster a membrane-inserted *active* state.²⁴ In this capacity, Bid acts more as a chaperone than a direct effector of cytochrome c release.

Thus, a role has emerged for Bid in at least some apoptotic pathways, in which it seems to function as a messenger for relaying signals induced by Fas or TNF at the cytoplasmic membrane, to the mitochondria. In other contexts, Bid may function more as a chaperone to promote Bax activation.

INHIBITOR OF APOPTOSIS PROTEIN FAMILY PROTEINS

Possibly one of the most ancient evolutionary pressures for a cell suicide program can be attributed to viruses. Death of the infected host cell stymies viral propagation—thereby protecting uninfected neighboring cells; however, viruses have co-evolved strategies for promoting cell survival by targeting conserved steps in the host cell-death program.^{13, 51, 134} Based on their antiapoptotic function during viral infection, inhibitor of apoptosis proteins (IAPs) were first discovered in baculovirus.^{9, 21} Intriguingly, ectopic expression of some Baculoviral IAPs in cultured mammalian cells suppressed cell death induced by several apoptotic stimulus, sug-

gesting that IAPs must block a conserved step in the cell-death program.

A novel 70 amino acid domain termed the *Baculoviral inhibitory repeat* (BIR) characterizes the IAP-protein family.^{9, 21} Based on sequence homology to the BIR domain, yeast, worms, flies, birds, mice, pigs, and humans all appear to encode IAP relatives, although many of these proteins have yet to be tested for antiapoptotic function. Six human IAP relatives have been identified including; NAIP, cIAP1/HIAP-2, cIAP2/HIAP-1, XIAP/hILP, Survivin, and BRUCE.^{2, 30, 40, 58, 87} NAIP, XIAP, cIAP1, cIAP2, and Survivin all exhibit antiapoptotic function in various experimental systems, thus confirming their membership in the IAP family of proteins.

Up to three tandem copies of the BIR domain can occur within the known human IAP-family proteins (Fig. 3). The repeated BIRs are highly conserved within a particular IAP and between distinct family members—ranging from approximately 30% to 90% identity. The conserved presence and spacing of cysteine and histidine residues present within the BIR domain coordinate zinc-binding.¹⁰⁸ Some IAPs also contain a zinc-binding ringfinger motif at their C-terminus. Although the role of the ringfinger motif with respect to IAP antiapoptotic function is unknown, in all cases at least one BIR is required for IAP-mediated suppression of cell death (reviewed in reference 25).

HOW DO INHIBITOR OF APOPTOSIS PROTEINS SUPPRESS APOPTOTIC PATHWAYS?

Recent studies have demonstrated that several of the human IAPs (XIAP, cIAP1, and cIAP2) directly inhibit caspases.^{26, 89} XIAP, cIAP1, and cIAP2 bind and potently inhibit caspase-3, to 7 and -9 but not caspase-1, 6, 8, or 10 or CED-3.^{79–81} Survivin also can be coimmunoprecipitated with caspase-3, -7, and -9, and it suppresses apoptosis induced by overexpression of these caspases, implying that Survivin also is a caspase inhibitor.¹¹³ More recently, IAPs from other species, including, *Drosophila*, *Lepidopteran*, and *Baculovirus*, have been shown to function by inhib-




Figure 3. Human IAP proteins. Highly conserved BIR domains that can be present at 1 to 3 copies (BIR 1–3) characterize IAPs. XIAP, cIAP1 and cIAP2, also contain a C-terminal ring domain (Ring). Survivin and BRUCE have a single BIR domain. In addition to a single N-terminal BIR domain, BRUCE also has ubiquitin conjugating (E2) domain (UBC) at its C-terminus; however, the functional significance of the BIR or UBC domain in BRUCE has yet to be reported.

iting specific caspases.* Thus, caspase inhibition appears to be a conserved mechanism by which IAP family proteins suppress cell death.

Caspase-8 and -9 represent the pinnacle caspases in the Fas/TNF-family death receptor and cytochrome c/Apaf-1 pathways, respectively (see Fig. 1). Although human IAPs (XIAP, cIAP1, and cIAP2) do not bind or inhibit caspase-8, they do bind to and inhibit its substrate, caspase-3, thereby arresting the proteolysis cascade and providing protection from Fas/caspase-8-induced apoptosis.^{26–27, 89} In contrast, in mitochondria-dependent pathways for caspase activation, XIAP, cIAP1, and cIAP2 directly bind to the apical caspase, procaspase-9, and prevent its processing and activation induced by cytochrome c, and Apaf-1.²⁷ Presumably, IAP interaction with procaspase-9 occurs on recruitment to the apoptosome complex, but this has yet to be determined.

Overexpression of IAP-family proteins has been shown to suppress apoptosis induced by Bax and other proapoptotic Bcl-2 family proteins, which are known for their ability to target mitochondria and induce cytochrome c release.^{10, 26, 48, 66, 125} The IAPs, however, do not interfere with Bax-mediated release of cytochrome c,^{48, 34} an observation that is consistent with data indicating that the

human IAPs (at least XIAP, cIAP1, cIAP2, and Survivin) block caspase activation and apoptosis downstream of Bax, Bik, Bak and cytochrome c.^{26–27, 31, 82, 89, 113}

At least for XIAP, the ability to inhibit caspase-3 and -7 has been attributed to its BIR2 domain and sequence just N-terminal to the BIR2 domain,^{108, 111} whereas the ability to inhibit caspase-9 localizes to the BIR3-ring region of XIAP.²⁸ Therefore, at least some IAPs have evolved distinct caspase inhibitory domains that may, in part, explain their versatility and effectiveness as antiapoptotic proteins.

IAPs and more specifically BIR domains, however, may have other functions. BIR-containing proteins have recently been identified in the yeast strains *Schizosaccharomyces pombe* and *Saccharomyces cerevisiae*. Because yeast do not appear to contain caspase-like proteases, yeast BIR proteins presumably have functions other than caspase inhibition. Consistent with this idea, yeast BIR proteins are reported to facilitate cell division.^{118, 131} Similarly, recent genetic analysis of a *C. elegans* BIR-containing gene demonstrated its essential role in cytokinesis, rather than apoptosis.³⁵

Interestingly, the single BIR-domain of the IAP family member Survivin, seems most closely related to the BIR domains found in yeast and worms, which as reviewed previously are reported to function in cell division and not in cell death. The scenario for

*References 26–28, 44, 49, 89, 108, 111, 113, 122

human Survivin, however, may not be as straight forward. Indeed, Survivin is expressed in the G₂/M phase of the cell cycle in a cycle-regulated manner.⁵⁵ At the beginning of mitosis, Survivin associates with microtubules and disruption of Survivin-microtubule interaction results in loss of Survivin's antiapoptotic function and increased caspase-3 activity. These and other results suggest that Survivin may countact a default induction of apoptosis at the G₂/M "checkpoint" of the cell cycle.⁵⁵ Thus, the human IAP Survivin appears to bridge the evolutionary gap between the nematode and yeast BIR proteins—which are regulators of cell division, and other viral, fly and human IAPs—that are antiapoptotic proteins.

INHIBITOR OF APOPTOSIS PROTEINS, SIGNAL TRANSDUCTION, AND APOPTOSIS

cIAP2 has been functionally implicated in TNF-induction of nuclear factor (NF- κ B) and protection from apoptosis.¹⁷ First, TNF- α has been shown to induce expression of cIAP2 though stimulation of NF- κ B. Second, overexpression of cIAP2, reportedly can also lead to NF- κ B activation. Third, cIAP2 expression suppresses cell death induced by TNF- α through the receptor TNFR-1. A dominant form of the NF- κ B inhibitor I- κ B (resistant to degradation), blocks these cIAP2 activities, implying that cIAP2 participates in a positive feedback mechanism regulating NF- κ B activation by targeting I- κ B for degradation (Fig. 4). Moreover, a mutant of cIAP2 lacking the C-terminal ring domain inhibited NF- κ B induction by TNF and enhanced TNF killing. Based on these findings, the authors¹⁷ suggested that cIAP2 is critically involved in TNF signaling events that induce NF- κ B, which are required for suppression of TNF-induced apoptosis.

Is the induction of IAP-family genes, however, critical for the antiapoptotic effect of NF- κ B? Studies of the effects of TNF- α on IAP-family gene expression in endothelial cells suggests the answer to this question may be difficult to obtain because of redundancy in IAP family genes. Transcription of cIAP1,




Figure 4. Antiapoptotic activity of NF- κ B. At least one proposed mechanism by which NF- κ B suppresses cell death is through its ability to induce the expression of IAPs, which in turn block apoptosis. IAPs also may facilitate the destruction of the NF- κ B inhibitor I- κ B, thereby potentiating a positive feedback loop that promotes cell survival.

cIAP2, and XIAP genes was found to be strongly up-regulated on treatment of endothelial cells with the TNF- α , interleukin (IL)-1 β , and LPS—reagents that lead to NF- κ B activation.¹⁰² In these studies, overexpression of I- κ B suppressed NF- κ B activation and prevented the induction of all these IAP-family genes. I- κ B overexpression also sensitized endothelial cells to TNF- α -induced apoptosis. Ectopic expression of at least one of the IAPs, XIAP, suppressed the I- κ B effect, thereby protecting endothelial cells from TNF- α -induced apoptosis, suggesting that XIAP represents one of the NF- κ B-regulated genes that can counteract the apoptotic signals caused by TNF- α -induced activation of caspase-8.¹⁰² Thus, although we do not know whether IAP expression is necessary for NF- κ B-mediated protection against TNF- α , it is sufficient. Based on these and similar reports, it may be worth considering whether dysfunctional regulation of the IAPs occurs in sepsis and some inflammatory conditions, where cytokine-induced endothelial cell death occurs.

INHIBITOR OF APOPTOSIS PROTEIN AND Bcl-2 FAMILY PROTEINS IN DISEASE

Misregulation of the balance between life and death at the cellular level, can contribute

to acute and chronic disease. Resistance to cell-death stimuli can result in an expanded population of diseased cells, as in the case of some carcinomas, and may play a role in angiogenesis and cardiovascular-related diseases. Excessive cell death, however, can contribute to autoimmune and neurodegenerative diseases and acute conditions, such as ischemia and excessive tissue damage following trauma. Therefore, it is perhaps not surprising that dysregulation of Bcl-2 and IAP family proteins is increasingly implicated in the pathology of human diseases.

HEART AND VASCULAR RELATED DISEASES

Nuclear factor κ B seems to play an important role in controlling the induction of apoptosis in human and rat vascular smooth muscle cells (SMCs).³³ Reportedly, SMCs in high-density culture are resistant to apoptosis, which correlated with the expression of cIAP1 and high NF- κ B activity. Transfection of I κ -B, inhibitor of NF- κ B, reduced human cIAP1 mRNA levels. These data suggest that NF- κ B activity increases expression of cIAP1, which confers protection from cell death. Consistent with this idea, antisense inhibition of IAP-1 sensitized high-density SMCs to cell-death induction.³³ Based on their data, the author³³ suggested that cIAP1 is transcriptionally regulated by NF- κ B and that SMCs at high density are protected by an antiapoptotic mechanism that involves increased expression of NF- κ B and cIAP1.

Using differential display, cIAP2 was reportedly one of the cytokine-responsive genes from endothelial cells that can be regulated by monocyte conditioned medium or TNF- α . Furthermore, in vivo expression of cIAP2 was detected in endothelial cells overlying lesions heavily infiltrated by monocytes and foam cells. These results suggest that cIAP2 may play an important role in the molecular processes involved in vascular diseases, such as atherosclerosis.⁴³

Several studies have detected the presence of Bcl-2 protein family members in cardiac myocytes. In rat heart, antiapoptotic Bcl-2 and Bcl-x_L were expressed to high levels in neona-

tal cardiac tissue and their presence was maintained throughout development. The proapoptotic proteins Bad and Bax, while present at high levels in neonatal hearts, were absent in adult hearts.¹⁹ Although the functional significance of these observations remains to be investigated, the presence of these proteins may suggest that they play roles developing, modeling and maintaining the adult heart by regulating apoptosis.

In this regard, reperfusion of ischemic myocardium causes cardiomyocyte apoptosis that reportedly occurs in concert with down-regulation of Bcl-2 gene expression.⁶⁹ In these studies, ischemic preconditioning (PC) mediated by cyclic episodes of short-term ischemia and reperfusion, reportedly reduced apoptotic cell death. PC was shown to initiate a signaling pathway by potentiating tyrosine kinase phosphorylation, which lead to the activation of p38 MAP kinase and MAPKAP kinase 2. Based on observations that NF- κ B plays a crucial role in this signaling pathway and can be a target of oxygen free radicals and that Bcl-2 is reported to be an antioxidant gene, the authors⁶⁹ hypothesized that reactive oxygen species might play a role in this signaling process. Alternatively, NF- κ B may influence the expression of other antiapoptotic proteins, such as the IAPs, thereby conferring protection against ischemic insult in cardiomyocytes.

Expression of p53 in ventricular myocytes was shown to result in a significant increase in Bax and was sufficient to trigger apoptosis.⁵⁰ In these studies, expression of Bcl-2 was sufficient to prevent p53-mediated apoptosis and p53-dependent transcription of Bax in ventricular myocytes.⁵⁰ These studies suggest that pro- and antiapoptotic proteins can influence ventricular remodeling after injury. This may have clinical significance since inappropriate loss of myocardial cells has been suggested to contribute to conduction defects and heart defects.

NEURONAL AND NEURODEGENERATIVE DISEASES

The *NAIP* gene was first identified because of its apparent deletion in patients with spinal

muscular atrophy (SMA), a hereditary motor neuron degenerative disease.^{58, 88} Though the primary genetic defect in SMA has been ascribed to an adjacent gene,⁶³ *SMN*, rather than *NAIP*, patients with the severest forms of this disease appear to harbor deletions at 5q13.1 that encompass the *SMN* and *NAIP* genes. Intriguingly, the survival motor neuron gene (*SMN*) protein has been reported to bind Bcl-2 and enhance Bcl-2-mediated protection from apoptosis,⁴⁵ raising the possibility that two survival genes may be lost in more severely affected individuals.

Consistent with the primary defect in SMA being attributed to the *SMN* gene, it recently was reported that *NAIP*-deleted mice develop normally. The survival of pyramidal neurons in the hippocampus after kainic acid-induced limbic seizures is, nevertheless, greatly reduced in the *NAIP* knock-out animals. The authors⁴² concluded that although *NAIP* is not necessary for normal development of the murine central nervous system; it is required for neuronal survival in pathological conditions.

NAIP also may be involved in adaptive responses to ischemia. Transient forebrain ischemia selectively elevates levels of *NAIP* in rat neurons that are resistant to ischemia-reperfusion.¹²⁷ Up-regulation of endogenous *NAIP* expression or intracerebral injection of *NAIP*-encoding adenoviruses reportedly reduces ischemic damage in the rat hippocampus, suggesting that *NAIP* may play a role in conferring resistance to ischemia-induced cell death.¹²⁷ In cell culture experiments, however, transfection of primary cerebellar granule cell neurons with adenoviruses encoding *NAIP*, *XIAP*, *cIAP1*, or *cIAP2* delayed but did not prevent apoptosis induced by K^+ depolarization and serum deprivation. Nonapoptotic cell death induced by L-glutamate was unaffected by these IAP-family proteins.¹⁰¹ Thus, IAPs are apparently insufficient to protect some types of neurons from insults often associated with ischemia.

Nevertheless, it was reported that adenovirus-mediated overexpression of *XIAP* prevented the production of catalytically active caspase-3 and degeneration of CA1 neurons after transient forebrain ischemia.¹²⁶ CA1 neurons protected in this manner appeared to

function normally, as assessed by immunohistochemical detection of the neuronal activity marker nerve growth factor inducible-A and by spatial learning performance in the Morris water maze. The authors¹²⁶ concluded that *XIAP* overexpression permits CA1 neurons to survive and operate properly after an ischemic insult.

CANCER

As described here and reviewed previously, the gene encoding Bcl-2 protein was first discovered because of its involvement in the t(14;18) chromosomal translocations commonly found in B-cell lymphomas.¹¹⁷ Bcl-2 contributes to neoplastic cell expansion by preventing cell turnover because of programmed cell death. In this regard, overexpression of antiapoptotic Bcl-2 and Bcl-x_L also has been documented in many other types of human tumors, including cancers of the prostate, colon, breast, and lung. Moreover, antiapoptotic Bcl-2 family proteins have been associated with chemoresistance and radioreistance in some types of malignancy (reviewed in references 46, 83, 85, 107).

The IAPs also are implicated increasingly in the oncogenic process. For example, the oncoprotein v-Rel, a member of the Rel/NF- κ B family of transcription factors, induces malignant transformation and inhibits apoptosis. The chicken homolog of cIAP1 (ch-IAP1) was found to be up-regulated following expression of v-Rel in fibroblasts, a B-cell line, and in spleen cells.¹³² Expression of exogenous ch-IAP1 in temperature-sensitive v-Rel-transformed spleen cells inhibited apoptosis in these cells at the nonpermissive temperature. Based on these results, it appears that ch-IAP1 is induced and functions as a suppressor of apoptosis in the v-Rel-mediated transformation process.¹³²

cIAP2 and a novel gene, named *MLT*, are recurrently rearranged in marginal zone cell lymphomas of mucosa-associated lymphoid tissue (MALT).²⁹ This t(11;18)(q21;q21) rearrangement appears to be the key genetic lesion and is found in approximately 50% of cytogenetically abnormal low-grade MALT lymphomas. Based on these data, it was sug-

gested that the resulting cIAP2-MLT fusion may have enhanced anti-apoptotic function, thereby contributing to the oncogenesis of MALT lymphoma.²⁹

The IAP member Survivin is expressed in a high proportion of the commonest human cancers but not in normal terminally differentiated adult tissues.² The assessment of Survivin expression in human tumor specimens included in situ RNA hybridization and immunohistochemical analysis, confirming expression in tumor cells but not admixed stromal cells or adjacent normal tissues.³ Thus, altered Survivin expression seems to define a common event associated with the pathogenesis of most human cancers. Moreover, reductions in Survivin expression achieved using antisense strategies cause apoptosis and sensitization to anticancer drugs, at least in some tumor cell lines, implying that Survivin expression can be important for cell survival or chemoresistance in carcinomas.³

Not all tumors, however, express Survivin and even within a given type of cancer, heterogeneity in Survivin expression may be observed. Immunohistochemical assessments of Survivin expression in tumors in which either immunointensity, percentage immunopositivity, or have been measured to segregate Survivin negative from positive (Survivin low from high) tumors suggest that Survivin expression (or higher levels of Survivin expression) is associated a poor prognosis in neuroblastomas, colon, and gastric cancers.^{1-3, 62} Though preliminary, assessments of Survivin expression may be of prognostic significance for patients with some types of cancer.

In this regard, a recent study revealed that Survivin expression was positive in 118 of 167 breast carcinoma cases (70.7%) having histological stages I to IH.¹¹⁴ In contrast, no Survivin expression was detected in adjacent normal tissue. Survivin-positive samples strongly correlate with Bcl-2 expression and exhibited reduced apoptosis (low apoptotic index). Patients with a low apoptotic index had lower survival rates than the group having a high apoptotic index. The authors¹¹⁴ suggest that apoptosis inhibition by Survivin alone, or in cooperation with Bcl-2, is a sig-

nificant prognostic parameter of worse outcome in breast carcinoma.

Endothelial cell activation and dysfunction can play a prominent role in physiological processes, such as angiogenesis, and in the pathophysiology of atherosclerosis.¹¹⁶ Vascular endothelial growth factor (VEGF) is a potent angiogenic factor that can act as an endothelial cell mitogen and seems to be a major survival agent for endothelial cells during angiogenesis and vasculogenesis. VEGF has been shown to mediate this latter function, in part through the induction of Bcl-2 expression and the activation of the P13 kinase-Akt/PKB signaling pathway.¹¹⁶ Additionally, VEGF was reported to increase XIAP and Survivin protein levels 2.9- and 19.1-fold, respectively, in human umbilical vein endothelial cells, suggesting that VEGF mediated survival may be, in part, mediated by inducing expression of these IAPs. The authors¹¹⁶ suggest that these results raise the possibility of therapeutically targeting XIAP or Survivin in antiangiogenic therapy as a means of suppressing tumor growth, in addition to directly targeting tumor cells that express these survival proteins.

Consistent with the above observations, a separate study reported that stimulation of quiescent endothelial cells with mitogens, including VEGF and basic fibroblast growth factor (bFGF), increased Survivin expression approximately 16-fold.⁷⁸ Survivin protein concentration was minimal in the endothelium of nonproliferating capillaries of normal skin, whereas it became massively up-regulated in newly formed blood vessels of granulation tissue in vivo. Ectopic expression of Survivin reduced caspase-3 activity and counteracted apoptosis induced by TNF- α /cycloheximide in endothelial cells⁷⁸ suggesting that anti-apoptotic proteins may play an important role in the angiogenic process.

IMMUNE DISEASE

As outlined above, increased activity or expression of antiapoptotic proteins can adversely influence the maintenance of healthy cells by suppressing apoptosis. In contrast, lack of antiapoptotic protein function can

result in excessive apoptosis. A recent example of this concept was described for cartilage-hair hypoplasia (CHH) syndrome—a rare autosomal recessive disease characterized by increased T-cell apoptosis and cell-mediated or combined immunodeficiency.¹²⁸ This study reported that CHH was associated with altered expression of Fas, Fas ligand (FasL), IAP, Bax, and Bcl-2. Increased apoptosis in CHH correlated with increased expression of Fas, FasL, and Bax and decreased expression of Bcl-2 and IAPs compared with the control. These data suggest that increased apoptosis of T cells contributes to lymphopenia and immunodeficiency in CHH, and that increased T-cell death, in this case, is mediated by altered expression of pro- and antiapoptotic proteins.¹²⁸

Changes in Fas, FasL, and Bcl-2 expression have also been reported in circulating T cells in patients with HIV infection,^{6, 23, 26} further suggesting a problem with regulation of apoptosis genes in immunodeficiency states. Conversely, autoimmune disorders are commonly characterized by a failure to remove autoreactive lymphocytes. In this context, studies of transgenic and knock-out mice have provided examples of autoimmunity that is caused by changes in the expression of Bcl-2, Bcl-x_L, and Fas.^{61, 65, 79, 112, 123} Alterations in the expression or function of apoptosis-regulating genes, such as Bcl-2 and Fas, also have been described in humans with lupus or other autoimmune disorders.^{7, 15, 80–81, 115} Also, the HIV protease reportedly cleaves Bcl-2.¹⁰⁶ Further, the HIV tat protein can sensitize T cells to Fas-dependent apoptosis.¹²⁴ Thus, defects in apoptosis regulation are intricately associated with immune system diseases.

SUMMARY

The balance between pro- and antiapoptotic proteins can determine cellular fate. In this regard, the Bcl-2 and IAP protein families have evolved as highly conserved regulators of cell death. A further testament to their critical roles in maintaining balance between cell life and death may be the increasing implication of Bcl-2 and IAP proteins in the

pathologies of human diseases. Although much has been learned about these families of proteins, future studies of the Bcl-2 and IAP families are sure to hold more exciting discoveries and will continue to reveal new strategies for combating human diseases.

References

1. Adida C, Crotty PL, McGrath J, et al: Developmentally regulated expression of the novel cancer anti-apoptosis gene survivin in human and mouse differentiation. *Am J Pathol* 152: 43–49, 1998
2. Ambrosini G, Adida C, Altieri D: A novel anti-apoptosis gene, survivin, expressed in cancer and lymphoma. *Nat Med* 3: 917–921, 1997
3. Ambrosini G, Adida C, Sirugo G, et al: Induction of apoptosis and inhibition of cell proliferation by survivin gene targeting. 273, 11177–11182, 1998
4. Antonsson B, Conti F, Ciavatta A, et al: Inhibition of Bax channel-forming activity by Bcl-2. *Science* 277: 370–372, 1997
5. Antonsson B, Montessuit S, Lauper S, et al: Bax oligomerization is required for channel-forming activity in liposomes and to trigger cytochrome c release from mitochondria. *Biochem J* 345: 271–278, 2000
6. Aries SP, Schaaf B, Muller C, et al: Fas (CD95) expression on CD4⁺ T cells from HIV-infected patients increases with disease progression. *Journal of Molecular Medicine* 73: 591–593, 1995
7. Aringer M, Wintersberger W, Steiner CW, et al: High levels of bcl-2 protein in circulating T lymphocytes, but not B lymphocytes, of patients with systemic lupus erythematosus. *Arthritis Rheumatoid* 37:423–430, 1994
8. Bernardi P, Scorrano L, Colonna R, et al: Mitochondria and cell death: Mechanistic aspects and methodological issues. *Eur J Biochem* 264: 687–701, 1999
9. Birnbaum MJ, Clem RJ, Miller LK: An apoptosis-inhibiting gene from a nuclear polyhedrosis virus encoding a polypeptide with Cys/His sequence motifs. *J Virol* 68: 2521–2528, 1994
10. Bossy-Wetzel E, Newmeyer DD, Green DR: Mitochondrial cytochrome c release in apoptosis occurs upstream of DEVD-specific caspase activation and independently of mitochondrial transmembrane depolarization. *EMBO J* 17: 37–49, 1998
11. Brenner C, Herve C, Helena LA, et al: Bcl-2 and Bax regulate the channel activity of the mitochondrial adenine nucleotide translocator. *Oncogene* 19, 329–336, 2000
12. Bullock JO, Cohen FS, Dankert JR, et al: Comparison of the macroscopic and single channel conductance properties of colicin E1 and its COOH-terminal tryptic peptide. *J Biol Chem* 258: 9908–9912, 1983
13. Bump NJ, Hackett M, Hugunin M, et al: Inhibition of ICE family proteases by baculovirus anti-apoptotic protein p35. *Science* 269: 1885–1888, 1995
14. Chang BS, Minn AJ, Muchmore SW, et al: Identification of a novel regulatory domain in Bcl-x_L and Bcl-2. *EMBO J* 16: 968–977, 1997
15. Cheng J, Liu C, Yang P, et al: Increased lymphocyte

- apoptosis in Fas ligand transgenic mice. *J Immunol* 159: 674–684, 1997
16. Chou JJ, Li H, Salvesen GS, et al: Solution structure of BID, an intracellular amplifier of apoptotic signaling. *Cell* 96: 615–624, 1999
 17. Chu ZL, McKinsey TA, Liu L, et al: Suppression of tumor necrosis factor-induced cell death by inhibitor of apoptosis c-IAP2 is under NF- κ B control. *Proc Natl Acad Sci USA* 94: 10057–10062, 1997
 18. González-García M, Pérez-Ballesteros R, Ding L, et al: Bcl-xL is the major bcl-x mRNA form expressed during murine development and its product localizes to mitochondria. *Development* 120: 3033–3042, 1994
 19. Cook SA, Sugden PH, Clerk A: Regulation of bcl-2 family proteins during development and in response to oxidative stress in cardiac myocytes: association with changes in mitochondrial membrane potential. *Circ Res* 5: 940–949, 1999
 20. Cramer WA, Heymann JB, Schendel SL, et al: Structure-function of the channel-forming colicins. *Annu Rev Biophys Biomol Struct* 24: 611–641, 1995
 21. Crook NE, Clem RJ, Miller LK: A apoptosis-inhibiting baculovirus gene with a zinc finger-like motif. *J Virol* 67: 2168–2174, 1993
 22. de Jong D, Prins FA, Mason DY, et al: Subcellular localization of the bcl-2 protein in malignant and normal lymphoid cells. *Cancer Res* 54: 256–260, 1994
 23. De Rossi A, Ometto L, Rancella S, et al: HIV-1 induces down-regulation of bcl-2 expression and death by apoptosis of EBV-immortalized B cells: A model for a persistent “self-limiting” HIV-1 infection. *Virology* 198: 234–244, 1994
 24. Desagher S, Osen-Sand A, Nichols A, et al: Bid-induced conformational change of Bax is responsible for mitochondrial cytochrome c release during apoptosis. *J Cell Biol* 155: 891–901, 1999
 25. Deveraux QL, Reed JC: IAP-family of proteins: suppressors of cell death. *Genes and Development* 13: 239–252, 1999
 26. Deveraux QL, Takahashi R, Salvesen GS, et al: X-linked IAP is a direct inhibitor of cell death proteases. *Nature* 388: 300–303, 1997
 27. Deveraux QL, Roy N, Stennicke HR, et al: IAPs block apoptotic events induced by caspase-8 and cytochrome c by direct inhibition of distinct caspases. *EMBO J* 17: 2215–2223, 1998
 28. Deveraux QL, Leo E, Stennicke HR, et al: Cleavage of human inhibitor of apoptosis protein XIAP results in fragments with distinct specificities for caspases. *EMBO* 18: 5242–5251, 1999
 29. Dierlamm J, Baens M, Wlodarska, et al: The apoptosis inhibitor gene API2 and a novel 18q gene, MLT, are recurrently rearranged in the t(11;18)(q21;q21) p53-associated with mucosa-associated lymphoid tissue lymphomas. *Blood* 93: 3601–3609, 1999
 30. Duckett CS, Li F, Tomaselli KJ, et al: A conserved family of cellular genes related to the baculovirus IAP gene and encoding apoptosis inhibitors. *EMBO J* 15: 2685–2689, 1996
 31. Duckett CS, Li F, Tomaselli KJ, et al: Human IAP-like protein regulates programmed cell death downstream of Bcl-X_L and cytochrome c. *Mol Cell Biol* 18: 608–615, 1998
 32. Duke RC, Ojcius DM, Young JD: Cell suicide in health and disease. *Scientific American* 275:80–87, 1996
 33. Erl W, Hansson GK, de Martin R, et al: Nuclear factor- κ B regulates induction of apoptosis and inhibitor of apoptosis protein-1 expression in vascular smooth muscle cells. *Circ Res* 84: 668–677, 1999
 34. Finucane DM, Bossy-Wetzel E, Cotter TG, Green DR: Bax-induced caspase activation and apoptosis via cytochrome c release from mitochondria is inhibitable by Bcl-XL. *J Biol Chem* 274: 2225–2233, 1999
 35. Fraser AG, James C, Evan GI, et al: *Caenorhabditis elegans* inhibitor of apoptosis protein (IAP) homologue BIR-1 plays a conserved role in cytokinesis. *Curr Biol* 9: 292–301, 1999
 36. Gougeon ML: Programmed cell death in HIV infection: Dysregulation of bcl-2 and fas pathways and contribution to AIDS pathogenesis. *Psychoneuroendocrinology* 22:7,1997
 37. Green D, Reed J: Mitochondria and apoptosis. *Science* 281: 1309–1312, 1998
 38. Gross A, McDonnell JM, Korsmeyer SJ: BCL-2 family members and the mitochondria in apoptosis. *Genes Dev* 13: 1899–1911, 1999
 39. Gross A, Jockel J, Mei MC, et al: Enforced dimerization of Bax results in its translocation, mitochondrial dysfunction and apoptosis. *EMBO J* 17: 3878–3885, 1998
 40. Hauser HP, Bardroff M, Pyrowolakis G, et al: A giant ubiquitin-conjugating enzyme related to IAP apoptosis inhibitors. *J Cell Biol* 141: 1415–1422, 1998
 41. Hirotsu M, Zhang Y, Fujita N, et al: NH₂-terminal BH4 domain of Bcl-2 is functional for heterodimerization with Bax and inhibition of apoptosis. *J Biol Chem* 274: 20415–20420, 1999
 42. Holcik M, Thompson CS, Yarghi Z, et al: The hippocampal neurons of neuronal apoptosis inhibitory protein 1 (NAIP1)-deleted mice display increased vulnerability to kainic acid-induced injury. *Proc Natl Acad Sci* 97:2286–2290, 2000
 43. Horrevoets AJ, Fantijn RD, van Zonneveld AJ, et al: Vascular endothelial genes that are responsive to tumor necrosis factor- α in vitro are expressed in atherosclerotic lesions, including inhibitor of apoptosis protein-1, stannin, and two novel genes. *Blood* 93: 3418–3431, 1999
 44. Huang Q, Deveraux QL, Maeda S, et al: Evolutionary conservation of apoptosis mechanisms: Lepidopteran and baculoviral inhibitor of apoptosis proteins are inhibitors of mammalian caspase-9. *Proc Natl Acad Sci USA* 97: 1427–1432, 2000
 45. Iwahashi H, Eguchi Y, Yasuhara N, et al: Synergistic anti-apoptotic activity between bcl-2 and smn implicated in spinal muscular atrophy. *Nature* 390: 413–417, 1997
 46. Jaattela M, Xie Z, Deveraux QL: Escaping cell death: Survival proteins in cancer. *Exp Cell Res* 248: 30–43, (1999)
 47. Joo P, Kuo CJ, Yuan J, et al: Essential requirement for caspase-8/FLICE in the initiation of the Fas-induced apoptotic cascade. *Curr Biol* 8: 1001–1008, 1998
 48. Jürgensmeier JM, Xie Z, Deveraux QL, et al: Bax directly induces release of cytochrome c from isolated mitochondria. *Proc Natl Acad Sci USA* 95: 4997–5002, 1998
 49. Kaiser WJ, Vucic D, Miller LK: The Drosophila inhibitor of apoptosis D-IAP1 suppresses cell death

- induced by the caspase drICE. *FEBS Lett* 440: 243–248, 1998
50. Kirshenbaum LA: Regulators of apoptosis in the heart: A matter of life and death. *Can J Cardiol* 14: 457–460, 1998
 51. Komiya T, Ray CA, Pickup DJ, et al: Inhibition of interleukin-1 beta converting enzyme by the cowpox virus serpin CrmA: An example of cross-class inhibition. *J Biol Chem* 269: 19331–19337, 1994
 52. Korsmeyer SJ, Shutter JR, Veis DJ, et al: Bcl-2/Bax: A rheostat that regulates an anti-oxidant pathway and cell death. *Canc Biol* 4: 327–332, 1993
 53. Krajewski S, Krajewski M, Ellerby LM, et al: Release of caspase-9 from mitochondria during neuronal apoptosis and cerebral ischemia. *Proc Natl Acad Sci USA* 96: 5752–5757, 1999
 54. Krajewski S, Tanaka S, Takayama S, et al: Investigation of the subcellular distribution of the bcl-2 oncoprotein: Residence in the nuclear envelope, endoplasmic reticulum, and outer mitochondrial membranes. *Cancer Res* 53: 4701–4714, 1993
 55. Li F, Ambrosini G, Chu EY, et al: Control of apoptosis and mitotic spindle checkpoint by survivin. *Nature* 396: 580–584, 1998
 56. Li H, Zhu H, Xu CJ, et al: Cleavage of BID by caspase 8 mediates the mitochondrial damage in the Fas pathway of apoptosis. *Cell* 94: 491–501, 1998
 57. Li P, Nijhawan D, Budihardjo I, et al: Cytochrome c and dATP-dependent formation of Apaf-1/Caspase-9 complex initiates an apoptotic protease cascade. *Cell* 91: 479–489, 1997
 58. Liston P, Roy N, Tamai K, et al: Suppression of apoptosis in mammalian cells by NAIP and a related family of IAP genes. *Nature* 379: 349–353, 1996
 59. Liu X, Kim CN, Yang J, et al: Induction of apoptotic program in cell-free extracts: Requirement for dATP and Cytochrome C. *Cell* 86: 147–157, 1996
 60. London E: Diphtheria toxin: Membrane interaction and membrane translocation. *Biochim Biophys Acta* 1113: 25–51, 1992
 61. Lopez-Hoyos M, Carrio R, Menno R, et al: Constitutive expression of Bcl-2 in B cells causes a lethal form of lupus-like autoimmune disease after induction of neonatal tolerance to H-2^b alloantigens. *J Exp Med* 183: 2523–2531, 1996
 62. Lu CD, Altieri DC, Tanigawa N: Expression of a novel anti-apoptosis gene, survivin, correlated with tumor cell apoptosis and p53 accumulation in gastric carcinomas. *Cancer Res* 58: 1808–1812, 1998.
 63. Liu Q, Fischer U, Wang F, et al: The spinal muscular atrophy disease gene product, SMN, and its associated protein SIP1 are in a complex with spliceosomal snRNP proteins. *Cell* 90: 1013–1021, 1997
 64. Luo X, Budihardjo I, Zou H, et al: Bid, a Bcl2 interacting protein, mediates cytochrome c release from mitochondria in response to activation of cell surface death receptors. *Cell* 94: 481–490, 1998
 65. Ma A, Pena JC, Chang B, et al: Bcl-x regulates the survival of double-positive thymocytes. *Proc Natl Acad Sci U S A* 92: 4763–4767, 1995
 66. Mahajan NP, Linder K, Berry G, et al: Bcl-2 and Bax interactions in mitochondria probed with green fluorescent protein and fluorescence resonance energy transfer. *Nature-Biotechnology* 16: 547–552, 1998
 67. Martin D, Siegel R, Zheng L, et al: Membrane oligomerization and cleavage activates the caspase-8 (FLICE/MACH α 1) death signal. *J Biol Chem* 273: 4345–4349, 1998
 68. Matsuyama S, Xu Q, Velours J, et al: Mitochondrial FOF1-ATPase proton-pump is required for function of pro-apoptotic protein bax in yeast and mammalian cells. *Mol Cell* 1: 327–336, 1998
 69. Maulik N, Engelmann RM, Rousou JA, et al: Ischemic preconditioning reduces apoptosis by upregulating anti-death gene Bcl-2. *Circulation* 100:11369–11375, 1999
 70. McDonnell JM, Fushman D, Milliman CL, et al: Solution structure of the proapoptotic molecule BID: A structural basis for apoptotic agonists and antagonists. *Cell* 96: 625–634, 1999
 71. Minn AJ, Kettlun CS, Liang H, et al: Bcl-xL regulates apoptosis by heterodimerization-dependent and -independent mechanisms. *EMBO* 18: 632–643, 1999
 72. Minn AJ, Velez P, Schendel SL, et al: Bcl-x_L forms an ion channel in synthetic lipid membranes. *Nature* 385: 353–357, 1997
 73. Montal M, Mueller P: Formation of bimolecular membranes from lipid monolayers and a study of their electrical properties. *Proc Natl Acad Sci USA* 69: 3561–3566, 1972
 74. Muchmore SW, Sattler M, Liang H, et al: X-ray and NMR structure of human Bcl-XL, an inhibitor of programmed cell death. *Nature* 381: 335–341, 1996
 75. Nagata S: Apoptosis by death factor. *Cell* 88: 355–365, 1997
 76. Narula J, Pandey P, Arbustini E, et al: Apoptosis in heart failure: Release of cytochrome c from mitochondria and activation of caspase-3 in human cardiomyopathy. *Proc Natl Acad Sci USA* 96: 8144–8149, 1999
 77. Nouraini S, Six E, Matsuyama S, et al: The putative pore-forming domain of bax regulates mitochondrial localization and interaction with bcl-X. *Mol Cell Biol* 20: 1604–1615, 2000
 78. O'Connor DS, Schechner JS, Adida C, et al: Control of apoptosis during angiogenesis by survivin expression in endothelial cells. *Am J Pathol* 156: 393–398, 2000 (in press)
 79. O'Connor L, Strasser A, O'Reilly LA, et al: Bim: A novel member of the bcl-2 family that promotes apoptosis. *EMBO J* 17: 384–395, 1998
 80. Ohasako S, Hara M, Harigai M, et al: The Bcl-2/Bax ratio of lymphocytes from human systemic lupus erythematosus patients. *Jpn J Rheumatol* 7: 305–313, 1997
 81. Ohasako S, Hara M, Harigai M, et al: Expression and function of Fas antigen and bcl-2 in human systemic lupus erythematosus lymphocytes. *Clin Immunol Immunopathol* 73: 109–114, 1994
 82. Orth K, Dixit VM: Bik and Bak induce apoptosis downstream of CrmA but upstream of inhibitor of apoptosis. *J Biol Chem* 272: 8841–8844, 1997
 83. Pellegrini M, Strasser A: A portrait of the Bcl-2 protein family: Life, death, and the whole picture. *J Clin Immunol* 19: 365–377, 1999
 84. Reed J: Chronic lymphocytic leukemia: A disease of dysregulated programmed cell death. *Clinical Immunology Newsletter* 17:125–140, 1998
 85. Reed JC: Double identity for proteins of the Bcl-2 family. *Nature* 387: 773–776, 1997
 86. Reed JC: Cytochrome C: Can't live with it; Can't live without it. *Cell* 91: 559–562, 1997
 87. Rothe M, Pan M-G, Henzel WJ, et al: The TNFR2-TRAF signaling complex contains two novel pro-

- teins related to baculoviral inhibitor of apoptosis proteins. *Cell* 83: 1243–1252, 1995
88. Roy N, Mahadevan QL, McLean M, et al: The gene for neuronal apoptosis inhibitory protein is partially deleted in individuals with spinal muscular atrophy. *Cell* 80: 167–178, 1995
 89. Roy N, Deveraux QL, Takahashi R, et al: The c-IAP-1 and c-IAP-2 proteins are direct inhibitors of specific caspases. *EMBO J* 16: 6914–6925, 1997
 90. Salvesen GS, Dixit VM: Caspases: Intracellular signaling by proteolysis. *Cell* 91: 443–446, 1997
 91. Sattler M, Liang H, Nettesheim D, et al: Structure of Bcl-xL-Bak peptide complex: Recognition between regulators of apoptosis. *Science* 275: 983–986, 1997
 92. Scaffidi C, Fulch S, Srinivasan A, et al: Two CD95 (APO-1/Fas) signaling pathways. *EMBO J* 17: 1675–1687, 1998
 93. Schendel S, Montal M, Reed JC: Bcl-2 Family proteins as ion-channels. *Cell Death Different* 5: 372–380, 1998
 94. Schendel SL, Xie Z, Montal MO, et al: Channel formation by anti-apoptotic protein Bcl-2. *Proc Natl Acad Sci USA* 94: 5113–5118, 1997
 95. Schendel SL, Azimov R, Pawlowski K, et al: Ion channel activity of the BH3 only Bcl-2 family member, BID. *J Biol Chem* 274: 21932–21936, 1999
 96. Schendel SL, Click EM, Webster RE, et al: The TolA protein interacts with colicin E1 differently than with other group A colicins. *J Bacteriol* 197: 3683–3690, 1997
 97. Schlesinger P, Gross A, Yin X-M, et al: Comparison of the ion channel characteristics of proapoptotic BAX and anti-apoptotic BCL-2. *Proc Natl Acad Sci USA* 94: 11357–11362, 1997
 98. Schulke N, Sepuri NB, Paine D: In vivo zippering of inner and outer mitochondrial membranes by a stable translocation intermediate. *Proc Natl Acad Sci USA* 94: 7314–7319, 1997
 99. Sedlak TW, Oltvai ZN, Yang E, et al: Multiple Bcl-2 family members demonstrate selective dimerizations with Bax. *Proc Natl Acad Sci USA* 92: 7834–7838, 1995
 100. Shimizu S, Narita M, Tsujimoto Y: Bcl-2 family proteins regulate the release of apoptogenic cytochrome c by the mitochondrial channel. *Nature* 399: 483–487, 1999
 101. Simons M, Beinroth S, Gleichmann M, et al: Adenovirus-mediated gene transfer of inhibitors of apoptosis proteins delays apoptosis in cerebellar granule neurons. *Neurochemistry* 72: 292–301, 1999
 102. Stehlik C, Martin RD, Kumabashiri I, et al: NF- κ B regulated x-chromosome-linked IAP gene expression protects endothelial cells from TNF- α induced apoptosis. *J Exp Med* 188: 211–216, 1998
 103. Steller H: Mechanisms and genes of cellular suicide. *Science* 267:1445–1449, 1995
 104. Stennicke HR, Jürgensmeier JM, Shin H, et al: Procaspase-3 is a major physiologic target of caspase-8. *J Biol Chem* 273: 27084–27090, 1998
 105. Stennicke HR, Salvesen GS: Properties of the caspases. *Biochem Biophys Acta* 1387: 17–31, 1998
 106. Strack PR, Frey MW, Rizzo CJ, et al: Apoptosis mediated by HIV protease is preceded by cleavage of BCL-2. *Proc Natl Acad Sci U S A* 93: 9571–9576, 1996
 107. Strasser A, Huang DCS, Vaux DL: The role of the *Bcl-2/ced-9* gene family in cancer and general implications of defects in cell death control for tumorigenesis and resistance to chemotherapy. *Biochim Biophys Acta* 1333: F151–F178, 1997
 108. Sun C, Cal M, Gunasekera AH, et al: NMR structure and mutagenesis of the inhibitor-of-apoptosis protein XIAP. *Nature* 401: 818–822, 1999
 109. Susin SA, Lorenzo HK, Zamzami N, et al: Mitochondrial release of caspase-2 and -9 during the apoptotic process. *J Exp Med* 189: 381–394, 1999
 110. Susin SA, Lorenzo HK, Zamzami N, et al: Molecular characterization of mitochondrial apoptosis-inducing factor. *Nature* 397: 441–446, 1999
 111. Takahashi R, Deveraux QL, Tamm I, et al: A single BIR domain of XIAP sufficient for inhibiting caspases. *JBC* 273: 7787–7790, 1998
 112. Takahashi T, Tanaka M, Brannan CI, et al: Generalized lymphoproliferative disease in mice, caused by a point mutation in the Fas ligand. *Cell* 76: 969–976, 1994
 113. Tamm I, Wang Y, Sausville E, et al: IAP-family protein survivin inhibits caspase activity and apoptosis induced by Fas(CD95), Bax, and anticancer drugs. *Cancer Res* 59: 5315–5320, 1998
 114. Tanaka K, Iwamoto S, Gan G, et al: Expression of survivin and its relationship to loss of apoptosis in breast carcinomas. *Clin Cancer Res* 6: 127–134, 2000 (in press)
 115. Tokano Y, Miyake S, Kayagaki N, et al: Soluble Fas molecule in the serum of patients with systemic lupus erythematosus. *Journal of Clinical Immunology* 16: 261–265, 1996
 116. Tran J, Rak J, Sheehan C, et al: Marked induction of the IAP family anti-apoptotic proteins survivin and XIAP by VEGF in vascular endothelial cells. *Biochem Biophys Res Commun* 264: 781–788, 1999
 117. Tsujimoto Y, Yunis J, Onorato-Showe L, et al: Molecular cloning of the chromosomal breakpoint of B-cell lymphomas and leukemias with the t(11;14) chromosome translocation. *Science* 224: 1403–1406, 1984
 118. Uren AG, Vaux DL, Beilharz T, et al: Role for yeast inhibitor of apoptosis (IAP)-like proteins in cell division. *Proc Natl Acad Sci USA* 96: 10170–10175, 1999
 119. Vander Heiden MG, Thompson CB: Bcl-2 proteins: Regulators of apoptosis or of mitochondrial homeostasis. *Nat Cell Biol* 1: 209–216, 1999
 120. Vander Heiden MG, Chandel NS, Williamson EK, et al: Bcl-X_L regulates the membrane potential and volume homeostasis of mitochondria. *Cell* 91: 627–637, 1997
 121. Wang K, Yin W-M, Chao DT, et al: BID: A novel BH3 domain-only death agonist. *Genes & Development* 10: 2859–2869, 1996
 122. Wang SL, Hawkins CJ, Yoo SJ, et al: The Drosophila caspase inhibitor DIAP1 is essential for cell survival and is negatively regulated by HID. *Cell* 98: 453–463 1999
 123. Watanabe-Fukunaga R, Brannan CI, Copeland NG, et al: Lymphoproliferation disorder in mice explained by defects in Fas antigen that mediates apoptosis. *Nature* 356:314–317, 1992
 124. Westendorp MO, Frank R, Ochsenbauer C, et al: Sensitization of T cells to CD95-mediated apoptosis by HIV-1 Tat and gp120. *Nature* 375: 497–500, 1995
 125. Wolter KG, Hsu YT, Smith CL, et al: Movement of bax from the cytosol to mitochondria during apoptosis. *J Cell Biol* 139: 1281–1292, 1997
 126. Xu DG, Bureau Y, McIntyre DC, et al: Attenuation of ischemia-induced cellular and behavioral deficits

- by X chromosome-linked inhibitor of apoptosis protein overexpression in the rat hippocampus. *Neurosci* 19: 5026–5033, 1999
127. Xu DG, Crocker SJ, Doucet JP, et al: Elevation of neuronal expression of NAIP reduces ischemic damage in the rat hippocampus. *Nat Med* 3: 997–1004, 1997
 128. Yel L, Aggarwal S, Gupta S: Cartilage-hair hypoplasia syndrome: increased apoptosis of T lymphocytes is associated with altered expression of Fas (CD95), Fas (CD95L), IAP, Bax, and Bcl2. *J Clin Immunol* 19: 428–434, 1999
 129. Yin XM, Oltvai ZN, Korsmeyer SJ: BH1 and BH2 domains of bcl-2 are required for inhibition of apoptosis and heterodimerization with bax. *Nature* 369: 321–333, 1994
 130. Yin X-M, Wang K, Gross A, et al: Bid deficient mice are resistant to Fas-induced hepatocellular apoptosis. *Nature* 400, 886–91 (1999).
 131. Yoon HJ, Carbon J: Participation of Bir1p, a member of the inhibitor of apoptosis family, in yeast chromosome segregation events. *Proc Natl Acad Sci USA* 96: 13208–13213, 1999
 132. You M, Ku PT, Hrdlickova R, et al: ch-IAP1, a member of the inhibitor-of-apoptosis protein family, is a mediator of the antiapoptotic activity of the v-rel oncoprotein. *Mol Cell Biol* 17:7328–7341, 1997
 133. Zha H, Aime-Sempe C, Sato T, et al: Proapoptotic protein Bax heterodimerizes with Bcl-2 and homodimerizes with Bax via a novel domain (BH3) distinct from BH1 and BH2. *J Biol Chem* 271: 7440–7444, 1996
 134. Zhou Q, Snipas S, Orth K, et al: Target protease specificity of the viral serpin CrmA: Analysis of five caspases. *J Biol Chem* 272: 7797–7800, 1997
 135. Zoratti M, Szabo I: Electrophysiology of the inner mitochondrial membrane. *Journal of Bioenergetics and Biomembranes* 26: 543–553, 1996
 136. Zou H, Henzel WJ, Liu X, et al: Apaf-1, a human protein homologous to *C. elegans* CED-4, participates in cytochrome c-dependent activation of caspase-3. *Cell* 90: 405–413, 1997

Address reprint requests to

John C. Reed, MD, PhD
The Burnham Institute
10901 N Torrey Pines Road
La Jolla, CA 92037

e-mail: jreed@burnham-inst.org