

Indications for Liver Transplant and AASLD Guidelines

To the Editor:

We read with great interest the recently updated AASLD Practice Guidelines published in the March issue of HEPATOLOGY titled "Evaluation for Liver Transplantation in Adults."¹ We commend Dr. Martin and colleagues for undertaking an extensive review of the literature. We think it important for your readers to be aware that, among the indications for liver transplant, other rare disorders should be included. Specifically, we write about growing interest in and a growing body of literature indicating that liver transplantation is curative for severe, unremitting acute intermittent porphyria (AIP).^{2,3} This was first described in 2004 and, thus far, most cases have been done in Europe.⁴ Recently, the first liver transplantation for AIP in the United States was performed with excellent success and prompt cure at Mt. Sinai Medical Center, NYC (Liu et al, pers. commun.).

It is also worth reminding readers that erythropoietic protoporphyria (EPP) with pigmentary cirrhosis is a well-established indication for liver or combined liver/bone marrow transplantation. Numerous case reports and publications in various peer-reviewed journals attest to the role of liver or combined liver/bone marrow transplantation in patients with chronic liver disease secondary to EPP.^{5,6} In fact, among the porphyrias, chronic liver disease in EPP is probably the best established indication for liver transplant.

We recently published a review article assessing the role of liver transplantation in porphyria.⁷ Because of the growing influence of AASLD guidelines for clinical practice and decisions regarding reimbursement, etc., we hope that the AASLD guidelines will soon be amended to take into account liver transplantation for carefully selected patients with porphyrias.

MANISH THAPAR, M.D.¹
HERBERT L. BONKOVSKY, M.D.²
¹Department of Medicine
Drexel University College of Medicine
Philadelphia, PA

²Department of Medicine
Carolinas HealthCare System and University of NC School
of Medicine
Charlotte, NC

References

1. Martin P, DiMartini A, Feng S, Brown R Jr, Fallon M. Evaluation for liver transplantation in adults: 2013 practice guideline by the American Association for the Study of Liver Diseases and the American Society of Transplantation. HEPATOLOGY 2014;59:1144-1165.
2. Dowman JK, Gunson BK, Mirza DF, Bramhall SR, Badminton MN, Newsome PN, et al. Liver transplantation for acute intermittent porphyria is complicated by a high rate of hepatic artery thrombosis. Liver Transpl 2012;18:195-200.
3. Dowman JK, Gunson BK, Bramhall SR, Badminton MN, Newsome PN. Liver transplantation from donors with acute intermittent porphyria. Ann Intern Med 2011;154:571-572.
4. Soonawalla ZF, Orug T, Badminton MN, Elder GH, Rhodes JM, Bramhall SR, et al. Liver transplantation as a cure for acute intermittent porphyria. Lancet 2004;363:705-706.
5. Wählin S, Stal P, Adam R, Karam V, Porte R, Seehofer D, et al. Liver transplantation for erythropoietic protoporphyria in Europe. Liver Transpl 2011;17:1021-1026.
6. McGuire BM, Bonkovsky HL, Carithers RL Jr, Chung RT, Goldstein JI, Lake JR, et al. Liver transplantation for erythropoietic protoporphyria liver disease. Liver Transpl 2005;11:1590-1596.
7. Singal A, Parker C, Bowden C, Thapar M, Liu L, McGuire B. Liver transplantation in the management of porphyria. HEPATOLOGY 2014; DOI: 10.1002/hep.27086.

Copyright © 2014 by the American Association for the Study of Liver Diseases.
View this article online at wileyonlinelibrary.com.

DOI 10.1002/hep.27176

Potential conflict of interest: Dr. Bonkovsky consults, advises, and received grants from Clinuvel. He consults and advises Recordati and Alynham. He received grants from Vertex.

Association Between Response to Pegylated Interferon/Ribavirin Therapy and Ribavirin Levels

To the Editor:

We read with great interest the article by Holmes et al.¹ published in HEPATOLOGY. The researchers conclude that inosine triphosphatase (ITPase) deficiency protects against ribavirin (RBV)-induced anemia, but that it is not associated with sustained virological response (SVR) to pegylated interferon (Peg-IFN)- α -2a/RBV therapy in HCV genotype 1 (HCV-1)-infected patients from the CHARIOT study.

Rapid virological response (RVR) has been considered the most important predictor for SVR, and an abbreviated treatment regimen was encouraged in patients with RVR and a low viral load.^{2,3} We wonder whether RVR (32.8%) had any effect on the result of the association between SVR and hemoglobin decline or upon RBV pharmacokinetics in the study by Holmes et al.¹ It is known that plasma RBV levels at week 8, rather than the actual hemoglobin levels or ITPase deficiency in patients with HCV-1 and genotype 2/3 infection,^{1,4} have been associated with SVR. Since the very important role of the genetic variation in interleukin (IL)28B on predicting response to Peg-IFN-based therapies either without^{3,5,6} or with new

direct antiviral agents⁷ has been recognized, it seems also necessary to take the IL28B genotype into consideration in elucidating the influence of RBV levels on SVR in HCV-1 patients, without otherwise detracting from the results by Holmes et al.¹

CHIA-YEN DAI, M.D., PH.D.^{1,2,3}
MING-LUNG YU, M.D., PH.D.^{1,2,3}
WAN-LONG CHUANG, M.D., PH.D.^{1,2}
¹Hepatobiliary Division
Department of Internal Medicine
Kaohsiung Medical University Hospital
Kaohsiung Medical University
Kaohsiung, Taiwan
²Hepatitis Center
Kaohsiung Medical University Hospital
Kaohsiung Medical University
Kaohsiung, Taiwan
³Center for Health Management
Kaohsiung Medical University Hospital
Kaohsiung Medical University
Kaohsiung, Taiwan