

- OKT-3 monoclonal antibody in pediatric kidney transplant recipients with recurrent and resistant allograft rejection. *J Pediatr* 1987; 111 : 45-50.
4. Bailey LL, Assad AN, Trimm RF, *et al.* Orthotopic transplantation during early infancy as therapy for incurable congenital heart disease. *Ann Surg* 1988; 3 : 279-286.
 5. Baum D, Stinson EB, Shumway NE. The place for heart transplantation in children. *Pediatr Cardiol* 1981; 4 : 741-747.
 6. Kemnitz J, Choritz H, Cohnert TR, *et al.* Predictive implications of bioptic diagnosis in cardiac allografts. *J Heart Transplant* 1989; 8 : 315-329.
 7. Oyer PE, Stinson EB, Bieber CP. Diagnosis and treatment of acute cardiac allograft rejection. *Transplant Proc* 1979; 11 : 296-303.
 8. Zerbe TR, Arena V. Diagnostic reliability of endomyocardial biopsy for assessment of cardiac allograft rejection. *Hum Pathol* 1988; 19 : 1307-1314.
 9. Sibley RK, Olivari MT, Ring WS, *et al.* Endomyocardial biopsy in the cardiac allograft recipient. *Ann Surg* 1986; 203 : 177-187.
 10. Esquivel CO, Iwatsuki S, Gordon RD, *et al.* Indication for pediatric liver transplantation. *J Pediatr* 1991; 119 : 1039-1045.
 11. Malatack JJ, Schaid DJ, Urbach AH, *et al.* Choosing a pediatric recipient for orthotopic liver transplantation. *J Pediatr* 1987; 111 : 479-489.
-

VARIATIONS IN THE NRAMP1 GENE AND SUSCEPTIBILITY TO TUBERCULOSIS

Genetic factors may affect the susceptibility to tuberculosis, but no specific genes governing susceptibility have been identified. In mice, natural resistance to infection with some mycobacteria is influenced by the gene for natural-resistance-associated macrophage protein 1 (Nramp 1), but the role of human homologue of this gene, NRAMP1 in TB is unknown.

A case-control study of TB in Gambia, W. Africa typed polymorphisms in NRAMP1 using sequence-specific oligonucleotide hybridisation and microsatellite analysis. Subjects were 410 adults (mean age, 34.7 years) with smear-positive pulmonary tuberculosis and 417 ethnically matched, healthy controls. Patients with HIV infection were excluded.

Results showed that four NRAMP1 polymorphisms were each significantly associated with TB. Subjects heterozygous for two NRAMP1 polymorphisms in intron 4 and the 3' untranslated region of gene were particularly overrepresented among those with TB, as compared with those with the common NRAMP1 genotype (odds ratio, 4.07; 95% CI, 1.86 to 9.12; chi-square = 14.58; $p < 0.001$). Genetic variation in NRAMP1 affects susceptibility to tuberculosis in W. Africans.

Abstracted from : N Engl J Med 1998; Vol 338 : No. 10; pp 640-644.