

# Giant Nernst effect in heavy-electron metals

K. Behnia<sup>a,\*</sup>, R. Bel<sup>a</sup>, A. Pourret<sup>a</sup>, K. Izawa<sup>b</sup>, J. Flouquet<sup>b</sup>, Y. Nakajima<sup>c</sup>, Y. Matsuda<sup>c</sup>,  
D. Kikuchi<sup>d</sup>, Y. Aoki<sup>d</sup>, H. Sugawara<sup>d</sup>, H. Sato<sup>d</sup>

<sup>a</sup>*Laboratoire de Physique Quantique (CNRS), ESPCI, F-75231 Paris, France*

<sup>b</sup>*DRFMC/SPSMS, Commissariat à l'Energie Atomique, F-38042 Grenoble, France*

<sup>c</sup>*Department of Physics, University of Kyoto, Kyoto 608-8502, Japan*

<sup>d</sup>*Department of Physics, Tokyo Metropolitan University, Tokyo 192-0397, Japan*

Available online 17 November 2006

## Abstract

Recent studies of the Nernst effect in a number of heavy-fermion systems have led to a previously unsuspected result. In some circumstances, the Nernst signal of quasi-particles becomes very large and can easily overwhelm the well-known Nernst effect produced by the movement of the superconducting vortices under the influence of a thermal gradient. In particular, the Nernst coefficient attains an exceptionally large magnitude in the ordered states of URu<sub>2</sub>Si<sub>2</sub> and PrFe<sub>4</sub>P<sub>12</sub>. In all these cases, the order of magnitude of the Nernst signal appears compatible with the Boltzmann picture which links the Nernst coefficient to the energy-dependence of the Hall angle.

© 2006 Published by Elsevier B.V.

PACS: 74.70.Tx; 72.15.Jf; 71.27.+a

Keywords: Thermoelectricity; Hidden order; Nernst effect; Quantum criticality

## 1. Introduction

The Nernst effect, the generation of a transverse electric field by a longitudinal thermal gradient has attracted much attention since the observation of a finite Nernst signal in the normal state of high- $T_c$  cuprates [1]. This uncommon transport coefficient is known to be enhanced in presence of vortex movement induced by thermal gradient in the vortex-liquid state of a type II superconductor [2]. In metals, on the other hand, due to a phenomenon dubbed Sondheimer cancelation [3], the Nernst coefficient, while much less explored, was believed to be small.

During the last few years, this belief has been countered by a host of experimental evidence. Several studies have established that Nernst effect in a metal can become quite large. While a rigorous interpretation of the Nernst data remains a challenge to the theory, the order of magnitude of the Nernst coefficient is compatible with the rough

estimate obtained using a Boltzmann picture. A simple argument indicates that a dilute liquid of heavy quasi-particles with a long mean-free-path should generate a large Nernst signal with possible applications for thermoelectric cooling at low temperatures.

## 2. Ambipolar Nernst effect

The study of NbSe<sub>2</sub> [4] revealed that a finite negative Nernst signal persists at temperatures well above  $T_c = 7.2$  K. The maximum in the Nernst signal occurs when the contribution of hole-like and electron-like carriers to the Hall coefficient cancel out and the system becomes accidentally compensated. This result recalls that in a multi-band metal the Sondheimer cancelation is not valid and the flow of quasi-particles of opposite charge in presence of a thermal gradient can lead to an enhanced Nernst signal. This phenomenon known as the ambipolar Nernst effect, has been known since a long time ago in semi-conductor physics [5].

\*Corresponding author. Tel.: +33 1 40 79 46 26; fax: +33 1 40 79 47 44.

E-mail address: [kamran.behnia@espci.fr](mailto:kamran.behnia@espci.fr) (K. Behnia).

### 3. Nernst effect in heavy-fermion metals

In general, heavy-fermion compounds are multi-band metals and therefore the existence of a finite Nernst signal is not a surprise by itself. As seen in Fig. 1, in many cases, the magnitude of the Nernst effect can become orders of magnitude larger than what was found in NbSe<sub>2</sub>. This simply reflects the smallness of the Fermi energy which sets an energy scale roughly the inverse of the thermoelectric coefficients of the system. In the case of the heavy-fermion superconductor CeCoIn<sub>5</sub>, the quasi-particle Nernst signal easily overwhelms the vortex signal associated with the superconducting transition [6]. This large Nernst coefficient ( $\sim 1 \mu\text{V}/\text{KT}$ ) appeared to be confined to a region of the (H,T) plane where other transport properties of the system display a non-Fermi liquid behavior. Further exploration of heavy-fermion metals led to the discovery of cases with an even larger Nernst coefficient. URu<sub>2</sub>Si<sub>2</sub> is notorious for the presence of a mysterious electronic order below  $T_0 = 17.5 \text{ K}$  [8]. It has been recently found that this ordering leads to a Nernst coefficient of a large ( $\sim 4 \mu\text{V}/\text{KT}$ ) magnitude [9]. The exploration of PrFe<sub>4</sub>P<sub>12</sub> which is believed to present an antiferroquadrupolar (AFQ) ordering below  $T_0 = 6.5 \text{ K}$  [10], revealed an even larger Nernst coefficient in the ordered state ( $\sim 50 \mu\text{V}/\text{KT}$ ) [11].

### 4. Correlation between the Nernst coefficient and the Hall angle

In the Boltzmann picture the Nernst coefficient tracks the energy dependence of the tangent of the Hall angle ( $\tan \theta_H = \rho_{xy}/\rho_{xx}$ ) at the Fermi level [3,12]:

$$v = \frac{1}{B} \frac{\pi^2 k_B^2 T}{3e} \frac{\partial \tan \theta_H}{\partial \varepsilon} \Big|_{\varepsilon_F}, \quad (1)$$

where  $k_B$  and  $e$  are, respectively, the Boltzmann constant and the fundamental charge. Since in a first approximation,




Fig. 1. Nernst coefficient of a number of heavy fermion compounds in the zero-field limit. For comparison, the figure displays data for gold and NbSe<sub>2</sub>, a multi-band metal.

this energy derivative is just proportional to  $\tan \theta_H/\varepsilon_F$ , a combination of a large Hall angle and a small Fermi energy can lead to a very large Nernst signal. The cases mentioned above seem to corroborate this picture.

The non-Fermi liquid state in CeCoIn<sub>5</sub> is characterized by a large Hall coefficient [13] and a large effective mass [7] (indicative of the proximity of a quantum critical point).

In both URu<sub>2</sub>Si<sub>2</sub> and PrFe<sub>4</sub>P<sub>12</sub>, ordering leads to a drastic drop in the carrier density and an enhancement of the electronic mean-free-path, the combination of these two factors leads to a spectacular increase in the Hall angle. In both cases, the residual quasi-particles of the ordered state still present a renormalized mass. The small Fermi wave-vector and the large effective mass of the quasi-particles in the ordered state have been directly observed by de Haas van Alphen studies [14,15]. These two features conspire to keep the Fermi energy small. Therefore, according to Eq. (1), a large Nernst coefficient is expected to emerge below the ordering temperature. The ordered state of these two compounds can be assimilated to a clean heavy-fermion semi-metal providing the three independent ingredients of a giant Nernst signal: a low carrier density, a large mass enhancement and a long mean-free-path. Further studies of the Nernst coefficient in more familiar heavy-fermion semi-metals such as CeNiSn would tell if these are the only ingredients needed to create a giant Nernst coefficient.

### 5. Towards an Ettingshausen refrigerator

Besides its fundamental aspect, the search for the origins of a giant Nernst signal is interesting due to its potential applications. An Ettingshausen cooler [16] can be conceived using a compound with a thermomagnetic figure of merit ( $Z'T = N^2 T/\kappa\sigma$ ) approaching unity, where  $N = v/B$  is the Nernst signal and  $\kappa$  and  $\sigma$  are, respectively, thermal and electric conductivities. For a metal, thermal conductivity is expected to be dominated by the electronic contribution at low temperatures and therefore, the Wiedemann–Franz law translates this requirement to  $N \approx 155 \mu\text{V}/\text{K}$ . The case of PrFe<sub>4</sub>P<sub>12</sub> indicates that this magnitude may be reached in a clean heavy-fermion semi-metal.

### References

- [1] Z.A. Xu, et al., Nature 406 (2000) 486.
- [2] R.P. Huebner, A. Seher, Phys. Rev. 181 (1969) 701.
- [3] Y. Wang, et al., Phys. Rev. B 64 (2001) 224519.
- [4] R. Bel, K. Behnia, H. Berger, Phys. Rev. Lett. 91 (2003) 066602.
- [5] R.T. Delves, Rep. Prog. Phys. 28 (1965) 249.
- [6] R. Bel, et al., Phys. Rev. Lett. 92 (2004) 217002.
- [7] C. Petrovic, et al., J. Phys. Condens. Matter 13 (2001) L337.
- [8] T.T.M. Palstra, et al., Phys. Rev. Lett. 55 (1985) 2727; W. Schlabbitz, et al., Phys. B 62 (1986) 171; M.B. Maple, et al., Phys. Rev. Lett. 56 (1986) 185.
- [9] R. Bel, et al., Phys. Rev. B 70 (2004) 220501(R).
- [10] H. Sato, et al., Phys. Rev. B 62 (2000) 15125; Y. Aoki, et al., J. Phys. Soc. Japan 74 (2005) 209.

- [11] A. Pourret, et al., Phys. Rev. Lett. 96 (2006) 176402.
- [12] V. Oganessian, I. Ussishkin, Phys. Rev. B 70 (2004) 054503.
- [13] Y. Nakajima, et al., J. Phys. Soc. Japan 73 (2004) 5.
- [14] H. Ohkuni, et al., Philos. Mag. B 79 (1999) 1045.
- [15] H. Sugawara, et al., Phys. Rev. B 66 (2002) 134411.
- [16] K. Scholz, et al., J. Appl. Phys. 75 (1994) 5406.