

Original article

CRNK gene transfer improves function and reverses the myosin heavy chain isoenzyme switch during post-myocardial infarction left ventricular remodeling

Davin L. Hart^a, Maria C. Heidkamp^a, Rekha Iyengar^a, Kalpana Vijayan^a, Erika L. Szotek^a,
John A. Barakat^a, Marysa Leya^a, Marcus Henze^b, Karie Scrogin^b,
Kyle K. Henderson^a, Allen M. Samarel^{a,c,*}

^a The Cardiovascular Institute, and the Department of Medicine, Loyola University Chicago Stritch School of Medicine, Maywood, Illinois 60153, USA

^b The Cardiovascular Institute, and the Department of Pharmacology, Loyola University Chicago Stritch School of Medicine, Maywood, Illinois 60153, USA

^c The Cardiovascular Institute, and the Department of Physiology, Loyola University Chicago Stritch School of Medicine, Maywood, Illinois 60153, USA

Received 17 August 2007; received in revised form 1 April 2008; accepted 7 April 2008

Available online 16 April 2008

Abstract

PYK2 is a Ca^{2+} -dependent, nonreceptor protein tyrosine kinase that is involved in the induction of left ventricular hypertrophy (LVH) and its transition to heart failure. We and others have previously investigated PYK2's function *in vitro* using cultured neonatal and adult rat ventricular myocytes as model systems. However, the function of PYK2 in the *in vivo* adult heart remains unclear. Here we evaluate the effect of PYK2 inhibition following myocardial infarction (MI) using adenoviral (Adv) overexpression of the C-terminal domain of PYK2, known as CRNK. First we demonstrate that CRNK functions as a dominant-negative inhibitor of PYK2-dependent signaling, presumably by displacing PYK2 from focal adhesions and costameres. Then, male Sprague–Dawley rats (~300 g) underwent permanent left anterior descending coronary artery ligation. One wk post-MI, either Adv-GFP ($n=34$) or Adv-CRNK ($n=28$) was administered (10^{10} pfu, 0.1 ml) via catheter-based, Optison[®]-mediated gene transfer. LV structure and function were evaluated by echocardiography 1 and 3 wk after gene transfer, and LV tissue was analyzed by real-time RT-PCR and Western blotting. CRNK overexpression was readily detected by Western blotting 1 wk following gene transfer. Adv-CRNK improved overall survival ($P=0.03$; Logrank Test) and LV fractional shortening ($23\pm2\%$ vs. $31\pm2\%$ for Adv-GFP vs. Adv-CRNK infected animals, respectively; $P<0.05$). Whereas MI hearts exhibited increased β -, and decreased α -myosin heavy chain (MHC) mRNA expression characteristic of LVH, Adv-CRNK reversed the MHC isoenzyme switch (3.3 ± 1.4 fold increase in α MHC; 0.4 ± 0.1 fold decrease in β MHC; $P<0.05$ for both). In summary, CRNK gene transfer improves survival, increases LV function, and alters MHC gene expression suggesting an attenuation of LV remodeling post-MI.

© 2008 Elsevier Inc. All rights reserved.

Keywords: Signal transduction; Heart failure; PYK2; Gene therapy; Focal adhesion kinase; Optison; Ultrasound

1. Introduction

Segmental loss of viable myocardium following myocardial infarction (MI) activates cardiomyocyte signal transduction pathways leading to pathological left ventricular hypertrophy (LVH). Pathological LVH is often associated with hemodynamic

overload and decreased systolic and/or diastolic performance, which can ultimately lead to the clinical syndrome of heart failure (HF). The progressive structural and functional alterations in both the infarcted and non-infarcted LV myocardium that occur following irreversible ischemic injury are referred to as post-MI ventricular remodeling [1]. Interventions that block the maladaptive cellular signaling leading to ventricular remodeling may also be useful in preventing or attenuating loss of ventricular performance seen in HF.

The Ca^{2+} -dependent, nonreceptor protein tyrosine kinase PYK2 has been previously implicated in cardiomyocyte cell

* Corresponding author. The Cardiovascular Institute, Loyola University Medical Center, Building 110, Room 5222, 2160 South First Avenue, Maywood, IL 60153, USA. Tel.: +1 708 327 2821; fax: +1 708 327 2849.

E-mail address: asamarel@lumc.edu (A.M. Samarel).

signaling pathways leading to LVH and HF [2–6]. PYK2 is a member of the focal adhesion kinase (FAK) family of nonreceptor PTKs. PYK2 coordinates integrin-, Ca^{2+} -, and PKC-dependent signal transduction in a number of tissues (for review, see [7]). PYK2 expression and phosphorylation are regulated by intracellular Ca^{2+} and protein kinase C (PKC) in cardiomyocytes and vascular smooth muscle cells [8–10]. As examined in various cell lines, PYK2 serves as an “activatable” scaffolding protein, and transduces signals from G-protein coupled receptors to the mitogen-activated protein kinases (MAPK) and the phosphoinositol-3-kinase (PI-(3)K)-PDK1-Akt signaling pathway depending upon which adaptor proteins bind to the phosphorylated enzyme [6,11–13]. PYK2 has also been shown to link a variety of stressful stimuli, including Ca^{2+} overload, UV irradiation, and H_2O_2 and TNF- α treatment to MAPK and Akt activation in many cell types.

Like FAK, the function of PYK2 is regulated by an endogenously expressed inhibitor, known as PYK2-Related Non-Kinase (PRNK) [14], also known as Cell Adhesion Kinase- β -Related Non-Kinase (CRNK) [15]. CRNK consists of the C-terminal portion of PYK2, containing the focal adhesion targeting sequence, but lacking the N-terminal autophosphorylation site and kinase domain. CRNK is structurally analogous to FAK-Related Non-Kinase (FRNK), the autonomously expressed C-terminal domain of FAK. CRNK transcripts are generated by either alternative splicing of the PYK2 hnRNA, or, as is the case of FRNK, by utilization of an intronic promoter [16]. CRNK is expressed at relatively high levels in the brain and other organs [14]. When ectopically expressed, CRNK can inhibit PYK2 (but not FAK) tyrosine autophosphorylation at Y₄₀₂, presumably by displacing PYK2 from its cytoskeleton-binding sites [15]. Thus CRNK, like its structurally homologous polypeptide FRNK, has been used as a tool to specifically inhibit PYK2-dependent signal transduction [17–20].

In this study we evaluated the effect of PYK2 inhibition using adenoviral (Adv)-mediated overexpression of CRNK in cultured cardiomyocytes and in an *in vivo* model of post-MI ventricular remodeling. Data are presented to indicate that CRNK prevents PYK2 autophosphorylation and downstream signaling in cultured heart cells. Furthermore, CRNK gene transfer *in vivo* improves survival, increases LV function, and alters MHC gene expression suggesting an attenuation of LV remodeling post-MI.

2. Methods

2.1. Reagents

PC-1 tissue culture medium was obtained from BioWhittaker (Walkersville, MD). Dulbecco's Modified Eagle Medium (DMEM) and Medium 199 were obtained from Gibco BRL (Grand Island, NY). Anti-FLAG M2 and anti-sarcomeric α -actinin monoclonal antibodies (mAb) were obtained from Sigma (St. Louis, MO). Rhodamine-conjugated goat anti-mouse IgG and FITC-conjugated phalloidin were obtained from Molecular Probes (Eugene, OR). Phospho-specific PYK2-Y₄₀₂, phospho-specific Akt-S₄₇₃, and Akt polyclonal Ab (pAb) were purchased from Cell Signaling Technology (Danvers, MA). N-terminal PYK2 pAb was

obtained from BioLegend (San Diego, CA). C-terminal PYK2/CRNK mAb was obtained BD Transduction Laboratories (San Jose, CA). Horseradish peroxidase-conjugated goat anti-rabbit and goat anti-mouse IgG were obtained from BioRad (Hercules, CA). Real-time RT-PCR reagents were obtained from GE Healthcare Biosciences (Piscataway, NJ), and Invitrogen (Carlsbad, CA). cDNA primers and probes were obtained from Integrated DNA Technologies, Inc. (Coralville, IA). All other reagents were of the highest grade commercially available and were obtained from Sigma, and Baxter S/P (McGaw Park, IL).

2.2. Cell culture

Animals used in these experiments were handled in accordance with the Guiding Principles in the Care and Use of Animals, approved by the Council of the American Physiological Society. Neonatal rat ventricular myocytes (NRVM) were isolated from the hearts of 2-day old Sprague-Dawley rats by collagenase digestion, as previously described [21]. Cells were pre-plated for 1 h in serum-free PC-1 medium to reduce non-myocyte contamination. The nonadherent NRVM were then plated at a density of 1600 cells per mm^2 onto collagen-coated chamberslides or 60 mm dishes, and left undisturbed in a 5% CO_2 incubator for 36 h. Unattached cells were removed by aspiration, washed twice in HBSS, and the attached cells were maintained in a solution of DMEM/Medium 199 (4:1) containing antibiotic/antimycotic solution. NRVM were infected (24 h) with replication-defective adenoviruses (Adv) diluted in DMEM/Medium 199. Medium was then replaced with virus-free DMEM/Medium 199, and the cells stimulated with agonists, or cultured for an additional 24 h. Adult rat ventricular myocytes (ARVM) were isolated according to Westfall et al. [22] with modification as described previously [9]. Myocytes were then either fixed and permeabilized, or plated in PC-1 medium onto laminin-coated (15 $\mu\text{g}/\text{ml}$) chamberslides for 60 min in a 5% CO_2 incubator. Unattached myocytes were removed and attached myocytes were maintained in PC-1 medium, and infected with Adv for 24 h.

2.3. Adenoviral constructs

A replication-deficient Adv encoding FLAG-tagged, human CRNK (Adv-CRNK) was kindly provided by Dr. Andrey Sorokin, Medical College of Wisconsin [18]. An Adv expressing GFP (Adv-GFP) was used to control for nonspecific effects of Adv infection [23]. An Adv expressing nuclear-encoded β -galactosidase (Adv-ne β gal; kindly provided by Dr. MK Rundell, Northwestern University) was used in some *in vivo* gene transfer experiments to demonstrate the regions of ventricular myocardium transduced by adenoviral infection. The multiplicity of viral infection (moi) was determined by dilution assay in HEK293 cells grown in 96 well clusters.

2.4. Immunolocalization

NRVM and ARVM were fixed and permeabilized as previously described [9,24]. Myocytes were stained with anti-

FLAG mAb followed by rhodamine-conjugated goat anti-mouse IgG, and counterstained with FITC-phalloidin. Fluorescently-labeled cells were then viewed using a Zeiss LSM 510 laser scanning confocal microscope.

2.5. Western blotting

Cultured cells and tissue samples were homogenized in lysis buffer [25], and equal amounts of extracted proteins (50–150 µg) were separated by SDS-PAGE and Western blotting on 10% polyacrylamide gels. Primary antibody binding was detected with horseradish peroxidase-conjugated goat anti-mouse or goat anti-rabbit secondary antibodies, and visualized by enhanced chemiluminescence (Pierce Biotechnology, Rockford, IL). Developed films were then scanned on a HP Deskjet 4890 Scanner, and band intensity was quantified using UN-SCAN-IT Gel Software, Ver. 6.1 (Silk Scientific, Orem UT).

2.6. Cellular composition

For the analysis of NRVM hypertrophy, cellular protein and DNA were quantitatively scraped from dishes in 0.2 N perchloric acid and collected by centrifugation (10,000 ×g, 10 min). The precipitate was re-dissolved by incubation (60 °C, 20 min) in 0.3 N KOH. Aliquots were used for analysis of total protein by the Lowry method with crystalline human serum albumin as standard, and for DNA using 33258 Hoechst dye and salmon sperm DNA as standard [26].

2.7. mRNA analysis

Total cellular RNA was isolated from NRVM using the RNeasy Mini Kit (Qiagen, Inc., Valencia, CA) or from LV tissue by the method of Chomczynski and Sacchi [27]. RNA was quantified by absorbance at 260 nm and its integrity was determined by examining the 28S and 18S rRNA bands in ethidium bromide-stained agarose gels. SERCA2, ANF, αMHC, and βMHC mRNAs were then analyzed by real-time RT-PCR, as previously described [5,28]. The mixture consisted of 1 µl of sample cDNA, 21 µl DEPC water, 25 µl Platinum Quantitative PCR SuperMix-UDG, and 3 µl of a primer/dual-labeled probe combination specific for each gene of interest (Table 1). TaqMan® Rodent GAPDH or 18S rRNA Control Reagents were obtained from Applied Biosystems, (Foster City, CA) and used to normalize mRNA expression levels. Probes were labeled at the 5' end with 6-FAM and at the 3' end with BHQ-1. Specific gene primers were 10 µM, and probe was

1 µM. Rodent GAPDH and 18S rRNA primers were 10 µM, and probe was 5 µM. PCR amplification was performed by cycling between 95 °C (15 s) and 60 °C (60 s) for 45 cycles, using the 6-FAM fluorophore for quantification. All samples were run in triplicate, and the results were averaged. Data were analyzed as previously described [28].

2.8. Coronary artery ligation

Adult male Sprague–Dawley rats weighing ~300 g were anesthetized with ketamine (70 mg/kg IM) and xylazine (10 mg/kg IM), intubated, and placed on a heated operating table. Ligation of the left anterior descending coronary artery was then performed as described by Pfeffer et al. [29]. Sham-operated animals underwent surgery without placement of the coronary ligature.

2.9. M-mode and 2-D echocardiography

Transthoracic 2-D guided, M-mode echocardiography was performed on anesthetized rats using an Acuson 15L8 Microson High Resolution Transducer and an Acuson Sequoia C256 Echocardiography System (Siemens Medical Solutions USA; Malvern, PA). Images were analyzed in real-time, or saved onto 630 mbyte magneto-optical discs for off-line analysis.

2.10. Adenoviral gene transfer

A modification of a catheter-based, Optison®-mediated gene transfer technique [30] was used to administer Adv-neβgal, Adv-GFP or Adv-CRNK to the myocardium of normal rats, and rats 1 wk after coronary artery ligation. Briefly, animals were anesthetized with ketamine/xylazine, and ventilated with O₂ administered via a conical facial mask. A right carotid artery cutdown was used to insert a custom-designed, 2.5 F double-lumen, 8 mm long × 4 mm diameter balloon catheter (Tyshak Mini Pediatric Valvuloplasty Catheter; NuMED, Inc., Hopkinton, NY) into the ascending aorta. With the tip of the catheter positioned under echocardiographic guidance exactly 3 mm above the aortic valve, the balloon was inflated, thus completely occluding the proximal aorta. Acetylcholine (1.5 µg in 100 µl saline) was injected to transiently arrest the heart and induce coronary vasodilation, followed immediately by a mixture of Adv (~10¹⁰ pfu in 100 µl)+Optison® echocontrast reagent (100 µl) and then 1 ml of saline flush. The balloon was deflated, and the 15L8 ultrasound transducer was positioned over the heart. The mechanical index of the transducer was raised to 1.9, and the

Table 1
Primers and probes used for real-time RT-PCR

cDNA	Sense primer	Antisense primer	Probe
SERCA2	5' TCT GTC ATT CGG GAG TGG GG 3'	5' GCC CAC ACA GCC AAC GAA AG 3'	5' TGG CCA CTC ATG ACA ACC CG 3'
ANF	5' CTT GCG GTG TGT CAC ACA GC 3'	5' GGG AGA GGT AAG GCC TCA CT 3'	5' CCT CCT GGA GCT GCA GCT TC 3'
αMHC	5' ACA GAG TGC TTC GTG CCT GAT 3'	5' CGA ATT TCG GAG GGT TCT GC 3'	5' ACA GTC ACC GTC TTG CCG TTT TCA GT 3'
βMHC	5' GCT ACC CAA CCC TAA GGA TGC 3'	5' TCT GCC TAA GGT GCT GTT TCA 3'	5' TGT GAA GCC CTG AGA CCT GGA GCC 3'

Probes were labeled at the 5' end with 6-FAM and at the 3' end with BHQ-1.

injected microbubbles were disrupted over the next 5 min. Finally, the catheter was removed, the carotid artery was tied off and the incision closed, and the animal allowed to recover.

2.11. Tissue fixation and β -galactosidase staining

Following induction of general anesthesia, the heart was removed, rinsed in ice-cold Krebs-Henseleit solution and the ascending aorta was cannulated with PE-240 tubing. The coronary circulation was pressure-perfusion flushed (5 min, 25 °C) with Krebs-Henseleit solution at 80 mm Hg and then fixed with Krebs-Henseleit solution containing 0.2% (v/v) glutaraldehyde (10 min, 25 °C). Hearts were embedded into OCT Compound (Sakura Finetek, Torrance, CA), frozen to –20 °C, and sectioned in the short axis to obtain 14 μ m thick sections. Sections were stained for β -galactosidase with X-gal (Takara Bio USA, Madison, WI) in a humidified chamber at 37 °C for 2 h in the dark, according to manufacturer's instructions. Specimens were washed 3 times with PBS, counterstained with eosin solution (Sigma; St. Louis, MO) for 30 s, and washed three times with PBS. Images were obtained with an Olympus microscope fitted with a Nikon digital camera.

2.12. Flow cytometry

Freshly isolated ARVM were fixed in 4% paraformaldehyde and permeabilized with 0.1% saponin. Cells were then stained with anti-sarcomeric α -actinin monoclonal antibody, followed by Alexa Fluor® goat anti-mouse IgG to distinguish the myocyte vs.

non-myocyte population. Flow cytometry was performed using a BD FACSCanto Flow Cytometer running FACSDiva software.

2.13. Data analysis

Results were expressed as means \pm SEM. Normality was assessed using the Kolmogorov–Smirnov test. Data were compared using one-way ANOVA followed by Student–Newman–Keuls test, one-way ANOVA on Ranks followed by Dunn's test, or unpaired *t*-test, where appropriate. Differences among means were considered significant at $P < 0.05$. Data were analyzed using SigmaPlot for Windows, Ver. 9.0 (Systat Software, San Jose, CA).

3. Results

3.1. CRNK overexpression and localization in NRVM and ARVM

Initial experiments were conducted in cultured cardiomyocytes to analyze FLAG-tagged, CRNK expression, and to verify its function as an inhibitor of PYK2-dependent signal transduction. NRVM and ARVM were infected (5 or 100 moi, respectively; 24 h) with Adv-CRNK and the transgene was localized by immunocytochemistry with anti-FLAG monoclonal antibody (1:10) and confocal microscopy. Of note, CRNK expression did not appear to affect cell morphology, nor did it cause cell detachment over this time period. As seen in Fig. 1A, CRNK was found in a reticular pattern throughout the cytoplasm




Fig. 1. Immunolocalization of CRNK in neonatal and adult cardiomyocytes. In Panel A, NRVM grown on Permax® chamberslides were infected with Adv-CRNK (5 moi, 24 h). Cells were then fixed, permeabilized, stained with anti-FLAG monoclonal antibody (1:10) followed by rhodamine-conjugated goat anti-mouse secondary antibody, and counterstained with FITC-conjugated phalloidin to detect actin filaments. Control chamberslides were handled in an identical fashion, except that the anti-FLAG antibody was omitted (No Primary Ab). Cells were then viewed with a confocal microscope. In Panel B, ARVM were infected (100 moi, 24 h) with Adv-CRNK. Cells were then fixed, permeabilized and stained with anti-FLAG antibody followed by rhodamine-conjugated goat-anti-mouse secondary antibody. Cells were then optically sectioned (1 μ m) using the confocal microscope.

of NRVM, but was excluded from the nucleus. In contrast, CRNK was predominantly localized to rib-like structures adjacent to the sarcoplasmic membrane of ARVM, consistent with the appearance of costameres (Fig. 1B). The difference in CRNK localization between NRVM and ARVM was reminiscent of the difference in PYK2 localization between neonatal and adult cardiomyocytes observed in our previous studies [5,9].

3.2. CRNK overexpression inhibits PYK2-dependent signal transduction

We next examined whether CRNK inhibited cardiomyocyte PYK2 activation in response to known activators of the kinase [2,6,9,31,32]. As seen in Fig. 2A, NRVM infected with a




Fig. 2. CRNK overexpression inhibits basal, and endothelin- and H₂O₂-induced PYK2 phosphorylation. In Panel A, NRVM were infected (5 moi, 24 h) with either Adv-GFP or Adv-CRNK. Cells were then left untreated, or stimulated (10 min) with endothelin-1 (ET, 100 nM) or H₂O₂ (100 μ M). Equal amounts of extracted cellular protein (50–150 μ g) were then separated by SDS-PAGE and Western blotting (WB). The position of molecular weight standards is indicated at the left of each blot. In Panel B, the quantitative analysis of 6–8 experiments is depicted. * P <0.05 vs. untreated, Adv-GFP infected cells; + P <0.05 for Adv-GFP vs. Adv-CRNK for each treatment group.




Fig. 3. CRNK overexpression inhibits basal and H₂O₂-induced Akt phosphorylation. In Panel A, NRVM were infected (5 moi, 24 h) with either Adv-GFP or Adv-CRNK. Cells were then left untreated, or stimulated (10 min) with endothelin-1 (ET, 100 nM) or H₂O₂ (100 μ M). Equal amounts of extracted cellular protein (50 μ g) were then separated by SDS-PAGE and Western blotting (WB). The position of molecular weight standards is indicated at the left of each blot. In Panel B, the quantitative analysis of 6 experiments is depicted. * P <0.05 vs. untreated, Adv-GFP infected cells; + P <0.05 for Adv-GFP vs. Adv-CRNK for each treatment group.

control adenovirus (Adv-GFP) expressed little endogenous CRNK, but contained readily detectable amounts of PYK2 that was phosphorylated at Y₄₀₂ under basal conditions. Endothelin-1 (ET; 100 nM, 10 min) and H₂O₂ (100 μ M, 10 min) further increased PYK2 phosphorylation 2–3 fold in cells expressing GFP. In contrast, CRNK overexpression significantly reduced basal PYK2 phosphorylation, and inhibited the increase in PYK2 phosphorylation induced by ET and H₂O₂. The quantitative analysis of 6–8 Western blotting experiments is depicted in Fig. 2B.

When phosphorylated at Y₄₀₂, PYK2 serves as a scaffolding protein for the PYK2- and Src-dependent tyrosine phosphorylation of PDK1 in vascular smooth muscle cells (VSMC) [13]. PYK2 colocalizes with PDK1 and paxillin in VSMC focal adhesions, and PDK1 phosphorylation is required for downstream activation of

Akt in response to angiotensin II [13]. A similar PYK2-dependent signaling pathway is required for phenylephrine (PE)-induced activation of Akt in NRVM, although this pathway appears to be tonically inhibited by novel PKCs in cardiomyocytes [6]. Therefore, we examined whether CRNK overexpression inhibited ET- or H₂O₂-induced Akt activation in NRVM. As seen in Fig. 3A, Akt was phosphorylated at S₄₇₃ under basal conditions in NRVM infected with Adv-GFP. H₂O₂ (but not ET) significantly increased Akt phosphorylation 2–3 fold. In contrast, CRNK overexpression reduced basal Akt phosphorylation, and also reduced Akt phosphorylation in ET- and H₂O₂-treated NRVM. The quantitative analysis of 6 Western blotting experiments is depicted in Fig. 3B.




Fig. 4. CRNK overexpression alters hypertrophic gene expression in NRVM. In Panel A, NRVM were maintained in control medium (UI), or infected (5 moi, 24 h) with either Adv-GFP or Adv-CRNK. Cells were then left untreated, or treated with phorbol myristate acetate (PMA, 200 nM) for an additional 24 h. Total protein and DNA were analyzed in cell extracts following perchloric acid precipitation. Data are the mean \pm SEM for 7 experiments. * P < 0.05 vs. untreated cells in each group. In Panel B, NRVM were infected (5 moi, 24 h) with either Adv-GFP or Adv-CRNK. Cell extracts were then analyzed for SERCA2, ANF, α MHC, and β MHC mRNA levels (relative to GAPDH mRNA) by real-time RT-PCR. * P < 0.05 vs. Adv-GFP infected cells for each gene of interest.




Fig. 5. Adenoviral gene transfer of nuclear-encoded β -galactosidase into normal hearts. Adv-ne β gal ($\sim 10^{10}$ pfu) was injected by an Optison-mediated gene transfer technique into the aortic root of anesthetized rats in order to define the regions of myocardium transduced by the procedure. Following glutaraldehyde perfusion of whole hearts 3–4 days after gene transfer, tissue sections (14 μ m thick) were processed to detect β -galactosidase activity by X-gal staining. As is evident from this section of the LV posterior wall, scattered regions of nuclear β -galactosidase expression was detected throughout the full thickness of the myocardium. A high power image (box and inset) indicated that both cardiomyocytes and vascular endothelial cells expressed the transgene. Bar = 25 μ m.

3.3. CRNK overexpression alters hypertrophic gene expression in NRVM

PYK2 has been implicated in signaling various aspects of pathological hypertrophy in cultured cardiomyocytes and the intact heart, including alterations in gene expression, increases in cell size and protein synthesis, the induction of apoptosis, and the production of reactive oxygen species in response to PE and ET stimulation [2–6,9,31–33]. Therefore, we tested whether CRNK overexpression prevented NRVM hypertrophy (as measured by total protein/DNA ratio) or altered the hypertrophic gene program in these cultured cells. As seen in Fig. 4A, CRNK overexpression (5 moi, 48 h) had no effect on total protein/DNA ratio as compared to uninfected NRVM, or NRVM infected with Adv-GFP. CRNK also did not prevent the increase in protein/DNA ratio elicited by treatment with phorbol myristate acetate (PMA; 200 nM, 48 h). However, CRNK overexpression increased SERCA2 mRNA levels, in keeping with PYK2's role in regulating SERCA2 gene transcription in NRVM (Fig. 4B) [5]. Furthermore, CRNK significantly reduced ANF mRNA levels, suggesting a role for PYK2 in mediating the PKC and/or Ca²⁺ dependence of ANF gene transcription in NRVM [33,34]. CRNK overexpression also appeared to affect MHC gene expression, although the observed increase in α MHC mRNA levels did not achieve statistical significance.

3.4. Adenoviral gene transfer of GFP and CRNK into normal hearts

Having determined that CRNK overexpression resulted in inhibition of PYK2-dependent signal transduction in cultured cardiomyocytes, we used a modification of a catheter-based, Optison[®]-mediated gene transfer technique [30] to introduce

adenoviral vectors into normal adult ventricular myocardium. First, we used Adv-ne β gal to define the regions of myocardium transduced by the Optison-mediated gene transfer technique. As seen in Fig. 5, injection of $\sim 10^{10}$ pfu of Adv-ne β gal into the aortic root of anesthetized Sprague–Dawley rats produced scattered regions of nuclear β -galactosidase expression 3–4 days after gene transfer. Transgene expression occurred throughout the

anterior as well as posterior walls of the normal LV. High power images indicated that both cardiomyocytes and vascular endothelial cells expressed the transgene. Then, $\sim 10^{10}$ pfu of Adv-GFP ($n=16$) or Adv-CRNK ($n=12$) was injected using the identical catheter-based technique. Five days after gene transfer of Adv-GFP, peak fluorescence in the myocyte (α -actinin positive) and non-myocyte (α -actinin negative) cell populations was assessed




Fig. 6. Adenoviral gene transfer of GFP and CRNK in normal hearts. Peak fluorescence of GFP in the myocyte and non-myocyte cell populations was assessed by flow cytometry in ventricular cells isolated from an uninfected heart, and a GFP-expressing heart 5 days following adenoviral gene transfer. Myocytes (α -actinin positive cells) showed a large shift in peak GFP fluorescence in the cells isolated from the Adv-GFP infected heart as compared to the cells from the uninfected heart (Panel B). Interestingly, in the non-myocyte population (α -actinin negative), there was also a significant increase in the peak fluorescence in cells isolated from the GFP heart (Panel A). In Panel C, freshly isolated cells were examined by phase contrast microscopy (left) and by epifluorescent microscopy (right). In Panel D, ventricular tissue homogenates derived from the LV apex of 3 rats infected with Adv-GFP and 3 rats infected with Adv-CRNK were compared by SDS-PAGE and Western blotting 1 wk after gene transfer. In Panels E and F, pre-operative and 7-day post-operative echo-derived LV/body weight ratios and LV fractional shortening were compared in animals infected with Adv-GFP ($n=16$) and animals infected with Adv-CRNK ($n=12$). * $P<0.05$ for pre- vs. post-operative measurements for each treatment group.




Fig. 7. Alterations in LV structure, function and gene expression during post-myocardial infarction LV remodeling. In Panel A, representative M-mode echocardiograms from Sham-operated and Infarct animals 5 days after surgery are depicted. In Panel B, echo-derived LV/body weight ratios, LV systolic dimensions, LV diastolic dimensions, and LV fractional shortening measurements from Sham-operated ($n=39$) and Infarct ($n=49$) animals at 5 days and 5 wk after surgery are depicted. In Panel C, LV tissue extracts from Sham-operated ($n=11$) and Infarct ($n=8$) animals 5 wks after surgery were analyzed for SERCA2, ANF, α MHC, and β MHC mRNA levels (relative to 18S rRNA) by real-time RT-PCR. * $P<0.05$ vs. Sham-operated for each gene of interest.




Fig. 8. Distribution of nuclear-encoded β -galactosidase following gene transfer into rats following coronary artery ligation. Optison-mediated gene transfer of Adv-ne β gal ($\sim 10^{10}$ pfu) was performed 7 days after ligation of the left anterior descending coronary artery. After removing the heart 3 days later, perfusion-fixation, tissue sectioning ($14\ \mu\text{m}$ thick sections), and X-gal staining was performed to identify regions of myocardium transduced by the procedure. As is evident from the figure, X-gal staining revealed scattered nuclear β -galactosidase activity throughout the non-infarcted septum and posterior wall, and at the border zone of this large, anterolateral wall infarct. No transgene was detected within the infarcted tissue; however, rare, nuclear β -galactosidase activity was detected in the rim of viable, epicardial muscle immediately adjacent to the infarct. Bar= $25\ \mu\text{m}$.

by flow cytometry, and compared to the background fluorescence of cells isolated from an uninfected heart. Myocytes showed a large shift in peak fluorescence in the cells isolated from the Adv-GFP infected LV myocardium (Fig. 6A). Interestingly, we also observed a significant shift in the peak fluorescence in the α -

actinin negative, non-myocyte population following Adv-GFP gene transfer (Fig. 6B), again indicating that both myocytes and nonmyocytes were infected and expressing the transgene. Phase-contrast and epifluorescent microscopy of freshly isolated cells 5d after gene transfer revealed that ~ 5 – 10% of the adult myocytes

were expressing GFP (Fig. 6C). LV tissue homogenates from 3 Adv-GFP and 3 Adv-CRNK infected hearts were then analyzed for CRNK expression by Western blotting. As seen in Fig. 6D, only trace amounts of CRNK were found in LV tissue samples derived from animals receiving Adv-GFP 1 wk after gene transfer.

However, LV CRNK protein levels were substantially increased in animals receiving Adv-CRNK. Transgene expression was readily detectable by Western blotting in the septum, and LV and RV free walls at both the apex and base of the heart. Furthermore, CRNK overexpression was still detectable up to 2 wk after gene




Fig. 9. Adenoviral gene transfer of Adv-GFP and Adv-CRNK after myocardial infarction. In Panel A, the gene transfer protocol is schematically outlined. Rats were subjected to coronary artery ligation and gene transfer with $\sim 10^{10}$ pfu of either Adv-GFP ($n=34$) or Adv-CRNK ($n=28$). Surviving animals were then subjected to general anesthesia, echocardiography, and euthanasia at either 2 or 4 wks following infarct surgery (i.e., 1 or 3 wks after gene transfer). In Panel B, survival curves for Adv-GFP and Adv-CRNK animals were generated by the method of Kaplan and Meier, and compared using the Logrank test ($P=0.032$ for Adv-GFP vs. Adv-CRNK). In Panel C, LV fractional shortening measurements were compared in Adv-GFP vs. Adv-CRNK infected rats each time point in the gene transfer protocol. In Panel D, the change in fractional shortening between the 1 wk and 2 wk echocardiograms of Adv-GFP and Adv-CRNK infected rats are depicted. * $P<0.05$ vs. Adv-GFP at each time point examined.

transfer, but was undetectable at 3 wk (data not shown). Of note, CRNK overexpression did not appear to affect animal survival. In fact, Adv-CRNK modestly increased LV weight/body weight ratio, and also modestly increased fractional shortening as determined by echocardiography 1 wk after gene transfer (Figs. 6E and F).

3.5. Alterations in LV structure, function and gene expression during post-infarction LV remodeling

Adult male rats were then subjected to either sham surgery ($n=39$) or left anterior descending coronary artery ligation ($n=49$). Approximately 25% of the rats died within the first 24 h after sham or infarct surgery. However, after the immediate post-operative period, all of the sham-operated animals survived, whereas 43 of 49 infarct animals survived the entire 5 wk observation period. M-mode and 2-D echocardiography was performed at 5 days and at 5 wk (i.e., immediately prior to euthanasia of the surviving animals). As seen in Fig. 7A, representative M-mode and 2-D echocardiographic images demonstrated anterior wall thinning and increased systolic and diastolic LV chamber dimensions as early as 5 days after coronary artery ligation. Thereafter, LV end-diastolic dimension progressively increased over time, consistent with post-MI LV remodeling (Fig. 7B). Fractional shortening was also significantly reduced at 5 days, and remained depressed for the 5wk observation period (Fig. 7B). Gene expression analysis conducted using a portion of the non-infarcted LV tissue from a subset of infarct animals (Fig. 7C) demonstrated the typical pattern of hypertrophic gene expression associated with pathological LVH. The significant alterations in SERCA2, ANF, α MHC and β MHC mRNA levels (relative to sham-operated controls) were consistent with the progressive increase in echo-derived LV/body weight ratio following coronary artery ligation (Fig. 7B).

3.6. Distribution of transgene following gene transfer into rats with coronary artery ligation


As seen in Fig. 8, Optison-mediated gene transfer of Adv-ne β gal into rats 3–4 days after coronary artery ligation revealed scattered transgene expression throughout the non-infarcted septum and posterior wall, and at the border zone of the anterolateral wall infarct. No transgene was detected within the infarcted tissue; however, rare, nuclear β -galactosidase activity was detected in the rim of viable, epicardial muscle immediately adjacent to the infarct.

3.7. Adenoviral gene transfer of Adv-GFP and Adv-CRNK after myocardial infarction

As depicted in the gene transfer protocol schematically outlined in Fig. 9A, another group of rats was then subjected to coronary artery ligation, and 1 wk later, received either Adv-GFP ($n=34$) or Adv-CRNK ($n=28$). Surviving animals were then subjected to general anesthesia, echocardiography, and euthanasia at either 2 or 4 wk following infarct surgery (i.e., 1 or 3 wk after gene transfer). Survival curves for Adv-GFP and Adv-

CRNK animals were generated by the method of Kaplan and Meier, and compared using the Logrank test. As seen in Fig. 9B, animals receiving Adv-CRNK demonstrated a significant survival advantage over animals receiving Adv-GFP, especially within the first 7 days after gene transfer. The improved survival of Adv-CRNK animals was associated with significantly improved LV fractional shortening at both 2 and 4 wk after infarct surgery (i.e., 1 and 3 wk after gene transfer) (Fig. 9C). Differences in function and survival were not due to differences in initial infarct size between the two groups, as LV fractional shortening at the time of gene transfer were similar ($28.7 \pm 1.9\%$ vs. $28.7 \pm 1.5\%$ for Adv-GFP vs. Adv-CRNK; $P=0.98$). Indeed, fractional shortening in the surviving Adv-GFP infected rats decreased during the first week after gene transfer (i.e., between 1 wk and 2 wk post-MI), whereas animals receiving Adv-CRNK slightly improved ($-6.2 \pm 2.9\%$ vs. $+2.7 \pm 2.6\%$ change in fractional shortening between 1 wk and 2 wk echocardiograms; $P=0.034$; Fig. 9D). Gene expression analysis revealed that the functional improvement in rats receiving Adv-CRNK was associated with a significant decrease in β MHC mRNA levels 1 wk after gene transfer (Fig. 10A). The reduction in β MHC mRNA persisted, and was also associated with a significant increase in α MHC mRNA levels 3 wk after gene transfer as compared to animals receiving Adv-GFP (Fig. 10B).

A. 1 wk after Gene Transfer



B. 3 wk after Gene Transfer




Fig. 10. CRNK overexpression alters hypertrophic gene expression during post-MI ventricular remodeling. LV tissue extracts were analyzed for SERCA2, ANF, α MHC, and β MHC mRNA levels (relative to 18S rRNA) by real-time RT-PCR 1 wk (Panel A) and 3 wk (Panel B) after gene transfer with Adv-GFP or Adv-CRNK. * $P<0.05$ vs. Adv-GFP infected tissue for each gene of interest.

4. Discussion

In this report, adenoviral gene transfer was used to over-express CRNK, the C-terminal domain of PYK2, in cultured cardiomyocytes and in the intact heart. CRNK overexpression reduced basal PYK2 phosphorylation, and interfered with PYK2-dependent activation of Akt in response to the pro-apoptotic agonist H_2O_2 . Furthermore, CRNK overexpression improved LV function and reversed some of the gene expression changes associated with post-MI LV remodeling. Thus, the results of these experiments help to define the role of PYK2-dependent signal transduction in mediating some of the adverse effects of PYK2 activation and overexpression that accompanies LVH and HF [2,4].

Like its homolog FAK, PYK2 contains a C-terminal focal adhesion targeting sequence that directs PYK2 to cardiomyocyte focal adhesions and costameres. Once localized, PYK2 undergoes autophosphorylation at Y₄₀₂, providing a docking site for Src-family protein tyrosine kinases to bind to, and phosphorylate PYK2 at other sites. Thus, PYK2 functions as a scaffold for the assembly of other signaling kinases and adaptor proteins within cardiomyocyte focal adhesions and costameres. In contrast to FAK, PYK2 appears to direct downstream signals to either pro- or anti-apoptotic signaling cascades depending on which molecules bind to the phosphorylated kinase. For instance, Melendez et al. [3] previously demonstrated that overexpression of wildtype PYK2 promoted cardiomyocyte apoptosis, perhaps via the preferential activation of JNKs [3,5,6,32] and direct activation of the mitochondrial apoptotic machinery [35]. PYK2-dependent apoptosis could be suppressed by prior overexpression of paxillin, which binds to the C-terminal domain of PYK2 [14,36] and prevented JNK activation [3]. Similarly, Guo et al. [6] recently demonstrated that PYK2 forms a signaling complex with PDK1 and paxillin, which when formed in the presence of a JNK inhibitor, promoted cell survival via preferential activation of Akt. Our results indicating that CRNK overexpression suppressed H_2O_2 -induced Akt phosphorylation are consistent with their results, as CRNK likely displaced endogenous PYK2 from the complex, and interfered with downstream signaling to Akt. However, CRNK overexpression alone did not appear to induce apoptosis or prevent cardiomyocyte hypertrophy, but rather reversed some of the PYK2-dependent alterations in gene expression observed in pathological LVH and HF [5,33].

Our results confirm earlier studies [15,17–20] and indicate that CRNK indeed functions as a dominant-negative inhibitor of PYK2 in cardiomyocytes. CRNK prevented PYK2 autophosphorylation at Y₄₀₂ in response to both ET and H_2O_2 treatment of NRVM. CRNK, like its homolog FRNK, appears to mediate its inhibitory effect by preventing PYK2 localization and autophosphorylation in response to a variety of extracellular signals. However, CRNK overexpression does not prevent FAK activation, presumably due to C-terminal sequence differences between FAK and PYK2 [15]. Conversely, FRNK overexpression can prevent both FAK and PYK2 activation in a variety of cell types, including cardiomyocytes [23]. Thus it is conceivable that some of the reported effects of FRNK overexpression are mediated by inhibition of PYK2, rather than FAK.

A novel aspect of this report was the modification of a previously described, catheter-based gene transfer method [30] to administer a potentially “therapeutic” adenovirus via selective injection into the coronary circulation. We showed that this catheter-based technique caused adenovirally mediated transgene expression in both the cardiomyocyte and nonmuscle cell populations of the rat heart, and induced the robust expression of a potent, polypeptide inhibitor of an important signaling kinase. CRNK overexpression clearly preserved LV function, and provided a significant survival advantage over animals receiving Adv-GFP, especially within the first 7 days after gene transfer. In fact, we believe that the beneficial effects of CRNK overexpression may have been underestimated, as a substantial number of animals receiving Adv-GFP did not survive and were therefore excluded from subsequent analysis of cardiac structure and function.

Nevertheless, the mechanism(s) responsible for this survival advantage remain unclear. Unlike other agents, CRNK overexpression did not appear to prevent the progressive LV dilatation associated with post-MI ventricular remodeling, as LV end-diastolic dimension was not significantly different between animals receiving Adv-GFP or Adv-CRNK at any time point in the gene transfer protocol (data not shown). Although β MHC mRNA levels were significantly reduced within 1 wk of CRNK gene transfer, it is unclear whether this was the effect of, rather than the cause of the improved LV function. Furthermore, the significant increase in α MHC mRNA was observed at time when transgene expression was undetectable (i.e., 3 wks after gene transfer). As CRNK was overexpressed long after the onset of myocardial necrosis, it seems unlikely that the transgene prevented further ischemic injury by augmenting signaling to cardiomyocyte cell survival pathways [6]. As indicated above, CRNK overexpression altered the hypertrophic gene program in the hemodynamically overloaded, non-infarcted LV myocardium. This hemodynamic overload likely stimulated integrin-dependent signal transduction at costameric sites within the cardiomyocyte population of the heart [37]. However, it is also conceivable that the inhibitor blocked some of the adverse effects of the gene transfer technique that occurred in the coronary endothelium. van Buul et al. [20] have shown that CRNK overexpression prevented the PYK2-dependent, tyrosine phosphorylation of vascular endothelial (VE)-cadherin in response to the addition of a VE-cadherin-blocking blocking antibody (c175) added to the medium of cultured human vascular endothelial cells. CRNK also markedly attenuated the increase in endothelial permeability induced by c175. As increased vascular permeability is required for efficient adenoviral gene transfer [38], CRNK overexpression in coronary microvascular endothelial cells may have promoted the restoration of barrier function, and improved coronary perfusion during the post-gene transfer period. It is also conceivable that the gene transfer technique itself induced cardiomyocyte apoptosis, and CRNK (but not GFP) reduced the loss of muscle cells, thus accounting for its beneficial effects on fractional shortening and survival. However, neither Beeri et al. [30] nor Bekeredjian et al. [39] reported any significant structural or functional effects of the microbubble-ultrasound

technique in delivering plasmids or adenoviruses via coronary artery injection, other than an increase in myocardial stiffness. Additional studies will be necessary, however, to determine the primary site of action of CRNK, and to ascertain the mechanisms responsible for PYK2-dependent deterioration of LV performance post-MI.

Acknowledgments

These studies were supported by NIH RO1 grant HL63711, and a grant to the Cardiovascular Institute from the Ralph and Marian Falk Trust for Medical Research.

References

- [1] Cohn JN, Ferrari R, Sharpe N. Cardiac remodeling-concepts and clinical implications: a consensus paper from an international forum on cardiac remodeling. Behalf of an International Forum on Cardiac Remodeling. *J Am Coll Cardiol* 2000;35:569–82.
- [2] Bayer AL, Heidkamp MC, Patel N, Porter MJ, Engman SJ, Samarel AM. PYK2 expression and phosphorylation increases in pressure overload-induced left ventricular hypertrophy. *Am J Physiol Heart Circ Physiol* 2002;283:H695–706.
- [3] Melendez J, Turner C, Avraham H, Steinberg SF, Schaefer E, Sussman MA. Cardiomyocyte apoptosis triggered by RAFTK/pyk2 via Src kinase is antagonized by paxillin. *J Biol Chem* 2004;279:53516–23.
- [4] Melendez J, Welch S, Schaefer E, Moravec CS, Avraham S, Avraham H, et al. Activation of pyk2/related focal adhesion tyrosine kinase and focal adhesion kinase in cardiac remodeling. *J Biol Chem* 2002;277:45203–10.
- [5] Heidkamp MC, Scully BT, Vijayan K, Engman SJ, Szotek EL, Samarel AM. PYK2 regulates SERCA2 gene expression in neonatal rat ventricular myocytes. *Am J Physiol Cell Physiol* 2005;289:C471–82.
- [6] Guo J, Sabri A, Elouardighi H, Rybin V, Steinberg SF. α 1-adrenergic receptors activate AKT via a Pyk2/PDK-1 pathway that is tonically inhibited by novel protein kinase C isoforms in cardiomyocytes. *Circ Res* 2006;99:1367–75.
- [7] Avraham H, Park SY, Schinkmann K, Avraham S. RAFTK/Pyk2-mediated cellular signalling. *Cell Signal* 2000;12:123–33.
- [8] Sabri A, Govindarajan G, Griffin TM, Byron KL, Samarel AM, Lucchesi PA. Calcium- and protein kinase C-dependent activation of the tyrosine kinase PYK2 by angiotensin II in vascular smooth muscle. *Circ Res* 1998;83: 841–51.
- [9] Bayer AL, Ferguson AG, Lucchesi PA, Samarel AM. Pyk2 expression and phosphorylation in neonatal and adult cardiomyocytes. *J Mol Cell Cardiol* 2001;33:1017–30.
- [10] Bayer AL, Heidkamp MC, Howes AL, Heller Brown J, Byron KL, Samarel AM. Protein kinase C ϵ -dependent activation of proline-rich tyrosine kinase 2 in neonatal rat ventricular myocytes. *J Mol Cell Cardiol* 2003;35:1121–33.
- [11] Blaukat A, Ivankovic-Dikic I, Gronroos E, Dolfi F, Tokiwa G, Vuori K, et al. Adaptor proteins Grb2 and Crk couple Pyk2 with activation of specific mitogen-activated protein kinase cascades. *J Biol Chem* 1999;274: 14893–901.
- [12] Pandey P, Avraham S, Kumar S, Nakazawa A, Place A, Ghanem L, et al. Activation of p38 mitogen-activated protein kinase by PYK2/related adhesion focal tyrosine kinase-dependent mechanism. *J Biol Chem* 1999;274:10140–4.
- [13] Taniyama Y, Weber DS, Roci P, Hilenski L, Akers ML, Park J, et al. Pyk2- and Src-dependent tyrosine phosphorylation of PDK1 regulates focal adhesions. *Mol Cell Biol* 2003;23:8019–29.
- [14] Xiong WC, Macklem M, Parsons JT. Expression and characterization of splice variants of PYK2, a focal adhesion kinase-related protein. *J Cell Sci* 1998;111:1981–91.
- [15] Li X, Dy RC, Cance WG, Graves LM, Earp HS. Interactions between two cytoskeleton-associated tyrosine kinases: calcium-dependent tyrosine kinase and focal adhesion tyrosine kinase. *J Biol Chem* 1999;274:8917–24.
- [16] Hayasaka H, Martin KH, Hershey ED, Parsons JT. Disruption of FRNK expression by gene targeting of the intronic promoter within the focal adhesion kinase gene. *J Cell Biochem* 2007;102:947–54.
- [17] Park JG, Bose A, Leszyk J, Czech MP. PYK2 as a mediator of endothelin-1/ $G\alpha_{11}$ signaling to GLUT4 glucose transporters. *J Biol Chem* 2001;276:47751–4.
- [18] Sorokin A, Kozlowski P, Graves L, Philip A. Protein-tyrosine kinase Pyk2 mediates endothelin-induced p38MAPK activation in glomerular mesangial cells. *J Biol Chem* 2001;276:21521–8.
- [19] Watson JM, Harding TW, Golubovskaya V, Morris JS, Hunter D, Li X, et al. Inhibition of the calcium-dependent tyrosine kinase (CADTK) blocks monocyte spreading and motility. *J Biol Chem* 2001;276:3536–42.
- [20] van Buul JD, Anthony EC, Fernandez-Borja M, Burridge K, Hordijk PL. Proline-rich tyrosine kinase 2 (Pyk2) mediates vascular endothelial-cadherin-based cell–cell adhesion by regulating β -catenin tyrosine phosphorylation. *J Biol Chem* 2005;280:21129–36.
- [21] Samarel AM, Engelmann GL. Contractile activity modulates myosin heavy chain- β expression in neonatal rat heart cells. *Am J Physiol* 1991;261: H1067–77.
- [22] Westfall MV, Rust EM, Albayya F, Metzger JM. Adenovirus-mediated myofibrillar gene transfer into adult cardiac myocytes. *Methods Cell Biol* 1997;52:307–22.
- [23] Heidkamp MC, Bayer AL, Kalina JA, Eble DM, Samarel AM. GFP-FRNK disrupts focal adhesions and induces anoikis in neonatal rat ventricular myocytes. *Circ Res* 2002;90:1282–9.
- [24] Eble DM, Strait JB, Govindarajan G, Lou J, Byron KL, Samarel AM. Endothelin-induced cardiac myocyte hypertrophy: role for focal adhesion kinase. *Am J Physiol Heart Circ Physiol* 2000;278:H1695–707.
- [25] Schlaepfer DD, Hunter T. Evidence for in vivo phosphorylation of the Grb2 SH2-domain binding site on focal adhesion kinase by Src-family protein-tyrosine kinases. *Mol Cell Biol* 1996;16:5623–33.
- [26] Strait JB, Martin JL, Bayer A, Mestrlil R, Eble DM, Samarel AM. Role of protein kinase C- ϵ in hypertrophy of cultured neonatal rat ventricular myocytes. *Am J Physiol Heart Circ Physiol* 2001;280:H756–66.
- [27] Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 1987;162:156–9.
- [28] Porter MJ, Heidkamp MC, Scully BT, Patel N, Martin JL, Samarel AM. Isoenzyme-selective regulation of SERCA2 gene expression by protein kinase C in neonatal rat ventricular myocytes. *Am J Physiol Cell Physiol* 2003;285:C39–47.
- [29] Pfeffer JM, Pfeffer MA, Braunwald E. Influence of chronic captopril therapy on the infarcted left ventricle of the rat. *Circ Res* 1985;57:84–95.
- [30] Beeri R, Guerrero JL, Supple G, Sullivan S, Levine RA, Hajjar RJ. New efficient catheter-based system for myocardial gene delivery. *Circulation* 2002;106:1756–9.
- [31] Kodama H, Fukuda K, Takahashi T, Sano M, Kato T, Tahara S, et al. Role of EGF receptor and Pyk2 in endothelin-1-induced ERK activation in rat cardiomyocytes. *J Mol Cell Cardiol* 2002;34:139–50.
- [32] Kodama H, Fukuda K, Takahashi E, Tahara S, Tomita Y, Ieda M, et al. Selective involvement of p130Cas/Crk/Pyk2/c-Src in endothelin-1-induced JNK activation. *Hypertension* 2003;41:1372–9.
- [33] Hirotsu Y, Higuchi Y, Nishida K, et al. Ca²⁺-sensitive tyrosine kinase Pyk2/CAK β -dependent signaling is essential for G-protein-coupled receptor agonist-induced hypertrophy. *J Mol Cell Cardiol* 2004;36: 799–807.
- [34] McDonough PM, Stella SL, Glembocki CC. Involvement of cytoplasmic calcium and protein kinases in the regulation of atrial natriuretic factor secretion by contraction rate and endothelin. *J Biol Chem* 1994;269: 9466–72.
- [35] Aoki H, Kang PM, Hampe J, Yoshimura K, Noma T, Matsuzaki M, Izumo S. Direct activation of mitochondrial apoptosis machinery by c-Jun N-terminal kinase in adult cardiac myocytes. *J Biol Chem* 2002;277: 10244–50.
- [36] Schaller MD, Sasaki T. Differential signaling by the focal adhesion kinase and cell adhesion kinase β . *J Biol Chem* 1997;272:25319–25.
- [37] Domingos PP, Fonseca PM, Nadruz Jr W, Franchini KG. Load-induced focal adhesion kinase activation in the myocardium: role of stretch and contractile activity. *Am J Physiol Heart Circ Physiol* 2002;282:H556–64.
- [38] Hajjar RJ, del Monte F, Matsui T, Rosenzweig A. Prospects for gene therapy for heart failure. *Circ Res* 2000;86:616–21.
- [39] Bekeredjian R, Chen S, Frenkel PA, Grayburn PA, Shohet RV. Ultrasound-targeted microbubble destruction can repeatedly direct highly specific plasmid expression to the heart. *Circulation* 2003;108:1022–6.