




Official Journal of the European Paediatric Neurology Society



Case study

Clinical and biochemical features in a Congolese infant with congenital disorder of glycosylation (CDG)-IIx

N.C. Nsibu^a, J. Jaeken^{b,*}, H. Carchon^b, M. Mampunza^c, L. Sturiale^d, D. Garozzo^d,
M.N.L. Mashako^a, M.P. Tshibassu^a

^aDepartment of Pediatrics, University of Kinshasa, Democratic Republic of the Congo

^bDepartment of Pediatrics, Center for Metabolic Disease, Katholieke Universiteit Leuven, Herestraat 49, BE 3000 Leuven, Belgium

^cDepartment of Psychiatry, University of Kinshasa, Democratic Republic of the Congo

^dCNR-Istituto per la Chimica e la Tecnologia dei Polimeri, Catania, Italy

ARTICLE INFO

Article history:

Received 14 May 2007

Received in revised form

23 July 2007

Accepted 26 July 2007

Keywords:

CDG-II

Congenital disorder of glycosylation

Congolese

MALDI-TOF

N-glycan

Transferrin isoelectrofocusing

ABSTRACT

We describe an infant girl with psychomotor retardation, growth retardation, mild facial dysmorphism, evidence of liver involvement and a type 2 pattern of serum sialotransferrins. Serum transferrin glycan analysis with MALDI-TOF showed an extremely altered N-glycan pattern with a large number of truncated asialoglycans pointing to a severely defective N-glycan processing. The basic defect in this patient with CDG-IIx has not yet been identified. © 2007 European Paediatric Neurology Society. Published by Elsevier Ltd. All rights reserved.

1. Introduction

Congenital disorders of glycosylation (CDG) are a large family of genetic diseases due to decreased or increased glycosylation of glycoconjugates.^{1–3} Some 29 defects have been identified since Jaeken et al. reported the first patients in 1980.⁴ There are four main groups of CDG: diseases of protein N-glycosylation, of protein O-glycosylation, of combined N- and O-glycosylation, and of lipid glycosylation. N-glycosylation defects can be due to defective assembly (cytosolic and endoplasmic reticulum (ER) defects) or to defective processing

(ER and Golgi defects). Clinical features of the different CDG types are heterogeneous and range from severe to mild, and from multisystem to mono-organ involvement, but neurological manifestations are mostly predominant.

Isoelectrofocusing (IEF) of serum transferrin (Tf) is the method of choice to screen for N-glycosylation defects accompanied by deficiency of sialic acid, a negatively charged sugar. The abnormal IEF patterns show a cathodal shift and can be grouped into two types: type 1 with a decrease of tetrasialoTf and an increase of disialo- and asialoTf pointing to an assembly defect, and type 2 with an increase also of the

*Corresponding author. Tel.: +32 16 34 38 20.

E-mail address: jaak.jaeken@uz.kuleuven.be (J. Jaeken).

uneven bands trisialo- and/or monosialoTf, pointing to a processing defect. In the latter case, intermediate and/or abnormal glycans are accumulating, and matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) is an important tool in the elucidation of these glycan structures and in the elaboration of a hypothesis about the basic defect.^{5,6}

Although most reported patients are Caucasians, patients with very diverse ethnic origins have been described.^{7–19} We believe that CDG are underdiagnosed by African clinicians, as their clinical phenotypes can bare similarity to diseases more common in tropical regions.

We report the first black African child with a CDG. Clinical and biochemical data point to CDG-II but the precise defect has not been elucidated.

2. Patient report

This Congolese girl was evaluated at 10 months of age for hypotonia and failure to thrive. She was born to a 37-year-old mother and a 46-year-old father after a 39 weeks uneventful gestation with normal Apgar score. Her birth weight was 3.900 kg (95th percentile), birth length was 50 cm (50th percentile) and head circumference was 35 cm (50th percentile). She was discharged at the age of 10 days after neonatal



Fig. 1 – Patient at the age of 16 months. Note hypertelorism and open mouth.

hyperbilirubinemia and readmitted at the age of 6 months for severe pneumonia treated with Erythromycin.

Physical examination at 10 months revealed a weight of 7 kg (3rd percentile), length of 68 cm (3rd percentile) and head circumference of 42.5 cm (3rd percentile). She had axial hypotonia, internal strabismus, hypertelorism, persistently open mouth with tongue protusion, and low-set ears. Her abdomen was soft with firm hepatomegaly (3 cm below the costal margin). She had a developmental delay (psychomotor level of about 2.5 months) and had tic-like head movements. Cardiovascular examination was normal. Skeletal radiography and echocardiography were normal. Electroencephalography showed bilateral temporal waves consistent with epileptiform activity.

The patient continued to show a growth retardation after the first year (at 15 months of age: weight 8 kg (<3rd percentile = 8.9); length 70 cm (<3rd percentile = 76); head circumference 43.5 cm (<3rd percentile = 45) (Fig. 1).




Fig. 2 – Serum Tf IEF pattern in a control (a), in the patient (b) and in three other patients with different type 2 IEF patterns ((c, e): CDG-IIx, (d): CDG-IIa). Numbers on the left indicate sialoTf isoforms.




Fig. 3 – MALDI-TOF analysis of intact serum Tf of the patient. The 79616.2 peak represents normal Tf, while the 78232.9 peak is abnormal Tf representing probably truncated asialoTf.

Blood cell count including platelets, as well as glycemia, ammonia levels and serum creatine kinase (CK) were normal. Serum bilirubin (7 mg/dl), glutamate-oxaloacetate transaminase (92 U/l) and glutamate-pyruvate transaminase (80 U/l) were increased while lactate dehydrogenase and albumin were decreased (76.9 U/l; reference range: 228–456 and 2.81 g/dl; Nl: 3.1–3.5, respectively). Thyroid function (T3, T4, TSH) and routine CSF analysis were normal. Coagulation studies, hepatic ultrasound and cerebral computed tomography were not performed.

A fibroblast culture from a skin biopsy was unsuccessful.

She was readmitted 5 months later with severe respiratory distress, seizures, pallor and jaundice resulting from a *Klebsiellae pneumoniae* infection. She died with a gastric bleeding after a week of intensive care at the age of 19 months. The parents refused a new skin biopsy.

3. Methods

IEF of serum Tf²⁰ and of apo C-III²¹ were performed as described.




Fig. 4 – (a) MALDI-TOF analysis of permethylated Tf glycans in positive reflector mode in patient: ■: N-acetylglucosamine; ●: mannose; ●: galactose; ◆: sialic acid; ▼: fucose. Note the large number of truncated asialoglycans (including a GlcNAc₂Man₃ glycan (boxed)) compared to control (b).

The N-glycans of Tf were released by PNGase F treatment. These N-linked glycans were purified by solid-phase extraction and subjected to permethylation in the presence of sodium hydroxide. The permethylated glycans were then analyzed by MALDI-TOF mass spectrometry (MS) in negative and positive ion mode.^{22,23}

The MGAT1 gene was analyzed by direct sequencing.

4. Results

Capillary zone electrophoresis and IEF of serum Tf showed a type 2 pattern with a decrease of tetrasialoTf and increased disialo-, trisialo-, mono- and asialoTf (Fig. 2). IEF of serum Tf after neuraminidase treatment showed only one Tf band excluding a Tf protein variant. IEF of serum apo C-III was normal.

MALDI-TOF analysis of intact serum Tf of the patient showed a prominent abnormal peak suggesting a large amount of truncated asialoglycans (Fig. 3). MALDI-TOF analysis of serum Tf in negative linear mode (not shown) and in positive reflector mode (Fig. 4a and b) confirmed a severely reduced amount of tetrasialoTf and an extremely altered N-glycan pattern with a large number of truncated asialo glycans (including a GlcNAc₂Man₃ glycan), pointing to severely defective glycan processing (Sturiale et al., in preparation).

Mutation analysis of MGAT1 did not show any pathogenic mutation at the level of genomic DNA.

5. Discussion

The clinical syndrome of this patient comprised psychomotor retardation, growth retardation, mild facial dysmorphism and evidence for liver involvement. She showed in addition a type 2 serum sialoTf pattern not due to a Tf protein variant. Since IEF of serum apo C-III was normal these data pointed to an N-glycosylation disorder without evidence for impairment of mucin type O-glycosylation. Serum Tf glycan analysis with MALDI-TOF showed a unique pattern not reported before.

CDG with a type 2 serum sialoTf pattern can be divided in CDG with an identified basic defect and CDG with a not yet identified basic defect. There are only few CDG with a type 2 pattern in which the biochemical defect has been identified (Table 1). Among these, CDG-IIa and CDG-IIId are N-glycosylation defects^{24,25} while the COG7, the COG1 and the COG8 defects are combined N- and O-glycosylation defects.^{13,19,33}

On the other hand, there is a rapidly growing group of reported and unreported CDG patients with a type 2 pattern and a still unclear basic defect (CDG-IIx). Among these are the patients with various combinations of rather aspecific symptoms (aspecific dysmorphism, psychomotor retardation, epilepsy, etc.),^{26–28} and on the other hand, several clinically recognizable groups such as isolated liver disease,^{29,30} cutis laxa syndromes in patients in whom the known cutis laxa defects have been excluded,³¹ and a syndrome with glomerulopathy and buphtalmia.³² The latter syndromes could be excluded in the present patients, and the combined clinical

Table 1 – CDG with a type 2 serum Tf IEF and identified basic defect

Disorder	Clinical features	First reported in (Ref.)
CDG-IIa	Psychomotor retardation, epilepsy, stereotype behaviour, dysmorphism	1994 ²⁴
CDG-IIId	Psychomotor retardation, Dandy-Walker malformation, myopathy, bleedings	2002 ²⁵
COG7 defect	Encephalopathy, growth retardation, liver disease, hyperthermia	2004 ¹⁹
COG1 defect	Mild psychomotor retardation, progressive microcephaly, growth retardation of the rhizomelic type	2006 ¹³
COG8 defect	Psychomotor retardation, neurological abnormalities (epilepsy, ataxia, etc.), mild dysmorphism	2007 ³³

and biochemical data permit us to exclude the elucidated CDG with a type 2 serum sialoTf pattern.

On the basis of the glycan structures, particularly the GlcNAc₂Man₃ glycan, we hypothesized that this patient might have suffered from a deficiency of N-acetylglucosaminyl-transferase 1 coded by MGAT1. However, MGAT1 analysis failed to show pathogenic mutations.

In conclusion, the present patient is the first reported black African with a CDG. The combined clinical and biochemical data point to a novel (primary or secondary?) CDG, belonging to the CDG-II group (processing defect).

Acknowledgments

We thank Prof. Gert Matthijs and Liesbeth Keldermans of the Center for Human Genetics (KULeuven) for DNA analysis.

REFERENCES

- Freeze HH. Genetic defects in the human glycome. *Nat Rev Genet* 2006;7:537–51.
- Jaeken J, Matthijs G. Congenital disorders of glycosylation: a rapidly expanding disease family. *Annu Rev Genomics Hum Genet* 2007, May 21. [epub ahead of print].
- Zeevaert R, Schollen E, Carchon H, Matthijs G, Jaeken J. Congenital disorders of glycosylation (CDG): update and perspectives. *Curr Pediatr* 2006;2:323–30.
- Jaeken J, Vanderschueren-Lodeweyckx M, Casaer P, et al. Familial psychomotor retardation with markedly fluctuating serum prolactin, FSH and GH levels, partial TBG deficiency, increased serum arylsulphatase A and increased CSF protein: a new syndrome? *Pediatr Res* 1980;14:179.
- Kang P, Mechref Y, Klouckova I, Novotny M. Solid-phase permethylation of glycans for mass spectrometric analysis. *Rapid Commun Mass Spectrom* 2005;19:3421–8.

6. Mills PB, Mills K, Mian N, Winchester BB, Clayton PT. Mass spectrometric analysis of glycans elucidating the pathogenesis of CDG type IIx. *J Inherit Metab Dis* 2003;26:119–34.
7. Adamowicz M, Matthijs G, Van Schaftingen E, et al. New case of phosphomannose isomerase deficiency. *J Inherit Metab Dis* 2000;23(Suppl. 1):184.
8. Artigas J, Cardo E, Pineda M, Nosas R, Jaeken J. Phosphomannomutase deficiency and normal pubertal development. *J Inherit Metab Dis* 1998;21:78–9.
9. De Michelena M, Franchi L, Summers P, De La Fuente C, Campos P, Jaeken J. Carbohydrate-deficient glycoprotein syndrome due to phosphomannomutase deficiency: the first reported cases from Latin America. *Am J Med Genet* 1999;84:481–3.
10. Di Rocco M, Barone R, Adam A, et al. Carbohydrate-deficient glycoprotein syndromes: the Italian experience. *J Inherit Metab Dis* 2000;23:391–5.
11. Du Plessis J, Pretorius P, Kholi D, et al. An atypical carbohydrate-deficient glycoprotein (CDG) syndrome patient in South Africa. *S Afr Med J* 2001;91:392–4.
12. Enns G, Steiner R, Buist N, et al. Clinical and molecular features of congenital disorder of glycosylation in patients with type 1 sialotransferrin pattern and diverse ethnic origins. *J Pediatr* 2002;141:695–700.
13. Foulquier F, Vasile E, Schollen E, et al. Conserved oligomeric Golgi complex subunit 1 deficiency reveals a previously uncharacterized congenital disorder of glycosylation type II. *Proc Natl Acad Sci USA* 2006;103:3764–9.
14. Imtiaz F, Worthington V, Champion M, et al. Genotypes and phenotypes of patients in the UK with carbohydrate-deficient glycoprotein syndrome type 1. *J Inherit Metab Dis* 2000;23:162–74.
15. Jaeken J, Matthijs G, Saudubray JM, et al. Phosphomannose isomerase deficiency: a carbohydrate-deficient glycoprotein syndrome with hepatic intestinal presentation. *Am J Hum Genet* 1998;62:1535–9.
16. Ohno K, Yuasa I, Akaboshi S, et al. The carbohydrate deficient glycoprotein syndrome in three Japanese children. *Brain Dev* 1992;14:30–5.
17. Pelletier VA, Galeano N, Brochu P, et al. Secretory diarrhea with protein-losing enteropathy, enterocolitis cystica superficialis, and congenital hepatic fibrosis: a new syndrome. *J Pediatr* 1986;108:61–5.
18. Tayebi N, Andrews D, Park J, et al. A deletion-insertion mutation in the phosphomannomutase 2 gene in an African American patient with congenital disorders of glycosylation-Ia. *Am J Med Genet* 2002;108:241–6.
19. Wu X, Steet R, Bohorov O, et al. Mutations of the COG complex subunit gene COG7 causes a lethal congenital disorder. *Nat Med* 2004;10:518–23.
20. Jaeken J, van Eijk HG, van der Heul C, et al. Sialic acid-deficient serum and cerebrospinal fluid transferrin in a newly recognized genetic syndrome. *Clin Chim Acta* 1984;144:245–7.
21. Wopereis S, Grunewald S, Morava E, et al. Apolipoprotein C-III isofocusing in the diagnosis of genetic defects in O-glycan biosynthesis. *Clin Chem* 2003;49:1839–45.
22. Ciucanu I, Kerek F. A simple and rapid method for the permethylation of carbohydrates. *Carbohydr Res* 1984;131:209–17.
23. Sturiale L, Barone R, Fiumura A, et al. Hypoglycosylation with increased fucosylation and branching of serum transferrin N-glycans in untreated galactosemia. *Glycobiology* 2005;15:1268–76.
24. Jaeken J, Schachter H, Carchon H, et al. Carbohydrate deficient glycoprotein syndrome type II: a deficiency in Golgi localized N-acetyl-glucosaminyltransferase II. *Arch Dis Child* 1994;71:123–7.
25. Hansske B, Thiel C, Lubke T, et al. Deficiency of UDP-galactose: N-acetylglucosamine beta-1,4-galactosyltransferase I causes the congenital disorder of glycosylation type IID. *J Clin Invest* 2002;109:725–33.
26. Cheillan D, Cognat S, Dorche C, et al. CDG-IIx with unusual phenotype. *J Inherit Metab Dis* 2004;27:103–4.
27. Miura Y, Tay SK, Aw MM, Eklund EA, Freeze HH. Clinical and biochemical characterization of a patient with congenital disorder of glycosylation (CDG)IIx. *J Pediatr* 2005;147:851–3.
28. Zentilini Boyer M, De Lonlay P, Seta N, et al. Failure to thrive and intestinal diseases in congenital disorders of glycosylation. *Arch Pediatr* 2003;10:590–5.
29. Butler M, Quelhas D, Critchley AJ, et al. Detailed glycan analysis of serum glycoproteins of patients with congenital disorders of glycosylation indicates the specific defective glycan processing step and provides an insight into pathogenesis. *Glycobiology* 2003;13:601–22.
30. Mandato C, Brive L, Miura Y, et al. Cryptogenic liver disease in four children: a novel congenital disorder of glycosylation. *Pediatr Res* 2006;59:293–8.
31. Wopereis S, Morava E, Coucke P, et al. Defective protein glycosylation in patients with cutis laxa syndrome. *Eur J Hum Genet* 2005;13:414–21.
32. Wilson M, Lewis D, Arbuckle S, et al. A new type II congenital disorder of glycosylation associated with progressive glomerulopathy, retinal dystrophy and neurological abnormalities. *J Inherit Metab Dis* 2003;26(Suppl. 2):128.
33. Foulquier F, Ungar D, Reynders E, et al. A new inborn error of glycosylation due to Cog8 deficiency reveals a critical role for the Cog1–Cog8 interaction in COG complex formation. *Hum Mol Gen* 2007;16:717–30.