

REVIEW

Articular cartilage destruction in experimental inflammatory arthritis: insulin-like growth factor-1 regulation of proteoglycan metabolism in chondrocytes

PERNETTE J. VERSCHURE¹, CORNELIS J. F. VAN NOORDEN², JAN VAN MARLE³ and WIM B. VAN DEN BERG¹

¹Department of Rheumatology, University Hospital Nijmegen, Geert Grooteplein Zuid 8, 6525 GA Nijmegen, The Netherlands

²Academic Medical Center, University of Amsterdam, Laboratory of Cell Biology and Histology, PO Box 22700, 1100 DE Amsterdam, The Netherlands

³Academic Medical Center, University of Amsterdam, Department of Electron Microscopy, PO Box 22700, 1100 DE Amsterdam, The Netherlands

Received 14 February 1996 and in revised form 30 July 1996

Summary

Rheumatoid arthritis, a disease of unknown aetiology, is characterized by joint inflammation and, in its later stages, cartilage destruction. Inflammatory mediators may exert not only suppression of matrix synthesis but also cartilage degradation, which eventually leads to severe cartilage depletion. Systemically and locally produced growth factors and hormones regulate cartilage metabolism. Alterations in levels of these factors or in their activity can influence the pathogenesis of articular cartilage destruction in arthritic joints. The main topic of the present review is the role of the anabolic factor insulin-like growth factor-1 in the regulation of chondrocyte metabolic functions in normal and in diseased cartilage. This is the most important growth factor that balances chondrocyte proteoglycan synthesis and catabolism to maintain a functional cartilage matrix. A brief overview of how chondrocytes keep the cartilage matrix intact, and how catabolic and anabolic factors are thought to be involved in pathological cartilage destruction precedes the review of the role of this growth factor in proteoglycan metabolism in cartilage.

Articular cartilage

Structure and function of articular cartilage

Articular cartilage forms the weight-bearing surface of the articular joint, providing it with a resilient, smoothly gliding surface. Articular cartilage consists of a highly organized extracellular matrix in which specialized cells, chondrocytes, are embedded. Aggregated proteoglycans and collagen fibrils are the two major structural components of the matrix, and form a three-dimensional framework giving the tissue a unique architecture.

Cartilage collagen has a very low basal rate of turnover, and the intact collagen network provides a permanent framework that gives the tissue tensile

strength. Cartilage collagen consists mainly of type II collagen, but cartilage also contains small amounts of type VI, IX and X collagen; type X is restricted to the deep part of the cartilage and type XI is part of the collagen fibril. Moreover, type X collagen is mainly found in hypertrophic chondrocytes or in osteoarthritic cartilage (Reichenberger *et al.*, 1991; Nerlich *et al.*, 1992; Von der Mark *et al.*, 1992; Aigner *et al.*, 1993, 1995).

Proteoglycans are complex macromolecules with a high turnover rate; they consist of a core protein to which one or more glycosaminoglycan side chains or N-O-linked oligosaccharides are attached (Handley *et al.*, 1985). Glycosaminoglycan chains are polymers of repeating disaccharides that can vary in

number of chains, their length as well as their pattern of sulphation; they play a dominant role in the determination of physical properties of the protein to which they are attached. High negatively charged glycosaminoglycans entrap solvent, thus creating a high osmotic pressure which is restrained by collagen. The swelling pressure, together with changed hydraulic permeability, enables cartilage to resist compressive loads with minimal deformation.

The major proteoglycan found in articular cartilage is aggrecan (Hardingham & Fosang, 1992, 1995). Aggrecan molecules contain three globular domains (G1, G2 and G3), of which the G1 domain can bind noncovalently to a single chain of hyaluronan, forming large macromolecular aggregates. This interaction is promoted and stabilized by link protein (Hardingham *et al.*, 1992a) (Fig. 1). Furthermore, cartilage contains nonaggregated small proteoglycans, such as biglycan, decorin and fibromodulin (Witsch-Prehm *et al.*, 1992). These smaller proteoglycans are studied intensively these days, but so far clear information on their function is scant.

An important aspect of the structural integrity of articular cartilage is the ability of chondrocytes to synthesize and degrade proteoglycans to maintain a functionally intact cartilage. Small perturbations in cartilage metabolism may lead to increased or

decreased local concentrations of matrix molecules which can alter physiological properties of the tissue. This may lead to rapid loss of tissue function and deterioration of the joint.

Three strikingly heterogeneous cartilage layers are generally recognized between the articular surface and the subchondral bone, i.e. surface, middle and deeper zones (Fig. 2) (Sampson & Cannon, 1986). Zonal subdivision is based on morphology, differences in cell density, the orientation, nature, content and distribution of proteoglycans, and the organization of the collagen fibrillar network (Maroudas *et al.*, 1990; Siczkowski & Watt, 1990; Aydelotte & Keuttner, 1991; Marles *et al.*, 1991). Chondrocytes in the surface zone are small, flattened, and oriented parallel to the articular surface, and they are rather inert with respect to matrix synthesis. Chondrocytes in the middle zone are larger and round, whereas chondrocytes in the deeper zones have the largest size and occur usually in groups. Chondrocytes in the middle and deeper zones of cartilage exhibit the highest metabolic activity and are responsible for cartilage homeostasis.

Destruction of articular cartilage

During chronic joint disease, such as rheumatoid arthritis, progressive joint inflammation results in irreversible cartilage destruction, leading to erosion




Fig. 1. The basic structural unit of aggrecan, consisting of a core protein to which highly negatively charged glycosaminoglycan chains or oligosaccharides are bound. The core protein is noncovalently attached to hyaluronic acid, and this binding is reinforced by link proteins.




Fig. 2. (A) Schematic drawing of histologically stained human cartilage, showing the three striking heterogeneous cartilage layers that are generally recognized between articular surface and subchondral bone, namely surface (I), middle (II) and deeper zones (III). Zonal subdivision is based on morphology, differences in cell density and orientation, nature, content and distribution of proteoglycans as well as organization of the collagen fibrillar network. (B–D) Schematic drawings of chondrocytes in the three cartilage layers. (B) Chondrocytes in the surface zone are small, flattened, oriented parallel to the articular surface, and are rather inert with respect to matrix synthesis. (C) Chondrocytes in the middle zone are larger and round. (D) Chondrocytes in the deeper zones are largest and usually occur in groups. Chondrocytes in the middle and deeper zones of cartilage exhibit highest metabolic activity and are responsible for cartilage homeostasis. (A, $\times 390$; B $\times 5226$; C, $\times 4680$; D, $\times 5616$).

of articular cartilage, injury of surrounding tissues and eventually permanent loss of joint function (Fig. 3) (Kelly *et al.*, 1989; Harris, 1990).

For the study of processes that determine joint destruction in arthritic joints, animal models have been developed in which arthritis is induced experimentally. Studies performed by our group have revealed that the main aspects of pathological cartilage depletion are increased degradation of articular cartilage, reduced matrix synthesis, and chondrocyte death (Van den Berg *et al.*, 1981; Schalkwijk *et al.*, 1989b). Synovial cells, chondrocytes, and large numbers of macrophages and lymphocytes that infiltrate arthritic joints generate vast quantities of mediators during inflammation,

causing invasion and erosion of cartilage and subsequent cartilage destruction. These mediators include cytokines, proteolytic enzymes and reactive oxygen species. In addition, chronic interference with factors that upregulate synthesis of matrix components in chondrocytes plays an important role in cartilage destruction during joint inflammation (Schalkwijk *et al.*, 1989a, b; Tyler, 1989; Verschure *et al.*, 1989). The importance of a balance between suppressive mediators, such as cytokines, proteolytic enzymes or reactive oxygen species, and anabolic factors, such as circulating hormones and locally produced growth factors for the structural integrity of articular cartilage is emphasized in the following paragraphs.




Fig. 3. Schematic representation of a normal and a rheumatoid arthritic knee joint. Joint inflammation is characterized by chronic synovitis leading to irreversible destruction of the articular cartilage. F = femur, T = tibia, SL = synovial lining, AC = articular cartilage.

Catabolic factors

Cytokines

Cytokines are thought to play an important role in the pathogenesis of rheumatoid arthritis. These signal molecules mediate cell-cell interactions and are able to induce, amplify and maintain inflammation. Cytokines have been implicated in the release of enzymes capable of cartilage destruction (Baici & Lang, 1990; Macnaul *et al.*, 1990; Shapiro *et al.*, 1990; Circolo *et al.*, 1991; Chandrasekhar & Harvey, 1992; Mort *et al.*, 1993; Wahl & Corcoran, 1993), but also have strong effects on metabolic functions of chondrocytes (Holt *et al.*, 1992). High levels of interleukin-1 (IL-1), interleukin-6 (IL-6), interleukin-8 (IL-8), interleukin-10 (IL-10), tumour necrosis factor α (TNF α), granulocyte-macrophage colony stimulating factor and leukocyte inhibitory factor respectively, have been detected in the synovial fluid of arthritic joints; interleukin-2 (IL-2), interleukin-4 (IL-4) and interferon γ are not elevated in arthritic joints (Nouri *et al.*, 1984; Di Giovanna *et al.*, 1988; Houssiau *et al.*, 1988; Saxne *et al.*, 1988; Xu *et al.*, 1989; Brennan *et al.*, 1990; Firestein & Zvaifler, 1990; Waring *et al.*, 1993). However, it must be realized that levels of cytokines in synovial fluid or plasma gives only limited information about their penetration and ultimate effects in the cartilage matrix. Moreover, the final effect of these cytokines is a balance between their concentrations and the concentrations of their specific inhibitors (Hopkins *et al.*, 1988; Dayer & Burger, 1994). Moreover, responses to cytokines are also dependent on the presence of receptors on their target cells.

Despite the complexity of the cytokine network involved in joint inflammation, IL-1 and TNF α are

identified as key mediators in the pathogenesis of rheumatoid arthritis (e.g. Bentzen *et al.*, 1985; Tyler, 1985; Benton & Tyler, 1988; Larrick & Kunkel, 1988; Verschure & Van Noorden, 1990; Henderson *et al.*, 1991; Van den Berg *et al.*, 1991; Elliot *et al.*, 1993; Arend & Dayer, 1995; Van de Loo *et al.*, 1995b; Joosten *et al.*, 1996). In our experimental models of arthritis, the arthritis-inducing stimulus determines whether TNF α drives IL-1 production, or whether IL-1 and TNF α are individually involved (Van den Berg, 1994; Joosten *et al.*, 1995). Moreover, IL-4 and IL-10 are able to interfere with TNF α and IL-1 production and they upregulate expression of their inhibitors, i.e. IL-1 receptor antagonist and soluble TNF receptor (Howard *et al.*, 1992; Wong *et al.*, 1993). IL-1 is a key mediator in the inhibition of chondrocyte proteoglycan synthesis in both antigen- and zymosan-induced models of rheumatoid arthritis (Van de Loo & Van den Berg, 1990; Van de Loo *et al.*, 1992, 1995a). Systemic treatment *in vivo* with anti-IL-1 antibodies or IL-1 receptor antagonist stimulates chondrocyte proteoglycan synthesis (Van de Loo *et al.*, 1992, 1995a; Lethwaite *et al.*, 1994). In the collagen-induced model of rheumatoid arthritis, systemic treatment *in vivo* with anti-IL-1 antibody or IL-1 receptor antagonist not only reduces inhibition of proteoglycan synthesis in chondrocytes, but also reduces cartilage destruction and joint inflammation (Wooley *et al.*, 1993; Van den Berg *et al.*, 1994). Evidence for a direct destructive role of TNF α *in vivo* is still lacking (Van de Loo *et al.*, 1995b; Joosten *et al.*, 1996), but it can induce cartilage damage *in vitro* (Saklatvala, 1986; Wilbrink *et al.*, 1991; Williams *et al.*, 1995). Its main effect in collagen-induced arthritis is probably induction of early joint swelling (Elliot *et al.*, 1993; Joosten *et al.*, 1996).

Proteolytic enzymes

High levels of degradative enzymes can cause breakdown of articular cartilage during joint inflammation. This topic is not discussed here because it has been reviewed extensively elsewhere (Evans, 1991; Firestein, 1992; Buttle, 1994; Henderson & Blake, 1994). Metalloproteinases such as collagenase and stromelysin are important proteolytic enzymes that may be involved in destructive processes (Nguyen *et al.*, 1989; Mort *et al.*, 1993). In addition, cysteine and serine proteinases seem to play a significant role as well (Van Noorden *et al.*, 1988; Maciewicz & Wotton, 1991; Ahmed *et al.*, 1992; Esser *et al.*, 1994). Recently, some doubt was raised about the relevance of the above mentioned proteinases, and it was hypothesized that a newly described enzyme, aggrecanase, could be of more importance (e.g., Flanary & Sandy, 1994; Fosang *et al.*, 1994; Hughes *et al.*, 1995; Lark *et al.*, 1995).

The activity of proteolytic enzymes is not only

determined by the presence of the enzyme molecules but also by post-translational activation processes and interactions of enzymes with their endogenous inhibitors (e.g. Roose & Van Noorden, 1995). In joint inflammation, post-translational activation may be increased, or an imbalance between the levels of matrix degrading enzymes and their inhibitors may occur (e.g. Ellis *et al.*, 1994; Martel-Pelletier *et al.*, 1994).

Reactive oxygen species

In addition to proteolytic breakdown, reactive oxygen species that are released in vast amounts by inflammatory and/or phagocytic cells can cause degradation of the articular cartilage matrix (Roberts *et al.*, 1989; Henrotin & Deby-Dupont, 1992). In previous studies, we and others have shown that *in vitro* exposure of articular cartilage to reactive oxygen causes severe inhibition of chondrocyte proteoglycan synthesis as well as degradation of the aggrecan proteoglycan (Bates *et al.*, 1985; Schalkwijk *et al.*, 1985a; Schalkwijk *et al.*, 1985b). However, the exact role of reactive oxygen species *in vivo* still has to be established. The ultimate effects of reactive oxygen species depends not only on their release but also on levels of endogenous scavenger enzymes and antioxidants (e.g. Schalkwijk *et al.*, 1985a).

At present, nitric oxide (NO), a reactive oxygen species that is produced by NO synthase, is receiving a lot of attention in arthritis research. There is increasing evidence that nitric oxide production contributes to pathological features of joint inflammation (Evans *et al.*, 1995; Sakurai *et al.*, 1995). Nitric oxide production is regulated by cytokines and growth factors (Palmer *et al.*, 1993; Blanco *et al.*, 1995; Lo & Cruz, 1995). The role of nitric oxide has been studied by applying NO synthase antagonists and inhibitors. IL-1 is known to cause inhibition of chondrocyte proteoglycan synthesis through nitric oxide production (Stadler *et al.*, 1991; Charles *et al.*, 1993; Hauselman *et al.*, 1994; Taskiran *et al.*, 1994; Fukuda *et al.*, 1995; Jarvinen *et al.*, 1995). Moreover, nitric oxide is thought to play a regulatory role in activation of metalloproteinases in articular chondrocytes (Murell *et al.*, 1995). However, it is unclear whether nitric oxide is a mediator of cartilage catabolism (Hanglow *et al.*, 1995; Stefanovic-Racic *et al.*, 1996).

Anabolic factors

Rates of chondrocyte proteoglycan synthesis and degradation in articular cartilage are the result of the effects of a number of hormones and locally produced growth factors (Lebovitz & Eisenbarth, 1975; Franchimont & Bassleer, 1991; Hardingham *et al.*, 1992b; Trippel, 1995).

Hormones such as steroids, and growth factors such as basic fibroblast growth factor (bFGF), epidermal growth factor (EGF), platelet-derived growth factor (PDGF), transforming growth factor β (TGF β) and insulin-like growth factor (IGF-1), are known to affect cartilage metabolism. IGF-1 is the most important growth factor that stimulates chondrocyte proteoglycan synthesis and inhibits proteoglycan catabolism (Tyler, 1989). In the following paragraphs, the role of steroid hormones and growth factors in chondrocyte proteoglycan metabolism will be briefly described and subsequently, the role of IGF-1 in cartilage metabolism will be reviewed.

Effect of growth factors and glucocorticoids on chondrocyte proteoglycan metabolism

Growth factors, such as EGF, bFGF, PDGF, TGF β and IGF-1, play a role in the regulation of chondrocyte proteoglycan synthesis and degradation. They are also important in repair processes during joint diseases (Cuevas *et al.*, 1988; Ogawa *et al.*, 1991; Harvey *et al.*, 1993). Controversy exists about their relative importance in modulating chondrocyte proteoglycan metabolism. It must be realized that most information has been gained from studies on isolated chondrocytes or from cartilage explants. Differences in experimental culture conditions, the origin and species from which chondrocytes or cartilage are obtained are all important factors that might determine the outcome of the experiments (Kato & Gospodarowicz, 1985b; Rosier *et al.*, 1988; Burton-Wuster & Lust, 1990; Hiraki *et al.*, 1990). Little is known of the role of growth factors in anatomically intact cartilage (Van der Kraan *et al.*, 1992a).

bFGF and EGF stimulate proteoglycan synthesis strongly in actively growing growth-plate chondrocytes, but only to a limited extent in chondrocytes that have become hypertrophic (Kato & Iwamoto, 1990; Iwamoto *et al.*, 1991; Trippel *et al.*, 1993; Sah *et al.*, 1994; Wroblewski & Edwall-Arvidsson, 1995). PDGF stimulates proteoglycan synthesis to a similar extent as IGF-1, whereas TGF β exerts both stimulatory and inhibitory effects on chondrocyte proteoglycan synthesis (Van der Kraan *et al.*, 1992a, b; Morales, 1994). Furthermore, *in vivo* experiments showed that repeated intra-articular injections of TGF β into mouse knee joints induce strong and long-lasting stimulation of chondrocyte proteoglycan synthesis (Van Beuningen *et al.*, 1993, 1994). Synergistic interactions between growth factors have been described as well in *in vitro* culture experiments (Inoue *et al.*, 1989; Nataf *et al.*, 1990; O'Keefe *et al.*, 1994). For example, IGF-1 synergistically enhances chondrocyte proteoglycan synthesis induced by bFGF or EGF in cultured chondrocytes (Kato *et al.*, 1983).

Glucocorticoids are involved in the regulation of long-term metabolic functions of chondrocytes, but their effects on chondrocyte proteoglycan metabolism remain controversial. Intra-articular injections of glucocorticoids suppress joint inflammation and result in pain relief and marked preservation of joint function (Hollander *et al.*, 1951; Holden & Hume, 1962; Standberg, 1964; Gray & Gottlieb, 1983; Weiss *et al.*, 1990). Low levels of glucocorticoids stimulate chondrocyte proteoglycan synthesis *in vitro* (Kato & Gospodarowicz, 1985a; Takano *et al.*, 1985; Takigawa *et al.*, 1988; Bellows *et al.*, 1989). They also have chondroprotective effects on proteoglycan-depleted cartilage *in vivo*, by decreasing both inhibition of proteoglycan synthesis, and loss of chondrocyte proteoglycan and osteophyte formation (Williams & Brandt, 1985; Joosten *et al.*, 1990; Van den Berg, 1991, 1992). Glucocorticoids, such as cortisone, prednisolone, hydrocortisone, dexamethasone and triamcinolone acetonide also exert inhibitory effects on chondrocyte proteoglycan synthesis in normal cartilage *in vitro* as well as *in vivo* (Mankin & Conger, 1966; Anastassiades & Dziewiatkowski, 1970; Behrens *et al.*, 1975; Weiss *et al.*, 1988; Barrueco *et al.*, 1989; Chunekamrai *et al.*, 1989; Anfeheld, 1992). IGF-1 and glucocorticoids have synergistic effects on chondrocytes (Itagane *et al.*, 1991). The effects of glucocorticoids on chondrocytes may be direct or via regulation of production of inflammatory mediators (Blondelon *et al.*, 1980; Sheppard *et al.*, 1982; Steinberg & Sledge, 1983; Silbermann & Maor, 1985; Shinmei *et al.*, 1988; Di Battista *et al.*, 1991; Itagane *et al.*, 1991; Price *et al.*, 1992; Schlaghecke *et al.*, 1992).

Insulin-like growth factor (IGF)

IGFs, which were originally defined as somatomedins, have been reported to be: a growth hormone-dependent serum factor able to stimulate ³⁵S-sulphate incorporation into cartilage *in vivo* ('sulphation factor'; Salmon & Daughaday, 1957), a cofactor extracted from serum able to stimulate replication of cells ('multiple stimulating activity'; Kato *et al.*, 1981), or a factor in serum exerting insulin-like activity on target tissues ('nonsuppressible insulin-like activity'; Zapf *et al.*, 1987).

Isolation and purification of IGF revealed that two forms exist: IGF-1 with a molecular mass of 7.5 kDa and pI of 8.5, and IGF-2 with a molecular mass of 7.4 kDa and pI of 7.0 (Clemmons & Shaw, 1986; Humbel, 1990). Structural analysis at biochemical and genetic levels revealed that IGF-1 and IGF-2 are both single polypeptides with significant sequence similarity with insulin (Daughaday & Rotwein, 1989; Rotwein, 1991; Adamo *et al.*, 1992; Van den Brande, 1992). IGF biosynthesis is regulated by various hormones from the pituitary gland, such as growth

hormone, adrenocorticotrophic hormone, luteinizing hormone, follicle-stimulating hormone and thyroid-stimulating hormone, and growth factors such as PDGF and bFGF. IGF is produced in significant amounts after tissue injury or during malnutrition (Smith *et al.*, 1989a; Isaksson *et al.*, 1991; Nissley & Lopaczynski, 1991). It was originally thought that this growth factor was produced by the liver and acted as a hormone (Hall & Tally, 1989). Now it is generally recognized that it is synthesized by numerous types of cells and tissues, suggesting that it acts via autocrine/paracrine as well as endocrine pathways (Daughaday, 1990; Phillips *et al.*, 1990; Le Roith & Roberts, 1991; Bang & Hall, 1992; Isgaard, 1992; Jennische *et al.*, 1992; Middleton & Tyler, 1992; Lazowski *et al.*, 1994). IGF levels in serum (20–80 nM) are higher than in tissues. However, IGF in serum is largely bound to specific IGF-binding proteins (IGFBPs) which prolong its half life (Adamo *et al.*, 1992). Therefore, the physiologically relevant levels of IGF may be tissue levels rather than serum levels. It has been suggested that IGF-1 is an important postnatal growth factor, affecting development and metabolism of many tissues other than cartilage (Demarquay *et al.*, 1990). IGF-2 plays a regulatory role in glucose metabolism and may be even more important as a fetal growth factor (Bhaumick & Bala, 1991).

IGF receptors

Biological action of IGF-1 and IGF-2 is initiated by binding to their specific cell surface receptors, which differ significantly in structure and peptide binding specificity (Fig. 4) (Czech *et al.*, 1989; Smith *et al.*, 1989b; Yokono *et al.*, 1989). IGF-1 binds preferentially to the type 1 receptor, and IGF-2 to the type 2 receptor, although both receptors can bind the other IGF with lower affinity (Le Roith *et al.*, 1991; Tollefsen *et al.*, 1991). Type 1 receptor is a heterotetrameric glycoprotein consisting of two extracellular α subunits with a molecular mass of 130 kDa, and two transmembrane β subunits with a molecular mass of 95 kDa. The α and β subunits are held together by disulphide bonds, thus forming a tetramer consisting of two $\alpha\beta$ dimers (Ullrich *et al.*, 1986; Le Roith & Raizada, 1989; Neely *et al.*, 1991). Type 1 receptor is formed as a precursor protein that needs a number of post-transcriptional and post-translational modifications including glycosylation, dimerization and proteolytic processing to yield the mature complex consisting of two $\alpha\beta$ subunits (Rechler & Nissley, 1990; Moxham & Jacobs, 1992). The ligand binding site is located in the extracellular α subunit, whereas the intracellular β subunit contains a tyrosine kinase domain responsible for signal transduction (Kjeldsen *et al.*, 1991; Zhang & Roth, 1991). Association of two $\alpha\beta$ dimers is required




Fig. 4. Structure of the type 1 and type 2 IGF receptors. The type 1 receptor is a heterotetrameric complex composed of two extracellular α subunits and two transmembrane β subunits held together by disulphide bonds. The α subunits bind IGF-1 and the β subunits contain tyrosine kinase activity in their cytoplasmic domain. The type 2 receptor has a short cytoplasmic tail and a large extracellular domain, consisting of many cysteine-rich regions.

for receptor signal transduction after ligand binding (Ikari *et al.*, 1988; Tollefsen & Thompson, 1988; Wilden *et al.*, 1989; Soos *et al.*, 1990). IGF-1 binding to the α subunit results in intracellular signal transduction by autophosphorylation of the β subunit and activation of the receptor tyrosine kinase. Tyrosine kinase is responsible for intracellular signal transduction, initiating a cascade of phosphorylation steps including activation of serine and threonine kinases (Corps, 1990). Recently, signal transduction from the IGF-1 type 1 receptor via tyrosine kinase receptor has been linked with the G protein-associated phosphatidylinositol hydrolysis pathway, in a way similar to that described for other growth factor receptors (Cantley *et al.*, 1991; Malcolm & Wright, 1993). After IGF binding and signal transduction, the type 1 receptor is internalized and recycled to the cell surface. The number of type 1 receptors at the cell surface is downregulated by increasing concentrations of IGF-1 (Schalch *et al.*, 1986).

Type 2 receptor is a single chain polypeptide with a molecular mass of 250 kDa consisting of a single transmembrane region and a large extracellular domain (Dahms *et al.*, 1987). This receptor does not seem to have signalling functions similar to the type 1 receptor (Haig & Graham, 1991). Type 2 receptor is identical to the cation-independent mannose-6-phosphate receptor, which binds lysosomal proteins and mediates their transport to the lysosomes (Poiraudau *et al.*, 1994). The mechanism of action of the type 2 receptor was also found to be

mediated via the G protein-associated phosphatidylinositol hydrolysis pathway (Nishimoto *et al.*, 1989).

IGF binding proteins (IGFBPs)

IGF binding proteins (IGFBPs) are proteins with high affinity for insulin-like growth factors. Their interaction plays an essential role in modulating biological functions of these growth factors (Baxter, 1993). Multiple binding proteins have been detected in plasma, amniotic fluid and spinal fluid and in a variety of tissues. Despite a large similarity in structure, the different binding proteins have distinctly different properties that might provide clues to elucidate their exact functions (Baxter, 1991; Lamson *et al.*, 1991). Six classes of these binding proteins, designated IGFBP-1 to 6, have been purified and cloned so far (Drop *et al.*, 1992; Bhaumick, 1993). In serum, they function as a storage depot allowing a ready supply of insulin-like growth factor to tissues. The proteins can bind the growth factor either to sequester it so as to prevent its binding to its receptors, or to present IGF to the receptor so as to improve ligand-receptor interactions, depending upon the structural and biochemical properties of the binding proteins. The ability of IGFBPs to adhere to the cell surface appears to be important in the latter process. IGFBP-1 and IGFBP-2 have RGD sequences (Arg-Gly-Asp) in their terminal region and are therefore structurally capable of binding to integrin receptors at the cell surface (Ruoslahti & Pierschbacher 1987; Brewer *et al.*, 1988). The agonistic or antagonistic nature of IGFBP

modulation of the biological functions of IGF depends on the cell or tissue types involved (Baxter & Martin, 1989; Sara & Hall, 1990).

Chondrocytes produce IGFBPs (Froger-Gaillard *et al.*, 1989; Tesch *et al.*, 1992; Olney *et al.*, 1993; Chevalier & Tyler, 1994). In osteoarthritis, production of IGFBP-3 and 5 by chondrocytes is increased (Dore *et al.*, 1994; Olney *et al.*, 1996; Tardif *et al.*, 1996; Chevalier & Tyler, 1996), which indicates that IGFBPs are involved in pathological cartilage destruction.

Effects of IGF on proteoglycan metabolism in cartilage

IGF-1 is the major growth factor that balances proteoglycan biosynthesis and breakdown in chondrocytes for the maintenance of the cartilage matrix (Schoenle *et al.*, 1982; Hascall *et al.*, 1983; Froesch *et al.*, 1985; McQuillan *et al.*, 1986; Luyten *et al.*, 1988; Schalkwijk *et al.*, 1989a; Curtis *et al.*, 1992; Middleton & Tyler, 1992). IGF-1 can stimulate chondrocyte proteoglycan synthesis in culture in the absence of foetal calf serum (Barone-Varelas *et al.*, 1991; Böhme *et al.*, 1992) and it can directly inhibit both basal and cytokine-induced degradation of proteoglycan in cartilage (Tyler, 1989). IGF-2 is less potent than IGF-1, requiring 50–100 times higher concentrations to achieve similar effects (Luyten *et al.*, 1988). Selective elimination of IGF-1 from serum or synovial fluid with neutralizing antibodies results in a distinct decrease in cartilage proteoglycan synthesis and enhanced proteoglycan breakdown (Schalkwijk *et al.*, 1989a). IGF-1 seems to inhibit production and release of proteases or protease activators, although evidence is not yet conclusive (Ballard *et al.*, 1986; Chandrasekhar *et al.*, 1992; Padayatty *et al.*, 1993).

IGF binding sites on chondrocytes in articular cartilage have been identified as specific IGF receptors (Postel-Vinay *et al.*, 1983; Trippel *et al.*, 1983, 1988; Watanabe *et al.*, 1985; Jansen *et al.*, 1989; Makower *et al.*, 1989; Yasunaga *et al.*, 1995). Trippel *et al.* (1988) calculated that an average of 3×10^4 type 1 receptors are present on the surface of each chondrocyte with a binding affinity to IGF-1 of 4×10^{-8} M.

The effects of IGF-1 and IL-1 are opposite: IGF-1 can speed up recovery from IL-1-induced inhibition of chondrocyte proteoglycan synthesis, while IL-1 promotes cartilage breakdown by inhibiting IGF-1-stimulated synthesis (Taylor *et al.*, 1988; Fosang *et al.*, 1991; Berenbaum *et al.*, 1994; Neidel *et al.*, 1994; Rayan & Hardingham, 1994). IL-1 downregulates IGF mRNA levels in chondrocytes and induces secretion of IGFBP and IGF-1 receptor, indicating that an autocrine loop regulates IGF-1 function (Lazarus *et al.*, 1993; Matsumoto *et al.*, 1994). These

findings suggest that IL-1 and IGF may play significant roles in the pathogenesis of inflammatory joint diseases.

IGF-1 nonresponsiveness of chondrocytes

Despite the importance of IGF in the regulation of proteoglycan metabolism in chondrocytes, only a limited number of studies have been focused on the role of IGF in rheumatoid arthritis. Measurements of IGF levels in serum or synovial fluid of patients with various types of arthritis yielded conflicting results (Coates *et al.*, 1977, 1978; Bennett *et al.*, 1988; Nuvert-Zwart *et al.*, 1988; Aitman *et al.*, 1989; Allen *et al.*, 1991; Keyzer *et al.*, 1995). However, it has now become clear that lower levels of IGF-1 in serum or synovial fluid and/or decreased availability of IGF-1 are not necessarily the main factors that regulate cartilage metabolism.

Recently, it was demonstrated in our laboratory that IGF-1 is essential to maintain chondrocyte proteoglycan synthesis during culture of intact cartilage (Schalkwijk *et al.*, 1989a). However, cultured, experimentally induced, arthritic cartilage obtained from inflamed joints was nonresponsive to IGF-1 (Schalkwijk *et al.*, 1989b). This phenomenon was not caused by total nonresponsiveness of the chondrocytes, since forskolin, which is an upregulator of the adenylate cyclase pathway, could stimulate chondrocyte proteoglycan synthesis. It was suggested that this nonresponsiveness to IGF contributes to cartilage destruction (Schalkwijk *et al.*, 1989b, Van den Berg *et al.*, 1989). Therefore, the exact role of IGF-1, and the mechanism by which IGF-1 regulates chondrocyte proteoglycan metabolism in normal and experimentally induced arthritic cartilage, were studied (Table 1). These studies are reviewed in the following paragraphs.

Regulation of IGF-1 nonresponsiveness by growth factors and glucocorticoids

The *in vitro* effects of IGF-1 and triamcinolone acetonide on the hydrodynamic volume of newly synthesized proteoglycans in normal and arthritic mouse cartilage have been investigated (Van der Kraan *et al.*, 1993; Verschure *et al.*, 1994a). *In vitro* IGF-1 stimulation of chondrocyte proteoglycan synthesis in normal cartilage resulted in the synthesis of hydrodynamically large proteoglycans, which is identical to the *in vivo* situation. In arthritic cartilage, IGF-1 was unable to stimulate chondrocyte proteoglycan synthesis and only small proteoglycan monomers were synthesized, demonstrating that in arthritic cartilage not only are the amounts of proteoglycan that are synthesized decreased, but also that their quality is inferior. In the presence of triamcinolone acetonide, hydrodynamically large proteoglycans were synthesized both in normal and

Table 1. Overview of the role of IGF-1 and the mechanism by which IGF-1 regulates chondrocyte proteoglycan metabolism in mouse normal and experimentally-induced arthritic and human osteoarthritic cartilage to provide insight in IGF-1-related processes underlying cartilage destruction

Normal cartilage

IGF-1 can maintain *in vitro* synthesis of proteoglycan monomers with large hydrodynamic size at similar levels as occur *in vivo*

Glucocorticoids are able to exert similar effects as IGF-1 on the regulation of *in vitro* chondrocyte proteoglycan metabolism

PDGF and bFGF have an additional effect on IGF-1 regulation of chondrocyte proteoglycan synthesis during culture. EGF and TGF β have an inhibitory effect on chondrocyte proteoglycan synthesis

IGF-1 receptors are located at the cell membrane of chondrocytes in the middle and deeper zones of cartilage, whereas intracellular labelling is found in surface zone chondrocytes

Arthritic cartilage

IGF-1 nonresponsiveness during arthritis is highly correlated with the severity of arthritis

The failure of IGF-1 to stimulate *in vitro* chondrocyte proteoglycan synthesis in arthritic cartilage accompanies the synthesis of small proteoglycan monomers, demonstrating that not only the amount but also the quality of synthesized proteoglycan has deteriorated in arthritic cartilage

Chondrocytes from arthritic cartilage do not respond to other growth factors besides IGF-1 during culture

Glucocorticoids are able to bypass disturbed IGF-1 responsiveness and may therefore play an important role in maintaining cartilage integrity in arthritic cartilage

Metalloproteinases and cysteineproteinases, reactive oxygen species, IL-1, and TNF α are not key factors in the maintenance of IGF-1 nonresponsiveness. IL-1 does not induce IGF-1 nonresponsiveness but is indirectly involved in the generation of this phenomenon

The correlation of chondrocyte IGF-1 receptor expression with *in vitro* IGF-1 stimulation and metabolic activity is disturbed in arthritic cartilage

Chondrocytes in arthritic cartilage do not exhibit decreased amounts of IGF-1 receptor on their cell membrane.

Therefore, IGF-1 nonresponsiveness cannot be explained by a decreased receptor expression

Osteoarthritic cartilage

IGF-1 has different effects on chondrocytes in the different cartilage zones in normal cartilage. This metabolic heterogeneity is lost in osteoarthritic cartilage. In such cartilage, chondrocyte proteoglycan synthesis and IGF-1 receptor distribution are dependent on the stage of osteoarthritis

arthritic cartilage. It was concluded that glucocorticoids may play an essential role in maintaining cartilage integrity in both normal and arthritic cartilage.

Furthermore, *in vitro* investigations were carried out to study the effects of other growth factors besides IGF-1, such as bFGF, EGF, PDGF and TGF β on the regulation of chondrocyte proteoglycan metabolism under normal and pathological conditions (Verschure *et al.*, 1994b). It was investigated whether these factors could compensate for IGF-1 nonresponsiveness of the chondrocyte either during repair or during early and/or later phases of experimental arthritis. In normal cartilage, none of the growth factors tested stimulated *in vitro* chondrocyte proteoglycan synthesis as IGF-1 did. EGF and TGF β even inhibited chondrocyte proteoglycan synthesis. Combinations of bFGF or PDGF with IGF-1 exerted significant additional stimulation of chondrocyte proteoglycan synthesis. Normal cartilage that was previously exposed to IL-1 *in vivo* was responsive to IGF-1 and to other growth factors in a similar way as, to normal cartilage. However, experimentally induced arthritic cartilage exhibited a state of nonresponsiveness to all individual

growth factors tested, as well as to combinations of growth factors. This suggests that during the inflammatory process in the joint, chondrocytes display a general deficiency of receptor functions. Apparently, inflammatory processes during experimental arthritis are more complex than processes caused by injecting IL-1 into the knee joints alone.

Regulation of IGF-1 nonresponsiveness by inflammatory mediators

Studies were carried out to see whether the state of nonresponsiveness to IGF-1 during arthritis was caused by either inflammatory mediators such as IL-1, TNF α , or by proteolytic enzymes or reactive oxygen species (Verschure *et al.*, 1995a). Mediators from tissues surrounding patellar cartilage were able to suppress *in vitro* responses of chondrocytes to IGF-1. Priming of normal cartilage with IL-1 did not completely induce IGF-1 nonresponsiveness, but enhanced the ability of suppressive mediators to downregulate the IGF-1 response. Treatment of the cartilage with antibodies against IL-1 α and IL-1 β , or IL-1 receptor antagonist, or antibodies against TNF α , or selective inhibitors of proteolytic enzymes or of reactive oxygen species, did not counteract the

nonresponsiveness to IGF-1 *in vitro*. However, treatment with anti-IL-1 α and anti-IL-1 β antibodies before induction of arthritis abolished nonresponsiveness to IGF-1. It was concluded that neither metalloproteinases and cysteine proteinases nor reactive oxygen species are key factors in IGF-1 nonresponsiveness, and that IL-1 and TNF α alone are not essential for the maintenance of nonresponsiveness. Moreover, IL-1 does not induce IGF-1 nonresponsiveness but is indirectly involved.

Regulation of IGF-1 nonresponsiveness by IGF-1 receptor

The lack of effects of IGF-1 on chondrocytes during arthritis may be due to a defect at the level of IGF-1 receptor binding or postreceptor signalling. Therefore, the amounts and distribution patterns of IGF-1 receptors on the cell membrane of chondrocytes were investigated in normal and arthritic cartilage by immunohistochemical localization and by quantification of IGF-1 receptor (Verschure *et al.*, 1994c). Immunohistochemical labelling of the IGF-1 receptor was performed on cryostat sections of mouse patellas using antibodies against the α subunit of the receptor, and was visualized with confocal laser scanning microscopy (CLSM). Optical sectioning using this technique resulted in improved spatial resolution of the localization of the IGF-1 receptor in chondrocytes, and allowed peripheral labelling to be distinguished quantitatively from intracellular labelling in mouse normal and arthritic cartilage (Verschure *et al.*, 1995b; Verschure *et al.*, 1996a), as well as in human normal and osteoarthritic cartilage (Verschure *et al.*, 1996b) (Figs 5 and 6). Discrimination of specific signals from background labelling, which is difficult when labelling is of low intensity, and background and autofluorescence levels are high, can be accomplished by the application of confocal microscopy (Wilson *et al.*, 1989; Van Oostvelt & Bauwens, 1990; Wells *et al.*, 1989). This is a problem that is inherent to articular cartilage, and therefore CLSM will become increasingly important in cartilage research. Most of the IGF-1 receptor fluorescence was found on chondrocytes of the middle and deeper zones of the cartilage, whereas surface zone chondrocytes exhibited negligible fluorescence. The heterogeneous distribution of IGF-1 receptor in chondrocytes of articular cartilage suggests that effects of IGF-1 on chondrocytes may be distinctly different in the specific cartilage zones.

Confocal laser scanning microscopy was applied to compare labelling of IGF-1 receptor by anti-IGF-1 receptor antibody with labelling of binding sites with biotinylated IGF-1 in the various cartilage zones of normal and experimental arthritic knee joints (Verschure *et al.*, 1996c). In normal cartilage, IGF-1 receptors were found at the cell membrane of chondrocytes of the middle and deep cartilage

zones with anti-IGF-1 receptor antibody, whereas intracellular labelling was highest in chondrocytes of the surface zone. After incubation with biotinylated IGF-1, distinct membrane labelling was not present. In normal cartilage, fluorescence was found to be distributed homogeneously in chondrocytes of the middle and deep zones but not in the surface zone. In cartilage from inflamed knee joints, labelling with anti-IGF-1 receptor antibody was not significantly different from normal cartilage, whereas labelling with biotinylated IGF-1 was increased in chondrocytes of the middle and deeper zones of arthritic cartilage as compared with normal cartilage. The lack of intracellular labelling with biotinylated IGF-1 of surface zone chondrocytes may suggest that intracellular IGF-1 binding sites have low affinity for IGF-1, or that the binding site of the receptor is shielded. Labelling patterns obtained with biotinylated IGF-1 also display IGF-1 binding proteins (IGFBPs). It was suggested that during joint inflammation larger amounts of these binding proteins are present.

Both the distribution patterns of IGF-1 receptor, and *in vitro* IGF-1 stimulation of chondrocyte proteoglycan synthesis were examined in articular cartilage of normal and experimental arthritic mouse knee joints during various phases of arthritis (Verschure *et al.*, 1995b). The distribution patterns of IGF-1 receptor expression in the different cartilage zones of normal cartilage reflected IGF-1 stimulation and proteoglycan synthetic activity of chondrocytes in these layers. Concurrently with the development of arthritis, cartilage lost its capacity to react to IGF-1, but when the inflammatory response was waning, IGF-1 responsiveness returned. Shortly after induction of experimental arthritis, a clear decrease in IGF-1 receptor expression was observed, whereas in later phases of arthritis receptor expression returned to normal levels. It was found that IGF-1 nonresponsiveness was highly correlated with severity of arthritis. The correlation of IGF-1 receptor expression with IGF-1 stimulation and metabolic activity is disturbed in arthritic cartilage, suggesting inadequate or abolished intracellular signal transduction in chondrocytes.

It was also studied whether chondrocyte proteoglycan synthesis correlates with chondrocyte IGF-1 receptor expression in human normal cartilage, as well as in cartilage known to display a shift in proteoglycan synthesis, such as cultured cartilage and osteoarthritic cartilage (Verschure *et al.*, 1996b). Osteoarthritic cartilage shows markedly enhanced proteoglycan synthesis. Cartilage specimens were obtained *post mortem* from human knees within 18 hours after the death of donors with no known clinical history of osteoarthritis. The samples were dissected from macroscopically normal regions as




Fig. 5. (A) Schematic drawing of a confocal laser scanning microscope (CLSM). It collects images that are almost free of out-of-focus signals. This type of analysis results in improved spatial resolution as compared with conventional light microscopy and allows optical sectioning of the specimen (White *et al.*, 1988; Wilson, 1989). (B) The possibility of making optical sections through the entire chondrocyte has been applied to the analysis of IGF-1 receptor labelling in the chondrocyte in the middle optical section.

well as from damaged regions. This procedure yielded a range of grades from mild to moderate and severe osteoarthritis. In normal cartilage, both chondrocyte proteoglycan synthesis and the amount of chondrocyte IGF-1 receptor were low in the surface zones, and high in the middle and deep zone of the cartilage. After culture, an increase in chondrocyte proteoglycan synthesis in the surface zone coincided with upregulation of IGF-1 receptor localisation. In mild osteoarthritis, particularly high levels of chondrocyte synthetic activity were found in the upper cartilage layer, whereas IGF-1 receptor expression was low in this layer, suggesting that factors other than IGF-1 are involved in enhancing chondrocyte proteoglycan synthesis. High proteo-

glycan synthesis activity and chondrocyte IGF-1 receptor labelling were found in moderate osteoarthritis, whereas low levels of proteoglycan synthesis and IGF-1 receptor labelling were observed in severe osteoarthritis.

Final considerations

A critical aspect of the structural integrity of articular cartilage is the ability of chondrocytes to synthesize and degrade proteoglycans. IGF-1 is the most important growth factor that regulates chondrocyte proteoglycan metabolism, by binding to the type 1 receptor. In the present review it is shown that IGF-1 receptor distribution patterns in surface, middle and




Fig. 6. Pseudo 3-D visualization of a CLSM data stack; consecutive optical sections of a chondrocyte located in the middle zone of mouse articular cartilage after being immunohistochemically labelled for the IGF-1 receptor. (A) Chondrocytes located in the middle zone of normal mouse cartilage after labelling with a polyclonal antibody against the α subunit of the IGF-1 receptor. The chondrocyte membrane contains large amounts of IGF-1 receptor fluorescence but intracellular labelling is weak. (B, C) Chondrocytes located in the middle zone of mouse normal (B) or arthritic articular cartilage (C) after being labelled with biotinylated IGF-1. Intracellular labelling is higher when using biotinylated IGF-1 than when using anti-IGF-1 receptor antibody. Chondrocytes in arthritic cartilage display strikingly higher fluorescence intensity than that observed in normal cartilage. High amounts of fluorescence are found directly below the cell membrane of the cells.

deeper zones of mouse articular cartilage appear to reflect a heterogeneity in IGF-1 stimulation and proteoglycan synthesis of chondrocytes in these layers. Intracellular IGF-1 receptors which are mainly found in chondrocytes of the surface zone, represent internalized, recycling or newly synthesized receptors. This intracellular pool of receptors may become important under conditions of stress. The cell surface receptors on chondrocytes of the middle and deeper zones may reflect receptors actively involved in triggering IGF-1 responses. After culture of human cartilage explants in the presence of serum, chondrocytes show markedly enhanced proteoglycan synthesis, mainly in the surface zone. This shift in metabolic activity coincides with an upregulation of IGF-1 receptor expression on the membrane of these cells (Korver *et al.*, 1990; Lafeber *et al.*, 1990, 1992; Maroudas *et al.*, 1990; Verschure *et al.*, 1996b). The quantities of receptors at the cell membrane or intracellularly in various cartilage zones may provide further insight regarding the regulation of chondrocyte metabolic activity. The application of confocal laser scanning microscopy provides a valuable tool to obtain more detailed information on IGF-1 receptor expression in articular cartilage chondrocytes. However, CLSM analysis of cartilage from diseased joints indicates that conclusions have to be drawn with caution from growth factor or

receptor expression patterns without evaluating biological responses triggered by receptor binding of the growth factor involved.

Chronic interference with IGF-1-dependent regulation of chondrocyte proteoglycan synthesis during induced arthritis in the mouse knee joint contributes to severe depletion of articular cartilage. Intra-articular administration of IGF-1 in the treatment of rheumatoid arthritis patients results in pain relief and improvement of joint function (Silveri *et al.*, 1994). However, the present investigation did not provide evidence for the supposed beneficial effects of IGF-1 treatment during induced arthritis in the mouse knee joint with respect to protection against articular cartilage damage. Treatment with glucocorticoids or blocking of IL-1 effects seem to provide a more promising therapeutic approach.

The exact mechanisms underlying IGF-1 nonresponsiveness during experimental arthritis in the mouse knee joint remain unclear. Many possible candidates, such as metalloproteinases, cysteine proteinases, reactive oxygen species, TNF α or IL-1 do not override IGF-1 signalling, although IL-1 seems to play an indirect role in the generation of IGF-1 nonresponsiveness (Joosten *et al.*, 1991; Lin *et al.*, 1992; Blount & Crawford, 1994; Verschure *et al.*, 1995a). The lack of IGF-1 response cannot be explained by receptor downregulation either. This

may indicate that, although IGF-1 receptors are present on the chondrocyte membrane, the membrane lipid bilayer composition may have changed, causing altered configuration of the transmembrane receptor, which may hamper the biological function of the receptor. Furthermore, receptors may be masked or damaged by agents produced by the inflamed synovium (Jansen *et al.*, 1989; Kwok *et al.*, 1989; Bergstedt & Wieloch, 1993; Kato *et al.*, 1993). For instance, IGFBPs may play a role in this process (Tesch *et al.*, 1992; Chevalier & Tyler, 1994). In osteoarthritic human cartilage, elevated IGFBP levels seem to be involved in capturing IGF-1 and therefore in reduced bioavailability of IGF (Dore *et al.*, 1994; Olney *et al.*, 1996; Chevalier & Tyler, 1996; Tardif *et al.*, 1996). Increased labelling in chondrocytes of arthritic cartilage after incubation with biotinylated IGF-1 may indicate increased levels of IGFBPs. However, even at unphysiologically high concentrations of IGF-1 chondrocyte proteoglycan synthesis is not stimulated *in vitro* and chondrocytes maintain their nonresponsiveness to IGF-1. Therefore, if IGFBPs are produced in high concentrations they may be involved in the IGF-1 nonresponsive state during arthritis.

Alternatively, IGF-1 receptors may be defective and unable to exhibit normal biological functions (Soos & Siddle, 1989; Raz *et al.*, 1991; Zhang & Roth, 1991; Attisano *et al.*, 1992; Miki *et al.*, 1992; Werner *et al.*, 1992). Yee *et al.* (1989) have found an alternative IGF-1 receptor transcript with a 3-base deletion in many tissues and cell lines. It would be interesting to investigate whether this alternatively spliced mRNA of IGF-1 receptor is also present in arthritic chondrocytes but not in normal chondrocytes. At present, this hypothesis is investigated by detecting the two different spliced IGF-1 receptor mRNAs with the use of the polymerase chain reaction. In addition, the composition of chondrocyte membranes may be changed and thus also the function of transmembrane proteins, such as the β subunits of the IGF-1 receptor, that are involved in signal transduction pathways. Therefore, possible defects in the phosphorylation pathways leading to receptor signalling will be another interesting item for further research. However, these experiments may introduce artefacts: isolation and subsequent culture of chondrocytes from arthritic knee joints may change the 'arthritic' phenotype of the cells. Moreover, analysis of whole chondrocyte populations leads to loss of information about individual chondrocytes located in the different cartilage layers.

Although arthritis seemed to cause a nonresponsive state to a whole range of potential anabolic growth factors, the glucocorticoid triamcinolone acetonide can maintain *in vitro* chondrocyte proteo-

glycan metabolism. Glucocorticoids may have an indirect effect on chondrocytes in arthritic cartilage; they can modulate the production of inflammatory mediators (Sheppard *et al.*, 1982; Steinberg & Sledge, 1983), and seem to have a priming effect on the biological action of IGF-1 in cartilage metabolism (Itagane *et al.*, 1991; Price *et al.*, 1992). In a previous study, it was shown that a single intra-articular injection in arthritic joints of a depot preparation of the steroid triamcinolone hexacetonide or rimexolone counteracts the severe suppression of chondrocyte proteoglycan synthesis in arthritic cartilage (Joosten *et al.*, 1990). These steroid-treated arthritic joints show normal responsiveness to IGF-1 (Joosten *et al.* unpublished results). In this respect, it will be interesting to analyse whether, at later stages of arthritis or after renewed exacerbation, chondrocytes still display their responsiveness induced by glucocorticoids.

Acknowledgements

The authors wish to thank Rob Kreuger and Rob Lutgerhorst for excellent technical assistance.

References

- ADAMO, M., ROBERTS, C.T. & LE ROITH, D. (1992) How distinct are the insulin and insulin-like growth factor-1 signalling systems? *Bio Factors* **3**, 151–7.
- AHMED, N.A., MARTIN, L.A., WATTS, L.M., PALMER, J., THORNBURG, L., PERIOR, J. & ESSER, R.E. (1992) Peptidyl fluoromethyl ketones as inhibitors of Cathepsin B. Implication for treatment of rheumatoid arthritis. *Biochem. Pharmacol.* **44**, 1201–7.
- AIGNER, T., REICHENBERGER, E., BERTLING, W., KIRSCH, T., STOSS, H. & VON DER MARK, K. (1993) Type X collagen expression in osteoarthritic and rheumatoid articular cartilage. *Virch. Arch. B* **63**, 205–11.
- AIGNER, T., DIETZ, U., STOSS, H. & VON DER MARK, K. (1995) Differential expression of collagen types I, II, III and X in human osteophytes. *Lab. Invest.* **73**, 236–43.
- AITMAN, T.J., PALMER, R.G., LOFTUS, J., ANSELL, B.M., ROYSTON, J.P., TEALE, J.D. & CLAYTON, R.N. (1989) Serum levels and growth failure in juvenile chronic arthritis. *Clin. Exp. Rheumatol.* **557**, 557–61.
- ALLEN, R.C., JIMENEZ, M. & COWELL, T. (1991) Insulin-like growth factor and growth hormone secretion in juvenile chronic arthritis. *Ann. Rheum. Dis.* **50**, 602–6.
- ANASTASSIADES, T. & DZIEWIATKOWSKI, D. (1970) The effect of cortisone on the metabolism of connective tissues in the rat. *J. Lab. Clin. Med.* **75**, 826–39.
- ANNEFELD, M. (1992) Changes in rat epiphyseal cartilage after treatment with dexamethasone and glycosaminoglycan-peptide complex. *Pathol. Res. Pract.* **188**, 649–52.
- AREND, W.P. & DAYER, J.M. (1995) Inhibition of the production of interleukin-1 and tumor necrosis factor

- α in rheumatoid arthritis. *Arthritis Rheum.* **38**, 151–60.
- ATTISANO, L., WRANA, J.L., CHELFETZ, S. & MASSAGUE, J. (1992) Novel activin receptors: Distinct genes and alternative mRNA generate a repertoire of serin/threonin kinase receptors. *Cell* **68**, 97–108.
- AYDELOTTE, M.B. & KEUTTNER, K.E. (1991) Heterogeneity of articular chondrocytes and cartilage matrix. In *Cartilage degradation: basic and clinical aspect* (edited by WOESSNER, J.F. & HOWELL, D.S.) pp. 37–9. New York: Raven Press.
- BAICI, A. & LANG, A. (1990) Effect of interleukin-1 β on the production of cathepsin B by rabbit articular chondrocytes. *FEBS Lett.* **277**, 93–6.
- BALLARD, F.J., READ, L.C., FRANCIS, G.L., BAGLEY, C.J. & WALLACE, J.C. (1986) Binding properties and biological potencies of insulin-like growth factors in L6 myoblasts. *Biochem. J.* **233**, 230–33.
- BANG, P. & HALL, K. (1992) Insulin-like growth factors as endocrine and paracrine hormones. In *The Insulin-like growth factors. Structure and biological functions* (edited by SCHOLFIELD, P.N.) pp. 151–68. Oxford: Oxford University Press.
- BARONE-VARELAS, J., SCHNITZER, T.J., MENG, Q., OTTEN, L. & THONAR, E.J.-M.A. (1991) Age-related differences in the metabolism of proteoglycans in bovine articular cartilage explants maintained in the presence of insulin-like growth factor 1. *Connect. Tissue Res.* **26**, 101–20.
- BARRUECO, J.L., GAZQUEZ, A., REDONO, E. & DURZN, E. (1989) A histopathological and morphometrical study of femur head cartilage in Wistar rats treated with prednisolone. *Histol. Histopathol.* **4**, 317–23.
- BATES, E.J., COLIN, C. & LOWTHER, D.A. (1985) Inhibition of proteoglycan synthesis by hydrogen peroxide in cultured bovine articular cartilage. *Biochim. Biophys. Acta* **838**, 221–8.
- BAXTER, R.C. & MARTIN, J.L. (1989) Binding proteins for the insulin-like growth factors: structure, regulation and function. *Prog. Growth Factor Res.* **1**, 49–68.
- BAXTER, R.C. (1991) Insulin-like growth factor (IGF) binding proteins: the role of serum IGF-BPs in the regulating IGF availability. *Acta Paediatr. Scand.* **372**, 107–14.
- BAXTER, R.C. (1993) Circulating binding proteins for the insulin-like growth factors. *Trends Endocrinol. Metabol.* **4**, 91–6.
- BEHRENS, F., SHEPARD, N. & MITCHELL, N. (1975) Alteration of rabbit articular cartilage by intra-articular injections of glucocorticoids. *J. Bone Joint Surg.* **57A**, 70–6.
- BELLOWS, C.G., HEERSCHE, J.N.M. & AUBIN, J.E. (1989) Effects of dexamethasone on expression and maintenance of cartilage in serum-containing cultures of calvaria cells. *Cell Tissue Res.* **256**, 145–51.
- BENNETT, A.E., SILVERMAN, E.D., MILLER, J.J. & HINTZ, R.L. (1988) Insulin-like growth factors in children with systemic onset of juvenile arthritis. *J. Rheumatol.* **15**, 655–8.
- BENTON, H.P. & TYLER, J.A. (1988) Inhibition of cartilage proteoglycan synthesis by interleukin-1. *Biochem. Biophys. Res. Commun.* **145**, 421–8.
- BENTZEN, K., PETERSEN, J., HALKJAER-KRISTENSEN, J. & INGEMANN-HANSEN, T. (1985) Interleukin-1-like activities in synovial fluids of patients with rheumatoid arthritis and traumatic synovitis. *Rheum. Int.* **5**, 79–82.
- BERENBAUM, F., THOMAS, G., POIRADEAU, S., BEREZIAT, G., CORVOL, M.T. & MASLIAH, J. (1994) Insulin-like growth factors counteract the effect of interleukin-1 β on type II phospholipase A2 expression and arachidonic acid release by rabbit articular chondrocytes. *FEBS Lett.* **340**, 51–5.
- BERGSTEDT, K. & WIELOCH, T. (1993) Changes in insulin-like growth factor 1 receptor density after transient cerebral ischemia in the rat: lack of protection against ischemic brain damage following injection of insulin-like growth factor 1. *J. Cereb. Blood Flow Metab.* **13**, 859–98.
- BHAUMICK, B. (1993) Insulin-like growth factor (IGF-1) binding protein and insulin-like growth factor secretion by cultured chondrocyte cells: identification, characterization and ontogeny during cell differentiation. *Regul. Pept.* **48**, 113–22.
- BHAUMICK, B. & BALA, R.M. (1991) Differential effects of insulin-like growth factor 1 and 2 on growth, differentiation and glucoregulation in differentiating chondrocyte cells in culture. *Acta Endocrinol.* **125**, 201–11.
- BLANCO, F.J., GENG, Y. & LOTZ, M. (1995) Differentiation-dependent effects of IL-1 and TGF β on human articular chondrocyte proliferation are related to inducible nitric oxide synthase. *J. Immunol.* **154**, 4018–26.
- BLONDELON, D., ADOLPHE, M., ZIZINE, L. & LECHAT, P. (1980) Evidence for glucocorticoid receptors in cultured rabbit articular chondrocytes. *FEBS Lett.* **117**, 195–8.
- BLOUNT, S. & CRAWFORD, A. (1994) Regulation of IGF-1 receptors on rabbit articular chondrocytes by inflammatory mediators. *Agents & Actions* **41**, C177–9.
- BÖHME, K., CONSCIENCE-EGLI, M., TSCHAN, T. & WINTERHALTER, K.H. (1992) Induction of proliferation or hypertrophy of chondrocytes in serum-free culture: the role of insulin-like growth factor-I, insulin, or thyroxine. *J. Cell Biol.* **116**, 1035–42.
- BRENNAN, F.M., ZACARIAE, C.O. & CHANTRY, D. (1990) Detection of interleukin-8 (IL-8) in synovial fluids from patients with rheumatoid arthritis and production of IL-8 mRNA by isolated synovial cells. *Eur. J. Immunol.* **20**, 2141–4.
- BREWER, M.T., STETLER, G.L., BUSBY, W.H. & CLEMMONS, D.R. (1988) Cloning, characterization and expression of a human insulin-like growth factor binding protein. *Biochem. Biophys. Res. Commun.* **152**, 1289–97.
- BURTON-WURSTER, N. & LUST, G. (1990) Fibronectin and proteoglycan synthesis in long term cultures of cartilage explants in Ham's F12 supplemented with insulin and calcium: Effects of the addition of TGF β . *Arch. Biochem. Biophys.* **283**, 27–33.
- BUTTLE, D.J. (1994) Lysosomal cysteine endopeptidases in the degradation of cartilage and bone. In *Immunopharmacology of joints and connective tissue* (edited by DAVIES, M.E. & DINGLE, J.T.) pp. 225–43. London: Academic Press.
- CANTLEY, L.C., AUGER, K.R., CARPENTER, C., DUCK-

- WORTH, B., GRARIANI, A., KAPPELLER, R. & SOLTOFF, S. (1991) Oncogenes and signal transduction. *Cell* **64**, 281–302.
- CHANDRASEKHAR, S., HARVEY, A.K. & HUBREY, P.S. (1992) Intra-articular administration of interleukin-1 causes prolonged suppression of cartilage proteoglycan synthesis in rats. *Matrix Biol.* **11**, 1–10.
- CHANDRASEKHAR, S. & HARVEY, A.K. (1992) Differential regulation of metalloproteinase steady-state mRNA levels by IL-1 and FGF in rabbit articular chondrocytes. *FEBS Lett.* **296**, 195–200.
- CHARLES, I.G., PALMER, P.M.J., HICKERY, M.J., BAYLISS, M.T., CHUBB, A.P., HALL, V.S.P., MOSS, D.W. & MONACA, S. (1993) Cloning, characterization and expression of a cDNA encoding an inducible nitric oxide synthase from the human chondrocyte. *Proc. Natl. Acad. Sci. USA* **90**, 11419–23.
- CHEVALIER, X. & TYLER, J.A. (1994) Production of IGF binding proteins by human cartilage explants and role of the IGFBP-3 in the modulation of proteoglycan synthesis caused by IGF-1. *Arthritis Rheum.* **37**, S 293.
- CHEVALIER, X. & TYLER, J.A. (1996) Production of binding proteins and role of the insulin-like growth factor 1 binding protein 3 in human articular cartilage explants. *Br. J. Rheumatol.* **35**, 515–22.
- CHUNEKAMRAI, S., KROOK, L.P., LUST, G. & MAYLIN, G.A. (1989) Changes in articular cartilage after intra-articular injections of methylprednisolone acetate in horses. *Am. J. Vet. Res.* **50**, 1733–41.
- CIRCOLO, A., WELGUS, H.G., PIERCE, G.F., KRAMER, J. & STRUNK, R.C. (1991) Differential regulation of the expression of proteinases/antiproteinases in fibroblasts. *J. Biol. Chem.* **266**, 12283–8.
- CLEMMONS, D.R. & SHAW, D.S. (1986) Purification and biologic properties of fibroblast somatomedin. *J. Biol. Chem.* **261**, 10293–8.
- COATES, C.L., BURWELL, G., BUTTERY, P.J., WALKER, G. & WOODWARD, P.M. (1977) Somatomedin activity in synovial fluid. *Ann. Rheum. Dis.* **36**, 50–5.
- COATES, C.L., BURWELL, R.G., LLOYD-JONES, K., SWANNELL, A.J., WALKER, G. & SELBY, C. (1978) Somatomedin activity in synovial fluid from patients with joint diseases. *Ann. Rheum. Dis.* **37**, 303–14.
- CORPS, A.N. (1990) Signalling at the insulin and insulin-like growth factor receptors: transduction probed by transfection. *J. Endocrinol.* **124**, 349–51.
- CUEVAS, P., BURGOS, J. & BAIRD, A. (1988) Basic fibroblast growth factor (FGF) promotes cartilage repair *in vivo*. *Biochem. Biophys. Res. Commun.* **156**, 611–18.
- CURTIS, A.J., KENG, C., HANDLEY, C.J. & ROBINSON, H.C. (1992) Effect of insulin-like growth factor-1 on the synthesis and distribution of link protein and hyaluronan in explant cultures of articular cartilage. *Biochem. Biophys. Acta* **1135**, 309–17.
- CZECH, M.P., LEWIS, R.E. & CORVERA, S. (1989) Multifunctional glycoprotein receptors for insulin and the insulin-like growth factors. In *Carbohydrate recognition in cellular function. Ciba Foundation Symposium* (edited by CZECH, A.J., KENG, C., HANDLEY, S.J. & ROBINSON, H.C.) pp. 27–44. Chichester: Wiley.
- DAHMS, N.W., LOBEL, P., BREITMEYER, J., CHIRGWIN, J. & KORNFELD, S. (1987) 46 kd 6-phosphate receptor: cloning, expression and homology to the 215 kd mannose 6-phosphate receptor. *Cell* **50**, 181–92.
- DAUGHADAY, W.D. (1990) Editorial: the possible autocrine/paracrine and endocrine roles of insulin-like growth factors of human tumors. *Endocrinology* **127**, 1–4.
- DAUGHADAY, W.H. & ROTWEIN, P. (1989) Insulin-like growth factor 1 and 2 peptide messenger ribonucleic acid and gene structures, serum and tissue concentrations. *Endocrine Rev.* **10**, 68–91.
- DAYER, J.M. & BURGER, D. (1994) Interleukin-1, tumor necrosis factor and their specific inhibitors. *Eur. Cytokine Network* **5**, 563–71.
- DEMARQUAY, D., DUMONTIER, M.F., TSAGRIS, L., BOURGUIGON, J., NATAF, V. & CORVOLL, M.T. (1990) *In vitro* insulin-like growth factor interaction with cartilage cells derived from postnatal animals. *Hormone Res.* **33**, 111–14.
- DI BATTISTA, J.A., MARTEL-PELLETIER, J., CLOUTIER, J.M. & PELLETIER, J.P. (1991) Modulation of glucocorticoid receptor expression in human articular cartilage chondrocytes by cAMP and prostaglandins. *J. Rheumatol.* **18**, 102–5.
- DI GIOVANNE, F.S., NUKI, G. & DUFF, G.W. (1988) Tumor necrosis factor in synovial exudates. *Ann. Rheum. Dis.* **47**, 786–72.
- DORE, S., PELLETIER, J-P, DIBATTISTA, J.A., TARDIF, G., BRAZEAU, P. & MARTEL-PELLETIER, J. (1994) Human osteoarthritic chondrocytes possess an increased number of insulin-like growth factor 1 binding sites but are unresponsive to its stimulation. *Arthritis Rheum.* **37**, 253–63.
- DROP, S.L.S., SCHULLER, A.G.P., LINDENBERGH-KORTLEVE, D.J., GROFFEN, C., BRINKMAN, A. & ZWARTHOFF, E.C. (1992) Structural aspects of the IGFBP family. *Growth Regul.* **2**, 68–79.
- ELLIOT, M.J., MAINI, R.N., FELDMAN, M., LONG-FOX, A., CHARLES, P., KATSIKIS, P., BRENNAN, M., WALKER, J., BIJL, H., GHRAIEB, J. & WOODY, J.N. (1993) Treatment of rheumatoid arthritis with chimeric monoclonal antibodies to tumor necrosis factor α . *Arthritis Rheum.* **12**, 1681–90.
- ELLIS, A.J., CURRY, V.A., POWELL, E.K. & CAWSTON, T.E. (1994) The prevention of collagen breakdown in bovine nasal cartilage by TIMP-1 and TIMP-2 and a low molecular weight synthetic inhibitor. *Biochem. Biophys. Res. Commun.* **201**, 94–100.
- ESSER, R.E.E., MURPHEY, A.M.D., WATTS, L.M., THORNBURG, L.P., PALMER, J.T., TALHOUK, J.W. & SMITH, R.E. (1994) Cysteine proteinase inhibitors decrease articular cartilage and bone destruction in chronic inflammatory arthritis. *Arthritis Rheum.* **2**, 236–47.
- EVANS, C.H. (1991) The role of proteinases in cartilage destruction. *Agents & Actions* **32**, 135–52.
- EVANS, C.H., STEFANOIC-RACIC, M. & LANCASTER, J. (1995) Nitric oxide and its role in orthopaedic disease. *Clin. Orthop.* **312**, 275–94.
- FIRESTEIN, G.S. (1992) Mechanism of tissue destruction and cellular activation in rheumatoid arthritis. *Curr. Opinion Rheumatol.* **4**, 348–54.
- FIRESTEIN, G.S. & ZVAIFLER, N.J. (1990) How important are T cells in chronic rheumatoid synovitis? *Arthritis*

- Rheum.* **33**, 768–73.
- FLANARY, C.R. & SANDY, J.D. (1994) Effects of proteinase inhibitors on aggrecan in chondrocyte cultures. *Ann. NY Acad. Sci.* **732**, 380–3.
- FOSANG, A.J., TYLER, J.A. & HARDINGHAM, T.E. (1991) Effect of interleukin-1 and insulin-like growth factor 1 on the release of proteoglycan components and hyaluronan from pig articular cartilage in explant culture. *Matrix Biol.* **11**, 17–24.
- FOSANG, A.J., LAST, K., NEAME, P.J., MURPHY, G., KNAUPER, V., TSCHESCHE, H., HUGHES, C.E., CATERSON, B. & HARDINGHAM, T.E. (1994) Neutrophil collagenase (MMP-8) cleaves at the aggrecanase site E373-A374 in the interglobular domain of cartilage aggrecan. *Biochem. J.* **304**, 347–51.
- FRANCHIMONT, P. & BASSLEER, C. (1991) Effects of hormones and local growth factors on articular chondrocyte metabolism. *J. Rheumatol.* **18**, 68–70.
- FROESCH, E.R., SCHMID, C., SCHWANDER, J. & ZAPF, J. (1985) Actions of insulin-like growth factor. *Annu. Rev. Physiol.* **47**, 443–67.
- FROGER-GAILLARD, B., HOSSENLOPP, P., ADOLPHE, M. & BINOUX, M. (1989) Production of insulin-like growth factors and their binding proteins by rabbit articular chondrocytes: relationship with cell multiplication. *Endocrinology* **124**, 2365–72.
- FUKUDA, K., KUMANO, F., SAITO, M., OTANI, K. & TANAKA, S. (1995) Insulin-like growth factors maintain steady-state metabolism of proteoglycans in bovine articular cartilage explants. *Arch. Biochem. Biophys.* **267**, 416–25.
- GRAY, R.G. & GOTTLIEB, N.L. (1983) Intra-articular corticosteroid: An updated assessment. *Clin. Orthop. Relat. Res.* **177**, 235–63.
- HAIG, D. & GRAHAM, C. (1991) Genomic imprinting and the strange case of the insulin-like growth factor II receptor. *Cell* **64**, 1045–6.
- HALL, K. & TALLY, M. (1989) The somatomedin-insulin-like growth factors. *J. Int. Med.* **225**, 47–54.
- HANDLEY, C.J., LOWTHER, D.A. & MCQUILLAN, D.J. (1985) The structure and synthesis of proteoglycans of articular cartilage. *Cell Biol. Int. Rep.* **9**, 753–82.
- HANGLOW, A.C., ROWAN, K., LUSCH, L. & COFFEY, J.W. (1995) Degradation of bovine cartilage proteoglycan *in vitro* is enhanced by inhibition of nitric oxide synthase. *Inflamm. Res.* **44** suppl, S151–2.
- HARDINGHAM, T.E. & FOSANG, A.J. (1992) Proteoglycans: many forms and many functions. *FASEB J.* **6**, 861–70.
- HARDINGHAM, T.E. & FOSANG, A.J. (1995) The structure of aggrecan and its turnover in cartilage. *J. Rheumatol.* **43**, 86–90.
- HARDINGHAM, T.E., FOSANG, A.J. & DUDHIA, J. (1992a) Aggrecan, the chondroitin sulfate/keratan sulfate proteoglycan form cartilage. In *Articular cartilage and osteoarthritis* (edited by KUETTNER, K.E., SCHLEYRBACH, R., PEYRON, J.G. & HASCALL, V.C.) pp. 5–20. New York: Raven Press.
- HARDINGHAM, T.E., BAYLISS, M.T., RAYAN, V. & NOBLE, D.P. (1992b) Effects of growth factors and cytokines on proteoglycan turnover in articular cartilage. *Br. J. Rheumatol.* **1**, suppl. 1, 1–6.
- HARRIS, E.D. (1990) Rheumatoid arthritis. Pathophysiology and implications for therapy. *New Eng. J. Med.* **322**, 1277–89.
- HARVEY, A.K., STACK, S.T. & CHANDRASEKHAR, S. (1993) Differential modulation of degradative and repair responses of interleukin-1 treated chondrocytes by platelet-derived growth factor. *Biochem. J.* **292**, 129–36.
- HASCALL, V.C., HANDLEY, C.J., MCQUILLAN, D.J., HASCALL, G.K., ROBINSON, H.C. & LOWTHER, D.A. (1983) The effect of serum on biosynthesis of proteoglycans by bovine articular cartilage in culture. *Arch. Biochem. Biophys.* **224**, 206–23.
- HAUSELMAN, H.J., OPPLIGER, L., MICHEL, B.A., STEFANOVIC-RACIC, M. & EVANS, C.H. (1994) Nitric oxide and proteoglycan biosynthesis by human articular chondrocytes in alginate culture. *FEBS Lett.* **352**, 361–4.
- HENDERSON, B. & BLAKE, S. (1994) Connective tissue destruction in rheumatoid arthritis: therapeutic potential of selective metalloproteinase inhibitors. In *Immunopharmacology of joints and connective tissue* (edited by DAVIES, M.E. & DINGLE, J.T.) pp. 199–223. London: Academic Press.
- HENDERSON, B., PETTIPHER, E.R. & MURPHY, G. (1990) Metalloproteinases and cartilage proteoglycan depletion in chronic arthritis. *Arthritis Rheum.* **33**, 241–6.
- HENDERSON, B., THOMPSON, R.C., HARDINGHAM, T.E. & LETHWAITE, J. (1991) Inhibition of interleukin-1 induced synovitis and articular cartilage proteoglycan loss in rabbit knee by recombinant human interleukin-1 receptor antagonist. *Cytokine* **3**, 246–9.
- HENROTIN, Y., DEBY-DUPONT, G., DEBY, C., FRANCHIMONT, P. & EMERIT, I. (1992) Active oxygen species, articular inflammation and cartilage damage. In *Free radicals and aging* (edited by EMERIT, I. & CHANG, B.) pp. 308–22. Basel: Birkhauser.
- HIRAKI, Y., INOUE, H., ASADA, A. & SUZUKI, F. (1990) Differential modulation of growth and phenotypic expression of chondrocytes in sparse and confluent cultures by growth factors in cartilage. *J. Bone Mineral Res.* **10**, 1077–85.
- HOLDEN, G. & HUME, K.P. (1962) The newer corticosteroids for local injection. *Ann. Phys. Med.* **6**, 178–82.
- HOLLANDER, J.L., BROWN JR, E.M., JESSAR, R.A. & BROWN, C.Y. (1951) Hydrocortisone and cortisone injected into arthritic joints: comparative effects of and use of hydrocortisone as a local antiarthritic agent. *J. Am. Med. Assn.* **147**, 1622–35.
- HOLT, I., COOPER, R.G. & DENTON, J. (1992) Cytokine inter-relationships and their association with disease activity in arthritis. *Br. J. Rheumatol.* **31**, 725–33.
- HOPKINS, S.J., HUMPHREYS, M. & JAYSON, M.I.V. (1988) Cytokines in synovial fluid. The presence of biologically active and immunoreactive IL-1. *Clin. Exp. Immunol.* **72**, 422–7.
- HOUSSIAU, F.A., DEVOGLAER, J.P., VAN DAMME, J. & DEDEUXCHAISNES, C.N. (1988) Interleukin-6 in synovial fluid: detection by RIA, bioassay and the presence of bioassay-inhibiting factors. *Rheumatol. Int.* **31**, 784–8.
- HOWARD, M., O'GARRA, A., ISHIDA, H., DE WAAL-MALEFIJ, R. & DE VRIES, J. (1992) Biological properties of Interleukin 10. *J. Clin. Immunol.* **12**, 239–47.
- HUGHES, C.E., CATERSON, B., FOSANG, A.J., ROUGHLEY,

- P.J. & MORT, J.S. (1995) Monoclonal antibodies that specifically recognize neopeptide sequences generated by 'aggrecanase' and matrix metalloproteinase cleavage of aggrecan: application to catabolism in situ and in vitro. *Biochem. J.* **305**, 799–804.
- HUMBEL, R. (1990) Insulin-like growth factors I and II. *Eur. J. Biochem.* **190**, 445–62.
- IKARI, N., YOSHINO, H., MOSES, A.C. & FLIER, J.S. (1988) Evidence that receptor aggregation may play a role in transmembrane signaling through the insulin-like growth factor-1 receptor. *Mol. Endocrinol.* **2**, 831–7.
- INOUE, H., KATO, Y., IWAMOTO, M., HIRAKI, Y., SAKUDA, M. & SUZUKI, F. (1989) Stimulation of cartilage-matrix proteoglycan synthesis by morphologically transformed chondrocytes grown in the presence of fibroblast growth factor and transforming growth factor-beta. *J. Cell. Physiol.* **138**, 329–37.
- ISAKSSON, O.G., OHLSSON, C., NILSSON, A., ISGAARD, J. & LINDAHL, A. (1991) Regulation of cartilage growth by growth hormone and insulin-like growth factor 1. *Pathology* **5**, 451–3.
- ISGAARD, J. (1992) Expression and regulation of IGF-1 in cartilage and skeletal muscle. *Growth Regul.* **2**, 16–22.
- ITAGANE, Y., INADA, H., FUJITA, K. & ISSHIKI, G. (1991) Interactions between steroid hormones and insulin-like growth factor-1 in rabbit chondrocytes. *Endocrinology* **128**, 1419–24.
- IWAMOTO, M., SHIMAZU, A., NAKASHIMA, K., SUZUKI, F. & KATO, Y. (1991) Reduction in basic fibroblast growth factor receptor is coupled with terminal differentiation of chondrocytes. *J. Biol. Chem.* **266**, 461–7.
- JANSEN, J., VAN BUUL-OFFERS, S.C., HOOGERBRUGGE, C.M., POORTER, T.L., CORVOL, M.T. & VAN DEN BRANDE, J.L. (1989) Characterization of specific insulin-like growth factor (IGF)-1 and IGF-11 receptors on cultured rabbit articular chondrocyte membranes. *Endocrinology* **129**, 245–9.
- JARVINEN, T.A.H., MOILANEN, T., JARVINEN, T.L.N. & MOILANEN, E. (1995) Nitric oxide mediates interleukin-1 induced inhibition of glycosaminoglycan synthesis in rat articular cartilage. *Mediators of Inflammation* **4**, 107–11.
- JENNISCHE, E., ISGAARD, J. & ISAKSSON, O.G.P. (1992) Local expression of insulin-like growth factors during tissue growth and regeneration. In *The Insulin-like growth factors. Structure and biological functions* (edited by SCHOLFIELD, P.N.) pp. 221–33. Oxford: Oxford University Press.
- JOOSTEN, L.A.B., HELSEN, M.A. & VAN DEN BERG, W.B. (1990) Protective effect of Rimexolone on cartilage damage in arthritic mice: a comparative study with Triamcinolone hexacetonide. *Agents & Actions* **31**, 135–42.
- JOOSTEN, L.A.B., HELSEN, H.A. & VAN DEN BERG, W.B. (1991) Transient chondrocyte non-responsiveness to insulin-like growth factor-1 upon H₂O₂ exposure is not related to IGF-1 receptor damage. *J. Rheumatol.* **18**, 585–90.
- JOOSTEN, L.A.B., HELSEN, M.M.A. & VAN DEN BERG, W.B. (1995) Dominant role of IL-1 in cartilage destruction in TNF driven joint inflammation. *Transactions of the 41st Annual Meeting of the Orthop. Res. Soc.* **20**, 100.
- JOOSTEN, L.A.B., HELSEN, H.A., VAN DE LOO, F.A.J. & VAN DEN BERG, W.B. (1996) Anti-cytokine treatment of established type collagen-induced arthritis in DBA/1 mice. *Arthritis Rheum.* **39**, 797–809.
- KATO, H., FARIA, T.N., STANNARD, B., LEVY-TOLEDANO, R., TAYLOR, S.I., ROBERTS JR, C.T. & LE ROITH, D. (1993) Paradoxical biological effects of overexpressed insulin-like growth factor-1 receptors in Chinese hamster ovary cells. *J. Cell Physiol.* **156**, 145–52.
- KATO, Y. & GOSPODAROWICZ, D. (1985a) Stimulation by glucocorticoid of the synthesis of cartilage-matrix proteoglycan synthesis in rabbit craniofacial chondrocytes in vitro. *Arch. Biol.* **33**, 893–9.
- KATO, Y. & GOSPODAROWICZ, D. (1985b) Sulfated proteoglycan synthesis by confluent cultures of rabbit costal chondrocytes. *J. Cell Biol.* **100**, 477–85.
- KATO, Y. & IWAMOTO, M. (1990) Fibroblast growth factor is an inhibitor of chondrocyte terminal differentiation. *J. Biol. Chem.* **265**, 5903–9.
- KATO, Y., NOMURA, Y., TSUJI, M., OHMAE, H., NAKAZAWA, T. & SUZUKI, F. (1981) Multiplication-stimulating activity (MSA) and cartilage-derived factor (CDF): Biological actions in cultured chondrocytes. *J. Biochem.* **90**, 1377–86.
- KATO, Y., HIRAKI, Y., INOUE, H., KINOSHITA, M., YUTANI, Y. & SUZUKI, F. (1983) Differential and synergistic actions of somatomedin-like growth factors, fibroblast growth factor and epidermal growth factor in rabbit costal chondrocytes. *Eur. J. Biochem.* **129**, 685–90.
- KELLY, W.N., HARRIS, E.D., RUDDY, S. & SLEDGE, C.B. (1989) *Textbook of rheumatology*. Philadelphia: W.B. Saunders.
- KEYZER, G.M., HEER, A.H., KRIEGSMAN, J., GEILER, T., KEYZER, C., GAY, R.E. & GAY, S. (1995) Detection of insulin-like growth factor 1 and 2 in synovial tissue specimens of patients with rheumatoid arthritis and osteoarthritis by in situ hybridization. *J. Rheumatol.* **22**, 275–81.
- KJELDSEN, T., ANDERSEN, A.S., WIBERG, F.C., RASMUSSEN, J.S., SCHAFFER, L., BALSCHMIDT, P., MOLER, K.B. & MOLLER, N.P.H. (1991) The ligand specificities of the insulin receptor and the insulin-like growth factor receptor reside in the different regions of common binding site. *Proc. Natl. Acad. Sci. USA* **88**, 4404–8.
- KORVER, G.H.V., VAN DE STADT, R.J., VAN KAMPEN, G.P., KILJAN, E. & VAN DER KORST, J.K. (1990) Composition of proteoglycans synthesized in different layers of cultured anatomically intact articular cartilage. *Matrix Biol.* **10**, 394–401.
- KWOK, C.F., GOLDSTEIN, B.J., MULLER-WEILAND, D.M., LEE, T-S, KAHN, R. & KING, G.L. (1989) Identification of persistent defects in insulin receptor structure and function in capillary endothelial cells from diabetic rats. *J. Clin. Invest.* **83**, 127–36.
- LAFABER, F.P.J.G., VAN DER KRAAN, P.M., ROY, J.L.A.M., HUBER-BRUNING, O. & BIJLSMA, J.W.J. (1990) Articular cartilage explant culture; an appropriate in vitro system to compare osteoarthritic and normal human cartilage. *Connect. Tiss. Res.* **29**, 287–99.
- LAFABER, F.P.J.G., VAN DER KRAAN, P.M., VAN ROY, H.L.A.M., VITTERS, E.L., HUBER-BRUNING, O., VAN

- DEN BERG, W.B. & BIJLSMA, J.W.J. (1992) Local changes in proteoglycan synthesis during culture are different for normal and osteoarthritic cartilage. *Am. J. Pathol.* **140**, 1421–9.
- LAMSON, G., GIUDICE, L.C. & ROSENFELD, R.G. (1991) Insulin-like growth factor binding proteins: Structural and molecular relationships. *Growth Factors* **5**, 19–28.
- LARK, M.W., GORDY, J.T., WEIDER, J.R., AYALA, J., KIMURA, J.H., WILLIAMS, H.R., MUMFORD, R.A., FLANNERY, C.R., CARLSON, S.S. & IWATA, M. (1995) Cell-mediated catabolism of aggrecan. Evidence that cleavage at the "aggrecanase" site (Glu 373-Ala374) is a primary event in proteolysis of the interglobular domain. *J. Biol. Chem.* **270**, 2550–6.
- LARRICK, J.W. & KUNKEL, S.L. (1988) The role of tumor necrosis factor and interleukin 1 in the immunoinflammatory response. *Pharm. Res.* **5**, 129–39.
- LAZARUS, D.D., LYLE, L., MOLDAWER & LOWRY, S.F. (1993) Insulin-like growth factor-1 activity is inhibited by interleukin-1 α , tumor necrosis factor α and interleukin-6. *Lymphokine Cytokine Res.* **12**, 219–23.
- LAZOWSKI, D.A., FRAHER, L.J., HODSMAN, A., STEER, B., MODROWSKI, D. & HAN, V.K.M. (1994) Regional variation of insulin-like growth factor-1 gene expression in mature rat bone and cartilage. *Bone* **15**, 563–76.
- LEBOVITZ, H.E. & EISENBARTH, G.S. (1975) Hormonal regulation of cartilage growth and metabolism. *Vitamins Hormones* **33**, 575–602.
- LE ROITH, D. & RAIZADA, M.K. (1989) *Molecular and cellular biology of insulin-like growth factors and their receptors*. New York: Plenum Press.
- LE ROITH, D. & ROBERTS, C.T. (1991) At the cutting edge: insulin-like growth factor-1 (IGF-1): a molecular basis for endocrine versus local action? *Mol. Cell. Endocrinol.* **77**, C57–61.
- LE ROITH, D., ADAMO, M., WERNER, H. & ROBERTS, C.T. (1991) Insulin-like growth factors and their receptors as growth regulators in normal physiology and pathologic states. *Trends Endocrinol. Metab.* **2**, 134–9.
- LETHWAITE, J.C., BLAKE, S.M., HARDINGHAM, T.E., WARDEN, P.J. & HENDERSON, B. (1994) The effect of recombinant human interleukin-1 receptor antagonist on the induction phase of antigen-induced arthritis in the rabbit. *Br. J. Rheumatol.* **21**, 467–72.
- LIN, T., WANG, D., NAGPAL, M.L., CHANG, W. & CALKINS, J.H. (1992) Down-regulation of Leydig cell insulin-like growth factor-1 gene expression by interleukin-1. *Endocrinology* **130**, 1217–24.
- LO, Y.Y. & CRUZ, T.F. (1995) Involvement of reactive oxygen species in cytokine and growth factor induction of c-fos expression in chondrocytes. *J. Biol. Chem.* **270**, 11727–30.
- LUYTEN, F.P., HACALL, V.C., NISSLEY, S.P., MORALES, T.I. & REDDI, A.H. (1988) Insulin-like growth factors maintain steady-state metabolism of proteoglycans in bovine articular cartilage explants. *Arch. Biochem. Biophys.* **267**, 416–25.
- MACIEWICZ, R.A. & WOTTON, S.F. (1991) Degradation of cartilage matrix components by the cysteine proteinases Cathepsins B and L. *Biomed. Biochem. Acta* **50**, 561–4.
- MACNAUL, K.L., CHARTRAIN, N., LARK, M., TOCCI, M.J. & HUTCHINSON, N.I. (1990) Discoordinate expression of stromelysin, collagenase and tissue inhibitor of metalloproteinases in rheumatoid human synovial fibroblasts. *J. Biol. Chem.* **265**, 17238–45.
- MAKOWER, A.-M., SKOTTNER, A. & WROBLEWSKI, J. (1989) Binding of insulin-like growth factor-1 (IGF-1) to primary cultures of chondrocytes from rat rib growth cartilage. *Cell Biol. Int. Rep.* **13**, 655–65.
- MALCOLM, R.A. & WRIGHT, N.A. (1993) Growth factors and growth factor receptors. *Br. J. Hosp. Med.* **49**, 774–86.
- MANKIN, H.J. & CONGER, K.A. (1966) The acute effects of intra-articular hydrocortisone on articular cartilage in rabbits. *J. Bone Surg.* **48A**, 1383–8.
- MARLES, P.J., HOYLAND, J.A., PARKINSON, R. & FREEMONT, A.J. (1991) Demonstration of variation in chondrocyte activity in different zones of articular cartilage: an assessment of the value of in-situ hybridisation. *Int. J. Exp. Pathol.* **72**, 171–82.
- MAROUDAS, A., SCHNEIDERMAN, R. & WEINBERG, C. (1990) Comparison between effects of serum and insulin on GAG synthesis in different zones of cultured human articular cartilage. *Trans. Orthop. Res. Soc.* **15**, 315.
- MARTEL-PELLETIER, J., MCCOLLUM, R., FUJIMOTO, N., OBATA, K., CLOUTIER, J.M. & PELLETIER, J.P. (1994) Excess of metalloproteinases over tissue inhibitor of metalloproteinase may contribute to cartilage degradation in osteoarthritis and rheumatoid arthritis. *Lab. Invest.* **70**, 807–15.
- MATSUMOTO, T., TSUKAZAKI, T., ENORMOTO, H., IWASAKI, K. & YAMASHITA, S. (1994) Effects of interleukin-1 β on insulin-like growth factor-1 autocrine/paracrine axis in cultured rat articular cartilage. *Ann. Rheum. Dis.* **53**, 128–33.
- MCQUILLAN, D.J., HANDLEY, C.J., CAMPBELL, M.A., BOLIS, S., MILWAY, V.E. & HERINGTON, A.C. (1986) Stimulation of proteoglycan synthesis by serum and insulin-like growth factor-1 in cultured bovine articular cartilage. *Biochem. J.* **240**, 424–30.
- MIDDLETON, J.F.S. & TYLER, J.A. (1992) Upregulation of insulin-like growth factor 1 gene expression in the lesions of osteoarthritic human articular cartilage. *Ann. Rheum. Dis.* **51**, 440–7.
- MIKI, T., BOTTARO, D.P., FLEMING, T.P., SMITH, C.L., BURGESS, W.H., CHAN, A.M.P. & AARONSON, S.A. (1992) Determination of ligand-binding specificity by alternative splicing: Two distinct growth factor receptors encoded by a single gene. *Proc. Natl. Acad. Sci. USA* **89**, 246–50.
- MORALES, T.I. (1994) Transforming growth factor beta and insulin-like growth factor-1 restore proteoglycan metabolism of bovine articular cartilage after depletion by retinoic acid. *Arch. Biochem. Biophys.* **315**, 190–8.
- MORT, J.S., DODGE, G.R., RHOUGHLEY, P.J., LIU, J., FINCH, G., DIPASQUALE, G. & POOLE, A.R. (1993) Direct evidence for active metalloproteinases mediating matrix degradation in interleukin-1 stimulated human articular cartilage. *Matrix Biol.* **102**, 95–102.
- MOXHAM, C. & JACOBS, S. (1992) Insulin-like growth

- factor receptors. In *Insulin-like growth factors. Structure and biological functions* (edited by SCHOLFIELD, P.N.) pp. 80–101. Oxford: Oxford University Press.
- MURELL, G.A., JANG, D. & WILLIAMS, R.J. (1995) Nitric oxide activates metalloproteinase enzymes in articular cartilage. *Biochem. Biophys. Res. Commun.* **206**, 15–21.
- NATAF, V., TSAGRIS, L., DUMOTIER, M.F., BONAVENTURE, J. & CORVOL, M. (1990) Modulation of sulfated proteoglycan synthesis and collagen gene expression by chondrocytes grown in the presence of bFGF alone or combined with IGF-1. *Reprod. Nutr. Develop.* **30**, 331–42.
- NEELY, E.K., BEUKERS, M.W., OH, Y., COHEN, P. & ROSENFELD, R.G. (1991) Insulin-like growth factor receptors. *Acta Paediatr. Scand.* **372**, 116–23.
- NEIDEL, J., SCHULZE, M. & SOVA, L. (1994) Insulin-like growth factor 1 accelerates recovery of articular cartilage proteoglycan synthesis in culture after inhibition by interleukin-1. *Arch. Orthop. Trauma Res.* **114**, 43–8.
- NERLICH, A.G., KIRSCH, T., WIEST, I., BETZ, P. & VON DER MARK, K. (1992) Localization of collagen type X in human fetal and juvenile articular cartilage and bone. *Histochemistry* **98**, 275–81.
- NGUYEN, G., MURPHY, G., ROUGHLY, P.J. & MORT, J.S. (1989) Degradation of proteoglycan aggregate by cartilage metalloproteinase enzymes. *Biochem. J.* **259**, 61–7.
- NISHIMOTO, I., MURAYAMA, Y., KATADA, T., UI, M. & OGATA, E. (1989) *J. Biol. Chem.* **264**, 14029–36.
- NISSLEY, P. & LOPACZYNSKI, W. (1991) Insulin-like growth factor receptors. *Growth Factors* **5**, 29–43.
- NOURI, A.M.E., PANAJI, G.S. & GOODMAN, S.M. (1984) Cytokines and the chronic inflammation of rheumatic disease. The presence of interleukin-1 in synovial fluids. *Clin. Exp. Immunol.* **55**, 295–302.
- NUVERT-ZWART, I., SCHALKWIJK, J., JOOSTEN, L.A.B., VAN DEN BERG, W.B. & VAN DE PUTTE, L.B.A. (1988) Effects of synovial fluid and synovial fluid cells on chondrocyte metabolism in short term tissue culture. *J. Rheumatol.* **15**, 210–6.
- OGAWA, Y., KSANDER, G.A., PRATT, B.M., SAWAMURA, S.J., ZIMAN, J.M., GERHARDT, C.O., AVIS, P.D., MURRAY, M.J. & MCPHERSON, J.M. (1991) Differences in the biological properties of transforming growth factor- β and platelet-derived growth factor *in vivo*. *Growth Factors* **5**, 57–68.
- O'KEEFE, J.O., CRABB, I.D., PUZAS, J.E. & ROSIER, R.N. (1994) Effects of transforming growth factor β 1 and fibroblast growth factor on DNA synthesis in growth plate chondrocytes are enhanced by insulin-like growth factor-1. *J. Orthop. Res.* **12**, 299–310.
- OLNEY, R.C., SMITH, R.L., KEE, Y. & WILSON, D.M. (1993) Production and hormonal regulation of insulin-like growth factor binding proteins in bovine chondrocytes. *Endocrinology* **133**, 563–70.
- OLNEY, R.C., TSUCHIYA, K., WILSON, D.M., MOHTAI, M., MALONEY, W.J., SCHURMAN, D.J. & SMITH, R.L. (1996) Chondrocytes from osteoarthritic cartilage have increased expression of insulin-like growth factor 1 (IGF-1) and IGF-binding protein-3 (IGFBP-3) and -5, but not IGF-2 or IGFBP-4. *J. Clin. Endocrinol. Metab.* **81**, 1096–103.
- PADAYATTY, S.J., ORME, S., ZENOBI, P.D., STICKLAND, M.H., BELCHTZ, P.E. & GRANT, P.J. (1993) The effects of insulin-like growth factor-1 on plasminogen activator inhibitor-1 synthesis and secretion: Results from *in vitro* and *in vivo* studies. *Thromb. Haemost.* **70**, 1009–13.
- PALMER, R.M.J., HICKERY, M.S., CHARLES, I.G., MANCADA, S. & BAYLISS, M.T. (1993) Induction of nitric oxide in human chondrocytes. *Biochem. Biophys. Res. Commun.* **193**, 398–405.
- PHILLIPS, L.S., HARP, J.B., GOLDSTEIN, S., KLEIN, J. & PAO, C-I. (1990) Regulation and action of insulin-like growth factors at the cellular level. *Proc. Nutr. Soc.* **49**, 451–8.
- POIRAUDEAU, S., KERGOSIEN, N., LIEBERHERR, M., AMOR, B. & CORVOL, M. (1994) Evidence that IGF2 but not IGF1 decreases chondrocyte lysosomal enzyme activities and increases calcium intracellular level through the type II IGF receptor (IGF2-M6P R). *Arthritis Rheum.* **37**, S294.
- POSTEL-VINAY, M.C., CORVOL, M.T., LANG, T., FRAUD, F., GUYDA, H. & POSNER, B. (1983) Receptors for insulin-like growth factors in rabbit articular and growth plate chondrocytes in culture. *Exp. Cell Res.* **148**, 105–16.
- PRICE, W.A., STILES, A. & MOATS-STAATS, B.M. (1992) Gene expression of insulin-like growth factors (IGF-1's), the type 1 IGF-1 receptor, and IGF-1-binding proteins in dexamethasone-induced fetal growth retardation. *Endocrinology* **130**, 1424–32.
- RAYAN, V. & HARDINGHAM, T.E. (1994) The recovery of articular cartilage in explant culture from interleukin-1 α : effects on proteoglycan synthesis and degradation. *Matrix Biol.* **14**, 263–71.
- RAZ, V., KELMAN, Z., ZVIVI, A., NEUFELD, G., GICOL, D. & YARDEN, Y. (1991) PCR based identification of new receptors: molecular cloning of a receptor for fibroblast growth factors. *Oncogene* **6**, 153–60.
- RECHLER, H.M. & NISSLEY, S.P. (1990) Insulin-like growth factors. In *Peptide growth factors and their receptors* (edited by SPORN, M.B. & ROBERS, A.B.) pp. 263–76. New York: Springer-Verlag.
- REICHENBERGER, E., AIGNER, T., VON DER MARK, K., STOSS, H., BERTLING, W. (1991) In situ hybridization studies on the expression of type X collagen in fetal human cartilage. *Dev. Biol.* **148**, 562–72.
- ROBERTS, C.R., ROUGHLY, P.J. & MORT, J.S. (1989) Degradation of human proteoglycan aggregate induced hydrogen peroxide. Protein fragmentation amino acid modification and hyaluronic acid cleavage. *Biochem. J.* **259**, 805–11.
- ROOSE, J.P. & VAN NOORDEN, C.J.F. (1995) Synthetic protease inhibitors: Promising compounds to arrest pathobiologic processes. *J. Lab. Clin. Med.* **125**, 433–41.
- ROSIER, R.N., O'KEEFE, R.J., CRABB, I.D., PUZAS, J.E. & BRAND, J.S. (1988) Articular and growth plate chondrocytes exhibit markedly different metabolic responses to growth plate factors. *J. Bone Mineral Res.* **S185**, 468.
- ROTWEIN, P. (1991) Structure, evolution, expression and regulation of insulin-like growth factors I and II.

Growth Factors 5, 3–18.

- RUOSLAHTI, E. & PIERSCHBACHER, M.D. (1987) New perspectives in cell adhesion: RGD and integrins. *Science* 238, 491–7.
- SAH, R.L., CHEN, A.C., GRODZINSKY, A.J. & TRIPPEL, S.B. (1994) Differential effects of bFGF and IGF-1 on matrix metabolism in calf and adult bovine cartilage explants. *Arch. Biochem. Biophys.* 308, 137–47.
- SAKLATVALA, J. (1986) Tumor necrosis factor α stimulates resorption and inhibits synthesis of proteoglycan in cartilage. *Nature* 322, 547–9.
- SAKURAI, H., KOHSAKA, H., LIU, M.F., HIGASHIYAMA, H., KANNO, K., SAITO, I. & MIYASAKA, N. (1995) Nitric oxide production and inducible nitric oxide synthase expression in inflammatory arthritides. *J. Clin. Invest.* 96, 2357–63.
- SALMON, W.D. & DAUGHADAY, W.H. (1957) A hormonally controlled serum factor which stimulates sulfate incorporation by cartilage *in vitro*. *J. Lab. Clin. Med.* 49, 825–36.
- SAMPSON, H.W. & CANNON, M.S. (1986) Zonal analysis of metabolic profiles of articular-epiphyseal cartilage chondrocytes. *Histochem. J.* 18, 265–70.
- SARA, V.R. & HALL, K. (1990) Insulin-like growth factors and their binding proteins. *Physiol. Rev.* 70, 591–614.
- SAXNE, T., PALLADINO, M.A., HEINEGARD, D., TALAL, N. & WOLLHEIM, F.A. (1988) Detection of tumor necrosis factor α in rheumatoid arthritis synovial fluid and serum. *Arthritis Rheum.* 31, 1041–5.
- SCHALCH, D.S., SESSIONS, C.M., FARLEY, A.C., MASAKAWA, A., EMLER, C.A. & DILLS, D.G. (1986) Interaction of insulin-like growth factor 1/Somatomedin with cultured rat chondrocytes: Receptor binding and internalization. *Endocrinology* 118, 1590–7.
- SCHALKWIJK, J., VAN DEN BERG, W.B., VAN DE PUTTE, L.B.A. & JOOSTEN, L.A.B. (1985a) Cationization of catalase, peroxidase and superoxide dismutase. Effect of improved intraarticular retention on experimental arthritis in mice. *J. Clin. Invest.* 76, 198–205.
- SCHALKWIJK, J., VAN DEN BERG, W.B., VAN DE PUTTE, L.B.A. & JOOSTEN, L.A.B. (1985b) Hydrogen peroxide suppresses the proteoglycan synthesis of intact articular cartilage. *J. Rheumatol.* 12, 205–10.
- SCHALKWIJK, J., JOOSTEN, L.A.B., VAN DEN BERG, W.B., WIJK, J.J. & VAN DE PUTTE, L.B.A. (1989a) Insulin-like growth factor stimulation of chondrocyte proteoglycan synthesis by human synovial fluid. *Arthritis Rheum.* 32, 66–71.
- SCHALKWIJK, J., JOOSTEN, L.A.B., VAN DEN BERG, W.B. & VAN DE PUTTE, L.B.A. (1989b) Chondrocyte non-responsiveness to insulin-like growth factor 1 in experimental arthritis. *Arthritis Rheum.* 32, 894–900.
- SCHLAGHECKE, R., KORNELY, E., WOLLENHAUPT, J. & SPECKER, C. (1992) Glucocorticoid receptors in rheumatoid arthritis. *Arthritis Rheum.* 35, 740–4.
- SCHOENLE, E., ZAPF, J., HUMBEL, R.E. & FROESCH, E.R. (1982) Insulin-like growth factor stimulates growth in hypophysectomized rats. *Nature* 296, 252–3.
- SHAPIRO, S.D., CAMPBELL, E.J., KOBAYASHI, D.K. & WELGUS, H.G. (1990) Immune modulation of metalloproteinase production in human macrophages. *J. Clin. Invest.* 86, 1210–40.
- SHEPPARD, L., PILSWORTH, L.C.M., HAZLEMEN, B. & DINGLE, J.T. (1982) Effects of antirheumatoid drugs on the production and action of porcine catabolin. *Ann. Rheum. Dis.* 41, 463–8.
- SHINMEI, M., KIKUCHI, T., MASUDA, K. & SHIMURA, Y. (1988) Effects of interleukin-1 and anti-inflammatory drugs on the degradation of human articular cartilage. *Drugs* 35 suppl. 1, 33–41.
- SICZKOWSKI, M. & WATT, F.M. (1990) Subpopulations of chondrocytes from different zones of pig articular cartilage: Isolation, growth and proteoglycan synthesis in culture. *J. Cell Sci.* 97, 349–60.
- SILBERMANN, M. & MAOR, G. (1985) Receptor-mediated glucocorticoid inhibition of cell proliferation in mouse growth cartilage *in vitro*. *Acta Endocrinol.* 108, 343–50.
- SILVERI, F., MOROSINI, P., BRECCIAROLI, D. & CERVINI, C. (1994) Intra-articular injection of somatostatin in knee osteoarthritis clinical results and IGF-1 serum levels. *Int. J. Clin. Pharm. Res.* 14, 79–85.
- SMITH, R.L., PALATHUMPAT, M.V., KU, C.W. & HINTZ, R.L. (1989a) Growth hormone stimulates insulin-like growth factor 1 actions on adult articular cartilage. *J. Orthop. Res.* 7, 198–207.
- SMITH, U., ERIKSSON, J. & LONNROTH, P. (1989b) Receptors for insulin-like growth factors. *J. Internal Med.* 225, 43–5.
- SOOS, M.A. & SIDDLE, K. (1989) Immunological relationships between receptors for insulin and insulin-like growth factor receptor 1: Evidence for structural heterogeneity of insulin-like growth factor 1 receptors involving hybrids with insulin receptors. *Biochem. J.* 263, 553–63.
- SOOS, M.A., WHITTAKER, J., LAMMERS, R., ULLRICH, A. & SIDDLE, K. (1990) Receptors for insulin and insulin-like growth factor-1 can form hybrid dimers. *Biochem. J.* 270, 383–90.
- STADLER, J., STEFANOFIC-RACIC, M., BILLAR, T.R., CURRAN, R.D., MCINTYRE, L.A., GEORGESCU, H.I., SIMMONS, R.L. & EVANS, C.H. (1991) Articular chondrocytes synthesize nitric oxide in response to cytokines and lipopolysaccharide. *J. Immunol.* 147, 3915–20.
- STANDBERG, B. (1964) Intra-articular steroid therapy: a survey and clinical report. *Acta Rheumatol. Scand.* 10, 29–54.
- STEFANOFIC-RACIC, M., MORALES, T.I., TASKIRAN, D., MCINTYRE, L.A. & EVANS, C.H. (1996) The role of nitric oxide in proteoglycan turnover by bovine articular cartilage organ cultures. *J. Immunol.* 156, 1213–20.
- STEINBERG, J. & SLEDGE, C.B. (1983) Synovial factors and chondrocyte mediated breakdown of cartilage: inhibition by hydrocortisone. *J. Orthop. Res.* 1, 13–21.
- TAKANO, T., TAKIGAWA, M. & SUZUKI, F. (1985) Stimulation by glucocorticoids of the differentiated phenotype of chondrocytes and the proliferation of rabbit costal chondrocytes in culture. *J. Biochem.* 97, 1092–100.
- TAKIGAWA, M., TAKANO, T., NAKAGAWA, K., SAKUDA, M. & SUZUKI, F. (1988) Hydrocortisone stimulation of proliferation and glycosaminoglycan synthesis in rabbit craniofacial chondrocytes *in vitro*. *Arch. Oral*

- Biol.* **33**, 839–99.
- TARDIF, G., REBOUL, P., PELLETIER, J.-P., GENG, C., CLOUTIER, J.-M. & MARTEL-PELLETIER, J. (1996) Normal expression of type 1 insulin-like growth factor receptor by human osteoarthritic chondrocytes with increased expression and synthesis of insulin-like growth factor binding proteins. *Arthritis Rheum.* **39**, 968–78.
- TASKIRAN, D., STEFANOVIC-RACIC, M., GEORGESCU, H. & EVANS, C. (1994) Nitric oxide mediates suppression of cartilage proteoglycan synthesis by interleukin-1. *Biochem. Biophys. Res. Commun.* **200**, 142–8.
- TAYLOR, A.M., DANDONA, P., MORRELL, D.J. & PREECE, M.A. (1988) Insulin-like growth factor-1, protein kinase-C, calcium and cyclic AMP: partners in the regulation of chondrocyte mitogenesis and metabolism. *FEBS Lett.* **236**, 33–8.
- TESCH, H.G., HANDLEY, C.J., CORNELL, H.J. & HERRINGTON, A.C. (1992) Effects of free and bound insulin-like growth factors on proteoglycan metabolism in articular cartilage explants. *J. Orthop. Res.* **10**, 14–22.
- TOLLEFSEN, S.E. & THOMPSON, K. (1988) The structural basis for insulin-like growth factor 1 receptor high affinity binding. *J. Biol. Chem.* **263**, 16267–73.
- TOLLEFSEN, S.E., STOSZEK, R.M. & THOMPSON, K. (1991) Interaction of the $\alpha\beta$ dimers of the insulin-like growth factor 1 receptor is required for receptor autophosphorylation. *Biochemistry* **30**, 48–54.
- TRIPPEL, S.B. (1995) Growth factor actions on articular cartilage. *J. Rheumatol.* **22**, 129–32.
- TRIPPEL, S.B., VAN WIJK, J.J., POSTER, M.B. & SVOBODA, M.E. (1983) Characterization of specific somatomedin C receptors on isolated bovine growth plate chondrocytes. *Endocrinology* **112**, 2128–34.
- TRIPPEL, S.B., CHERNAUSEK, S.D., VAN WIJK, J.J., MOSES, A.C. & MANKIN, H.J. (1988) Demonstration of type 1 and type 2 somatomedin receptors in bovine growth plate chondrocytes. *J. Orthop. Res.* **6**, 817–26.
- TRIPPEL, S.B., WROBLEWSKI, J., MAKOWER, A.M., SCHOENFELD, D. & DOCTROW, S.R. (1993) Regulation of growth-plate chondrocytes by insulin-like growth factor 1 and basic fibroblast growth factor. *J. Bone Joint Surg., American Volume* **75**, 177–89.
- TYLER, J.A. (1985) Pig interleukin-1 (catabolin) induces resorption of cartilage and prevents synthesis of cartilage proteoglycan and collagen. *Br. J. Rheumatol.* **24**, 150–5.
- TYLER, J.A. (1989) Insulin-like growth factor 1 can decrease degradation and promote synthesis of proteoglycan in cartilage exposed to cytokines. *Biochem. J.* **260**, 543–8.
- ULLRICH, A., GRAY, A., TAM, A.W., YANG-FENG, T., TSUBOKAWA, M., COLLINS, C., HENREL, W., LE BON, T., KATHURIA, S., CHEN, E., JACOBS, S., FRANCKE, U., RANCHANDRAN, J. & FUJITA-YAMAGUCHI, Y. (1986) Insulin-like growth factor 1 receptor primary structure: comparison with insulin receptor suggests structural determinants that define functional specificity. *EMBO J.* **5**, 2503–12.
- VAN BEUNINGEN, H.M., VAN DER KRAAN, P.M., ARNTZ, O.J. & VAN DEN BERG, W.B. (1993) Protection from interleukin-1 induced destruction of articular cartilage by transforming growth factor β : Studies in anatomically intact cartilage *in vitro* and *in vivo*. *Ann. Rheum. Dis.* **52**, 185–91.
- VAN BEUNINGEN, H.M., VAN DER KRAAN, P.M., ARNTZ, O.J. & VAN DEN BERG, W.B. (1994) Transforming growth factor- β 1 stimulates articular chondrocyte proteoglycan synthesis and induces osteophyte formation in the murine knee joint. *Lab. Invest.* **71**, 279–90.
- VAN DE LOO, A.A.J. & VAN DEN BERG, W.B. (1990) Effects of murine recombinant IL-1 on synovial joints in mice: Quantification of patellar cartilage metabolism and joint inflammation. *Ann. Rheum. Dis.* **49**, 238–45.
- VAN DE LOO, A.A.J., ARNTZ, O.J., OTTERNESS, I.G. & VAN DEN BERG, W.B. (1992) Protection against cartilage proteoglycan synthesis inhibition by anti-interleukin 1 antibodies in experimental arthritis. *J. Rheumatol.* **19**, 348–56.
- VAN DE LOO, A.A.J., ARNTZ, O.J., BAKKER, A.C., VAN LENT, P.L.E.M., JACOBS, M.J.M. & VAN DEN BERG, W.B. (1995a) Role of interleukin-1 in antigen-induced exacerbations of murine arthritis. *Am. J. Pathol.* **146**, 239–49.
- VAN DE LOO, A.A.J., JOOSTEN, L.A.B., VAN LENT, P.L.E.M., ARNTZ, O.J. & VAN DEN BERG, W.B. (1995b) Role of interleukin-1, tumor necrosis factor α and interleukin-6 in cartilage proteoglycan metabolism and destruction: effect of in situ blocking in murine antigen- and zymosan-induced arthritis. *Arthritis Rheum.* **38**, 164–72.
- VAN DEN BERG, W.B. (1991) Impact of NSAID and steroids on cartilage destruction in murine antigen induced arthritis. *J. Rheumatol.* **18**, 122–3.
- VAN DEN BERG, W.B. (1994) Cytokine profile dependent on the type of arthritis: dominant role of IL-1 in cartilage destruction. *Arthritis Rheum.* **37** (suppl), S 193.
- VAN DEN BERG, W.B., KRUIJSEN, M.W.M., VAN DE PUTTE, L.B.A. & ZWARTS, W.A. (1981) Antigen-induced arthritis and zymosan-induced arthritis in mice: Studies on *in vivo* cartilage proteoglycan synthesis and chondrocyte death. *Br. J. Exp. Pathol.* **62**, 308–16.
- VAN DEN BERG, W.B., JOOSTEN, L.A.B., SCHALKWIJK, J., VAN DE LOO, F.A.J. & VAN BEUNINGEN, H.M. (1989) Mechanism of cartilage destruction in experimental arthritis: lack of IGF-1 responsiveness. In *Therapeutic Approaches to Inflammatory Diseases* (edited by LEWIS, A.J., DOHERTY, N.S. & ACKERMAN, N.R.) pp. 47–54. New York: Elsevier.
- VAN DEN BERG, W.B., VAN DE LOO, A.A.J., OTTERNESS, I., ARNTZ, O.J. & JOOSTEN, L.A.B. (1991) *In vivo* evidence for a key role of IL-1 in cartilage destruction in experimental arthritis. *Agents & Actions suppl* **32**, 159–63.
- VAN DEN BERG, W.B., JOOSTEN, L.A.B., VAN DE LOO, F.A.J., DE VRIES, B.J., VAN DER KRAAN, P.M. & VITTERS, E.L. (1992) Drug evaluation on normal and arthritic mouse patellas. In *Articular Cartilage and Osteoarthritis* (edited by KEUTTNER, K.E., SCHLEYER-BACH, R., PEYRON, J.G. & HASCALL, V.C.) pp. 583–96. New York: Raven Press.
- VAN DEN BERG, W.B., JOOSTEN, L.A.B., HELSEN, M.M.A. &

- VAN DE LOO, A.A.J. (1994) Amelioration of established murine collagen-induced arthritis with IL-1 treatment. *Clin. Exp. Immunol.* **95**, 273–343.
- VAN DEN BRANDE, J.L. (1992) Structure of the human insulin-like growth factors: relationship to function. In *Insulin-like growth factors. Structure and biological functions* (edited by SCHOLFIELD, P.N.) pp. 12–37. Oxford: Oxford University Press.
- VAN DER KRAAN, P.M., VITTERS, E.L. & VAN DEN BERG, W.B. (1992a) Inhibition of proteoglycan synthesis by transforming growth factor β in anatomically intact articular cartilage of murine patellae. *Ann. Rheum. Dis.* **51**, 643–7.
- VAN DER KRAAN, P., VITTERS, E. & VAN DEN BERG, W.B. (1992b) Differential effect of transforming growth factor β on freshly isolated and cultured articular chondrocytes. *J. Rheumatol.* **19**, 140–5.
- VAN DER KRAAN, P.M., VITTERS, E.L., VERBUNT, J. & VAN DEN BERG, W.B. (1993) Maintenance of the synthesis of large proteoglycans in anatomically intact murine articular cartilage by steroids and insulin-like growth factor-1. *Ann. Rheum. Dis.* **52**, 734–41.
- VAN NOORDEN, C.J.F., SMITH, R.E. & RASNICK, D. (1988) Cysteine proteinase activity in arthritic rat knee joints and the effects of a selective inhibitor, Z-Phe-AlaCH₂F. *J. Rheumatol.* **15**, 1525–35.
- VAN OOSTVELT, P. & BAUWENS, S. (1990) Quantitative fluorescence in confocal microscopy. *J. Microscopy* **158**, 121–32.
- VERSCHURE, P.J. & VAN NOORDEN, C.J.F. (1990) The effects of interleukin-1 on articular cartilage destruction as observed in arthritic diseases and its therapeutic control. *Clin. Exp. Rheumatol.* **8**, 303–13.
- VERSCHURE, P.J., VAN NOORDEN, C.J.F. & DIJKSTRA, C.D. (1989) Macrophages and dendritic cells during the early stages of antigen-induced arthritis in rats: Immunohistochemical analysis of cryostat sections of the whole knee joint. *Scand. J. Immunol.* **29**, 371–81.
- VERSCHURE, P.J., VAN DER KRAAN, P.M., VITTERS, E.L. & VAN DEN BERG, W.B. (1994a) Stimulation of proteoglycan synthesis by triamcinolone acetonide and insulin-like growth factor in normal and arthritic murine articular cartilage. *J. Rheumatol.* **21**, 920–5.
- VERSCHURE, P.J., JOOSTEN, L.A.B., VAN DER KRAAN, P.M. & VAN DEN BERG, W.B. (1994b) Responsiveness of articular cartilage from normal and inflamed mouse knee joints to various growth factors. *Ann. Rheum. Dis.* **53**, 455–60.
- VERSCHURE, P.J., VAN MARLE, J., JOOSTEN, L.A.B. & VAN DEN BERG, W.B. (1994c) Localization and quantification of the insulin-like growth factor-1 receptor in mouse articular cartilage by confocal laser scanning microscopy. *J. Histochem. Cytochem.* **42**, 765–73.
- VERSCHURE, P.J., JOOSTEN, L.A.B., VAN DE LOO, F.A.J. & VAN DEN BERG, W.B. (1995a) IL-1 has no direct role in the IGF-1 nonresponsiveness during experimentally-induced arthritis. *Ann. Rheum. Dis.* **54**, 976–82.
- VERSCHURE, P.J., VAN MARLE, J., JOOSTEN, L.A.B. & VAN DEN BERG, W.B. (1995b) Chondrocyte IGF-1 receptor expression and responsiveness to IGF-1 stimulation in mouse articular cartilage during various phases of experimentally-induced arthritis. *Ann. Rheum. Dis.* **54**, 645–53.
- VERSCHURE, P.J., VAN MARLE, J., VAN NOORDEN, C.J.F. & VAN DEN BERG, W.B. (1996a) The contribution of quantitative confocal scanning laser microscopy in cartilage research. Chondrocyte insulin-like growth factor-1 receptors in health and disease. *Microsc. Res. Techniq.* (in press).
- VERSCHURE, P.J., JOOSTEN, L.A.B., HELSEN, M.M.A., LAFEVER, F.P.J.G. & VAN DEN BERG, W.B. (1996b) Insulin-like growth factor-1 receptor immunolocalization by confocal scanning laser microscopy in human normal and osteoarthritic cartilage in comparison with proteoglycan synthesis and content. *Br. J. Rheum.* **35**, (in press).
- VERSCHURE, P.J., VAN MARLE, J., JOOSTEN, L.A.B. & VAN DEN BERG, W.B. (1996c) Histochemical analysis of IGF-1 binding sites in normal and experimentally-induced arthritis. *Histochem. J.* **28**, 13–23.
- VON DER MARK, K., KIRSCH, T., NERLICH, A., KUSS, A., WESELOH, G., GLUCKERT, K. & STOSS, H. (1992) Type X collagen synthesis in osteoarthritic cartilage. Indication of chondrocyte hypertrophy. *Arthritis Rheum.* **35**, 806–11.
- WAHL, L.M. & CORCORAN, M.L. (1993) Regulation of monocyte/macrophage metalloproteinase production by cytokines. *J. Periodontol.* **64**, 467–73.
- WARING, P.M., CARROLL, G.J. & KANDIAH, D.A. (1993) Increased levels of leukemia inhibitory factor in synovial fluid from patients with rheumatoid arthritis and other inflammatory arthritides. *Arthritis Rheum.* **36**, 911–5.
- WATANABE, N., ROSENFELD, R.G., HINTZ, R.L., DOLLAR, L.A. & SMITH, R.L. (1985) Characterization of a specific insulin-like growth factor-1/Somatomedin-C receptor on high density primary monolayer cultures of bovine articular chondrocytes: regulation of receptor concentration by somatomedin, insulin and growth hormone. *J. Endocrinol.* **107**, 275–83.
- WEISS, A., LIVNE, E., BRANDEIS, E. & SILBERMANN, M. (1988) Triamcinolone impairs the synthesis of collagen and noncollagen proteins in condylar cartilage of newborn mice. *Calcif. Tissue Int.* **42**, 63–9.
- WEISS, A., LIVNE, E. & SILBERMANN, M. (1990) Growth and differentiation of murine cartilage cells *in vitro* following a short-time exposure to triamcinolone acetonide. *Cell Tissue Res.* **260**, 513–20.
- WELLS, S.K., SANDISON, D., STRICKLER, J. & WEBB, W.W. (1989) Quantitative fluorescence imaging with laser scanning confocal microscopy. In *The handbook of confocal microscopy* (edited by PAWLEY, J.) pp. 27–39. Madison: IMR Press.
- WERNER, S., DUAN, D-S, DE VRIES, C., PETERS, K.G., JOHNSON, D.E. & WILLIAMS, L.T. (1992) Differential splicing in the extracellular region of fibroblast growth factor receptor 1 generates receptor variants with different ligand-binding specificity. *Mol. Cell Biol.* **12**, 89–102.
- WHITE, J. G., AMOS, W. B. & FORDHAM, H. (1987) An evaluation of confocal versus conventional imaging of biological structures by fluorescent light microscopy. *J. Cell. Biol.* **105**, 41–5.
- WILBRINK, B., NIETFELD, J.J. & DEN OTTER, W. (1991) Role

- of TNF α in relation to IL-1 and IL-6 in the proteoglycan turnover of human articular cartilage. *Br. J. Rheumatol.* **30**, 265–71.
- WILDEN, P.A., TREADWAY, J.L., BRIAN, D., MORRISON, D. & PESSIN, J.E. (1989) IGF-1 dependent subunit communication of the IGF-1 holoreceptor: interactions between $\alpha\beta$ heterodimeric receptor halves. *Biochemistry* **28**, 9734–40.
- WILLIAMS, J.M. & BRANDT, K.D. (1985) Triamcinolone hexacetonide protects against fibrillation and osteophyte formation following chemically induced articular cartilage damage. *Arthritis Rheum.* **28**, 1267–74.
- WILLIAMS, R.O., FELDMAN, M. & MAINI, R.N. (1995) Anti tumor necrosis factor α and interleukin-6 in cartilage proteoglycan metabolism and destruction: effect of in situ blocking in murine antigen-induced arthritis. *Arthritis Rheum.* **38**, 164–72.
- WILSON, T. (1989) The role of the pinhole in confocal imaging systems. In *The handbook of confocal microscopy* (edited by PAWLEY, J.) pp. 126–33. Madison: IMR Press.
- WITSCH-PREHM, P., MIEHLKE, R. & KRESSE, H. (1992) Presence of small proteoglycan fragments in normal and arthritic human cartilage. *Arthritis Rheum.* **35**, 1042–52.
- WONG, H.L., COSTA, G.L., LOTZE, M.T. & WAHL, S.M. (1993) Interleukin (IL) 4 differentially regulates monocyte IL-1 family gene expression and synthesis *in vitro* and *in vivo*. *J. Exp. Med.* **177**, 775–81.
- WOOLEY, P.H., WHALEN, J.D., CHAPMAN, D.L., BERGER, A.E., RICHARD, K.A., ASPAR, D.G. & STAITE, N.D. (1993) The effect of an interleukin-1 receptor antagonist protein on type 11 collagen induced arthritis in mice. *Arthritis Rheum.* **36**, 1304–14.
- WROBLEWSKI, J. & EDWALL-ARVIDSSON, C. (1995) Inhibitory effects of basic fibroblast growth factor on chondrocyte differentiation. *J. Bone Mineral Res.* **10**, 735–42.
- XU, W.D., FIRESTEIN, G.C. & TAETLE, R. (1989) Cytokines in chronic inflammatory arthritis. Granulocyte-macrophage colony stimulating factor in rheumatoid synovial effusions. *J. Clin. Invest.* **83**, 876–82.
- YASUNAGA, Y., OCHI, M. & IKUTA, Y. (1995) Localization of insulin-like growth factor-1 binding to rabbit articular chondrocytes in situ. *Eur. J. Exp. Musculoskeletal Res.* **3**, 29–36.
- YEE, D., LEBOVIC, G.S., MARCUS, R.R. & ROSEN, N. (1989) Identification of an alternate insulin-like growth factor β subunit mRNA transcript. *J. Biol. Chem.* **264**, 21439–41.
- YOKONO, K., YONEZAWA, K., HARI, J. & BABA, S. (1989) Regulation of biological functions by insulin and insulin-like growth factor I via their own distinct receptors. *Diabetes Res. Clin. Pract.* **7**, S15–20.
- ZAPF, J., RINDERKNECHT, E., HUMBEL, R.E. & FROESCH, E.R. (1987) Nonsuppressible insulin-like activity (NSILA) from human serum: Recent accomplishments and their physiologic implications. *Metab. Clin. Exp.* **27**, 1803–28.
- ZHANG, B. & ROTH, R.A. (1991) Binding properties of chimeric insulin receptors containing the cysteine rich domain of either the insulin-like growth factor I receptor or the insulin receptor related receptor. *Biochemistry* **30**, 5113–7.