

Cobalt
I 7200

DOI: 10.1002/chin.201237020

37- 020

Synthesis and Structural Characterization of a Tetra-Cobalt(II)-Substituted Sandwich-Type Tungstophosphate, (1,3-H₂dap)₂H₄[Co₄(1,3-Hdap)₂(α -B-PW₉O₃₄)₂]-3H₂O. — The title compound (1,3-dap: 1,3-diaminopropane) is hydrothermally synthesized from a mixture of Na₂WO₄, NaH₂PO₄, CoCl₂, Co(OAc)₂, 1,3-dap, and H₂O (autoclave, 160 °C, 4 d, 16% yield). It crystallizes in the monoclinic space group P2₁/n with Z = 2 (single crystal XRD). The [Co₄(1,3-Hdap)₂(PW₉O₃₄)₂]⁸⁻ anion contains two lacunary α -B-[PW₉O₃₄]⁹⁻ Keggin units linked by a rhomb-like Co₄O₁₄(1,3-dap)₂ cluster. A three-dimensional framework is formed by interaction of protonated 1,3-dap units with the [PW₉O₃₄]⁹⁻ polyanions via hydrogen bonds. — (WANG*, Z.-L.; XI, H.-P.; Z. Naturforsch., B: Chem. Sci. 67 (2012) 5, 495-498, <http://dx.doi.org/10.5560/znb.2012-0077> ; Coll. Chem. Chem. Eng., Henan Univ., Kaifeng 475001, Peop. Rep. China; Eng.) — W. Pewestorf