

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/11685678>

A Convenient Method for the Preparation of (Z)- α,β -Unsaturated Carbonyl Compounds

ARTICLE *in* THE JOURNAL OF ORGANIC CHEMISTRY · MAY 1996

Impact Factor: 4.72 · DOI: 10.1021/jo952098j · Source: PubMed

CITATIONS

85

READS

12

4 AUTHORS, INCLUDING:



R Jason Herr

AMRI

38 PUBLICATIONS 1,049 CITATIONS

SEE PROFILE


A Convenient Method for the Preparation of (*Z*)- α,β -Unsaturated Carbonyl Compounds

Douglass F. Taber,* R. Jason Herr,
Shawn K. Pack,¹ and John M. Geremia¹

Department of Chemistry and Biochemistry, University of Delaware, Newark, Delaware 19716

Received November 28, 1995

(*Z*)- α,β -Unsaturated carbonyl compounds **2** are important building blocks in organic synthesis.² Several procedures have been developed for the general synthesis of these compounds,³ a few of which are known to exhibit substantial (*Z*)-selectivity (>90%).⁴ Ganem has shown that α -diazo esters **1** undergo β -hydride elimination with rhodium(II) acetate to produce (*Z*)-enoates **2**,⁵ but only in cases where no competing 1,5-insertion was possible (eq 1).^{6,7} We have found that the more reactive rhodium(II) trifluoroacetate complex efficiently catalyzes the β -hydride elimination process, at low temperatures, without carbocycle (**3**) formation.



The starting α -diazo esters and α -diazo ketones can be prepared by direct diazo transfer, or by homologation

(1) Undergraduate research participant.

(2) (a) Kelly, S. E. In *Comprehensive Organic Synthesis*, Pergamon Press: Oxford, 1991; Vol. 1, p 729. (b) Boyd, G. V. In *The Chemistry of Enones*; John Wiley & Sons: New York, 1989; Pt. 1, p 281.


(3) For some general routes to α,β -unsaturated esters and ketones, see: (a) Addition/elimination reactions with selenium derivatives: (i) Sharpless, K. B.; Lauer, R. F.; Teranishi, A. Y. *J. Am. Chem. Soc.* **1973**, *95*, 6137. (ii) Reich, H. J.; Wollowitz, S. In *Organic Reactions*; John Wiley & Sons: New York, 1993; Vol. 44, p 1. (iii) Hooz, J.; Oudenes, J. *Synth. Commun.* **1980**, *10*, 667. (b) Addition/elimination reactions with sulfur derivatives: (i) Trost, B. M.; Salzmann, T. N. *J. Am. Chem. Soc.* **1973**, *95*, 6840. (ii) Trost, B. M.; Salzmann, T. N.; Hiroi, K. *J. Am. Chem. Soc.* **1976**, *98*, 4887. (iii) Resek, J. E.; Meyers, A. I. *Tetrahedron Lett.* **1995**, *36*, 7051. (c) Condensation reactions with stabilized phosphonium ylides (review article): Maryanoff, B. E.; Reitz, A. B. *Chem. Rev.* **1989**, *89*, 863. (d) The Horner–Wadsworth–Emmons reaction (review article): see ref 2a. (e) Carbonylations of vinylic halides: Urata, H.; Maekawa, H.; Shigeharu, T.; Fuchikami, T. *J. Org. Chem.* **1991**, *56*, 4320. (f) Carbonylations of vinylmercurials: Larock, R. C.; Narayanan, K. *J. Org. Chem.* **1984**, *49*, 3411.

(4) For (*Z*)-stereoselective syntheses of α,β -unsaturated esters: (a) With bis(trifluoroethyl)phosphonoesters: Still, W. C.; Gennari, C. *Tetrahedron Lett.* **1983**, *24*, 4405. (b) With trimethylphosphonium-propionates: Schmid, G.; Fukuyama, T.; Akasaka, K.; Kishi, Y. *J. Am. Chem. Soc.* **1979**, *101*, 260. (c) With bis(phenyl)phosphonoesters: Ando, K. *Tetrahedron Lett.* **1995**, *36*, 4105. For (*Z*)-stereoselective syntheses of α,β -unsaturated ketones with stabilized ylides: (d) Pietrusiewicz, K. M.; Monkiewicz, J. *Tetrahedron Lett.* **1986**, *27*, 739. (e) Moorhoff, C. M.; Schneider, D. F. *Tetrahedron Lett.* **1987**, *28*, 4721. (f) McKenna, E. G.; Walker, B. J. *J. Chem. Soc., Chem. Commun.* **1989**, 568.

(5) Ikota, N.; Takamura, N.; Young, S. D.; Ganem, B. *Tetrahedron Lett.* **1981**, *22*, 4163.

(6) β -Hydride elimination has been shown to be competitive with rhodium-catalyzed 1,5-insertion reactions: (a) Taber, D. F.; Hennessy, M. J.; Louey, J. P. *J. Org. Chem.* **1992**, *57*, 436. (b) Hennessy, M. J. Ph.D. Dissertation, The University of Delaware, 1989. (c) Galeazzi, E.; Guzman, A.; Pinedo, A.; Saldana, A.; Torre, D.; Muchowski, J. M. *Can. J. Chem.* **1983**, *61*, 454.

of alkyl halides and tosylates. We used the latter approach to prepare the substrates listed in Table 1. Thus, α -alkylation of alkyl acetoacetates **4**^{6a} with alkyl halides and tosylates produced α -substituted β -keto esters **5a–e**, which were converted to the α -diazo esters **1a–e** using one or the other of our established methods (eq 2).⁸



Subjecting the diazo precursors **1a–e** to rhodium(II) trifluoroacetate dimer in CH_2Cl_2 at -78°C for 1 h resulted in the selective formation of (*Z*)- α,β -unsaturated esters **2a–e** in good yields (Table 1). At most, only slight traces of the C–H 1,5-insertion products were detected in the crude mixtures.^{9,10}

Subjecting the α -diazo ketone **1f**^{8c} to the same conditions also resulted in the formation of (*Z*)-enone **2f**, without any trace of cyclization. This illustrates a new regioselective method for the construction of (*Z*)- α,β -unsaturated ketones.

The preference for elimination rather than cyclization under what are only slightly modified conditions is striking. With the very reactive rhodium tetrakis(trifluoroacetate) catalyst, we speculate that the *enthalpy* of activation for these reactions is slight, allowing for *entropy* of activation to dominate. β -Hydride elimination (1,2-insertion), with a smaller entropy of activation, might then proceed more readily than 1,5-insertion, with a larger (more negative) entropy of activation.

This new method for constructing (*Z*)- α,β -unsaturated esters and ketones should be of general utility in organic synthesis. The excellent (>95%)¹⁰ stereoselectivity is especially interesting.


Experimental Section¹¹

Preparation of Methyl 2-Acetylundecanoate (5a). Sodium hydride (60% in mineral oil, 0.80 g, 20.0 mmol) was washed

(7) (a) A rhodium(II) acetate-catalyzed method for the elimination of α -diazo β -methoxy ketones to (*E*)- β -methoxy enones has also been reported: Hudlicky, T.; Olivo, H. F.; Natchus, M. G.; Umpierrez, E. F.; Pandolfi, E.; Volonterio, C. *J. Org. Chem.* **1990**, *55*, 4767. (b) It has also been shown that AgNO_3 -promoted decomposition of acyldiazoethanes can be used to prepare enones: Duggleby, P. McC.; Holt, G.; Hope, M. A.; Lewis, A. *J. Chem. Soc., Perkin Trans. 1* **1972**, 3020.

(8) (a) Taber, D. F.; Ruckle, R. E., Jr.; Hennessy, M. J. *J. Org. Chem.* **1986**, *51*, 4077. (b) Taber, D. F.; You, K.; Song, Y. *J. Org. Chem.* **1995**, *60*, 1093. (c) Taber, D. F.; Gleave, D. M.; Herr, R. J.; Moody, K.; Hennessy, M. J. *J. Org. Chem.* **1995**, *60*, 2283.

(9) However, subjecting the α -diazo ester **1** to the same conditions resulted in exclusive C–C insertion product **ii**.



(10) All unsaturated products (**2a–f**) were clean by ^1H and ^{13}C NMR spectroscopy, showing no trace of the (*E*)-isomer in the crude product mixture before chromatography. Early reactions run at higher temperatures showed significant contamination by the (*E*)-isomer.

(11) For a summary of general experimental procedures see: Taber, D. F.; Houze, J. B. *J. Org. Chem.* **1994**, *59*, 4004.

Table 1. β -Hydride Elimination of α -Diazo Carbonyl Compounds **1**

Entry	α -Diazocarbonyl Compound	α,β -Unsaturated Carbonyl Compound	Yield
1			80%
2			85%
3			91%
4			82%
5			94%
6			92%

three times with petroleum ether and then suspended in 10 mL of dry DME and cooled to 0 °C under nitrogen. Methyl acetoacetate (2.16 mL, 20.0 mmol) was added dropwise to give a solid mass. $n\text{Bu}_4\text{NI}$ (0.21 g, 1.0 mmol) was added, followed by 1-bromononane (1.91 mL, 10.0 mmol), and the reaction was stirred at 80 °C for 20 h. The mixture was cooled to rt, diluted with 10 mL of 5% aqueous HCl and extracted three times with 20 mL of petroleum ether. The combined organic extracts were dried over Na_2SO_4 , concentrated *in vacuo*, and chromatographed to give the α -alkylated β -keto ester **5a** as a colorless oil (1.99 g, 82%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.73; ^1H NMR (250 MHz, CDCl_3) δ 3.62 (s, 3H), 3.33 (t, 1H, J = 7.4 Hz), 2.11 (s, 3H), 1.71 (m, 2H), 1.15 (m, 14H), 0.77 (m, 3H); ^{13}C (62.9 MHz, CDCl_3) δ up: 202.8, 170.1, 31.6, 29.1 (\times 2), 28.0 (\times 2), 27.2 (\times 2), 22.4; down: 59.4, 52.0, 28.4, 13.8; IR (film) 2924, 2855, 1746, 1721, 1645, 1435, 1358, 1152 cm^{-1} ; EI MS m/z (rel intensity) 242 (M^+ , 1), 200 (7), 157 (6), 129 (11), 116 (60), 87 (100), 74 (23), 59 (12).

Ethyl 2-Acetylundecanoate (5b). From 1.00 mL of 1-bromononane was obtained the β -keto ester **5b** as a colorless oil (1.16 g, 86%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.54; ^1H NMR (250 MHz, CDCl_3) δ 4.19 (q, 2H, J = 7.2 Hz), 3.40 (t, 1H, J = 7.5 Hz), 2.22 (s, 3H), 1.82 (m, 2H), 1.25 (m, 15H), 0.87 (m, 3H); ^{13}C (62.9 MHz, CDCl_3) δ up: 203.2, 169.8, 61.1, 31.8, 29.4, 29.2, 28.1, 27.3, 22.6; down: 59.8, 28.6, 14.0; IR (film) 2926, 2855, 1743, 1718, 1644, 1466, 1358, 1242, 1151, 1120 cm^{-1} ; EI MS m/z (rel intensity) 257 (M^+ , 3), 214 (32), 171 (12), 157 (16), 143 (23), 130 (97), 101 (100), 72 (66); HRMS (calcd for $\text{C}_{15}\text{H}_{28}\text{O}_3$) 256.2038, found 256.2058.

tert-Butyl 2-Acetylundecanoate (5c). From 4.78 mL of 1-bromononane was obtained the β -keto ester **5c** as a colorless oil (5.11 g, 72%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.69; ^1H NMR (250 MHz, CDCl_3) δ 3.21 (t, 1H, J = 7.4 Hz), 2.12 (s, 3H), 1.67 (m, 2H), 1.38 (s, 9H), 1.17 (m, 12H), 0.79 (m, 3H); ^{13}C (62.9 MHz, CDCl_3) δ up: 203.0, 168.5, 81.0, 31.2, 28.9, 28.7 (\times 2), 28.6, 27.5, 26.7, 22.0; down: 60.4, 27.9, 27.3 (\times 3), 13.4; IR (film) 2925, 2855, 1714, 1641, 1460, 1368, 1250, 1148, 848 cm^{-1} ; EI MS m/z (rel intensity) 211 (M^+ - $t\text{BuO}$, 1), 158 (1), 112 (3), 103 (13), 102 (15), 98 (16), 57 (100).

Ethyl 2-Acetyl-4-methylheptanoate (5d). From 1.67 g of 2-methylpentanyl iodide was obtained an inseparable mixture of β -keto ester diastereomers **5d** as a colorless oil (1.31 g, 78%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.49; ^1H NMR (250 MHz, CDCl_3) δ 4.17 (q, 2H, J = 7.2 Hz), 3.47 (dd, 1H, J = 5.2, 7.9 Hz), 2.18 (s, 3H), 1.85 (m, 1H), 1.66 (m, 1H), 1.28 (m, 4H), 1.23 (t, 3H, J = 7.2 Hz), 1.07 (m, 1H), 0.84 (d, 6H, J = 6.4 Hz); ^{13}C (62.9 MHz, CDCl_3) δ up: 203.3, 170.0, 61.2, 39.2, 35.3, 19.8; down: 58.1, 30.6, 28.5, 19.3, 14.1, 14.0; IR (film) 2959, 2874, 1746, 1715, 1464, 1359, 1242, 1183, 1025 cm^{-1} ; EI MS m/z (rel intensity) 215 (M^+ + 1, 1), 172 (26), 143 (17), 131 (22), 130 (67), 115 (41), 101 (100), 97 (15), 84 (14), 73 (77); HRMS (calcd for $\text{C}_{12}\text{H}_{23}\text{O}_3$) 215.1647, found 215.1652.

Methyl (E)-2-Acetyl-5-phenylpentanoate (5e). From 3.00 g of cinnamyl bromide was obtained the β -keto ester **5e** as a colorless oil (3.29 g, 93%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.49; ^1H NMR (250 MHz, CDCl_3) δ 7.31 (m, 5H), 6.46 (d, 1H, J = 15.8 Hz), 6.12 (dt, 1H, J = 7.1, 15.8 Hz), 3.74 (s, 3H), 3.62 (t, 1H, J = 7.4 Hz), 2.76 (t, 2H, J = 7.3 Hz), 2.26 (s, 3H); ^{13}C (62.9 MHz, CDCl_3) δ up: 202.1, 169.6, 136.9, 31.5; down: 132.7, 128.4 (\times 2), 127.3, 126.1 (\times 2), 125.6, 57.3, 52.3, 29.2; IR (film) 2926, 2855, 1743, 1718, 1644, 1466, 1358, 1242, 1151, 1120 cm^{-1} ; EI MS m/z (rel intensity) 232 (M^+ , 34), 190 (26), 189 (51), 168 (33), 156 (100), 131 (54), 130 (39), 119 (69), 116 (56), 104 (18), 91 (88), 77 (18); HRMS (calcd for $\text{C}_{14}\text{H}_{16}\text{O}_3$) 232.1099, found 232.1082.

Preparation of Methyl 2-Diazoundecanoate (1a). Method A: Sodium hydride (60% in mineral oil, 0.151 g, 3.78 mmol) was washed three times with petroleum ether and then suspended in 10 mL of dry diethyl ether at 0 °C under nitrogen. To this solution was then added dropwise 0.611 g (2.52 mmol) of β -keto ester **5a** in 3 mL of diethyl ether. After the mixture had stirred for 10 min, 0.915 g (7.56 mmol) of methanesulfonyl azide was added dropwise, and the mixture was allowed to warm to rt over 5 h. The mixture was then diluted with 10 mL of 10% aqueous NaOH and extracted three times with 25 mL of ethyl acetate. The organic layers were then washed with brine, dried over Na_2SO_4 , concentrated *in vacuo*, and chromatographed to give the α -diazo ester **1a** as a yellow oil (0.531 g, 93%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.83; ^1H NMR (250 MHz, CDCl_3) δ 3.67 (s, 3H), 2.22 (t, 2H, J = 7.0 Hz), 1.42 (m, 2H), 1.19 (m, 12H), 0.80 (m, 3H); ^{13}C (62.9 MHz, CDCl_3) δ up: 167.9, 31.8, 29.4, 29.1, 28.6, 27.5, 23.0 (\times 2), 22.6; down: 51.6, 13.9; IR (film) 2926, 2855, 2360, 2079, 1699, 1436, 1351, 1136, 739 cm^{-1} ; EI MS m/z (rel intensity) 199 (M^+ - N_2 , 2), 167 (6), 137 (2), 124 (8), 113 (100), 100 (30), 87 (43), 81 (60), 69 (37).

Ethyl 2-Diazoundecanoate (1b). Method A: From 1.284 g of β -keto ester **5b** was obtained the α -diazo ester **1b** as a yellow oil (0.915 g, 76%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.88; ^1H NMR (250 MHz, CDCl_3) δ 4.14 (q, 2H, J = 7.1 Hz), 2.22 (t, 2H, J = 7.1 Hz), 1.43 (m, 2H), 1.20 (m, 15H), 0.81 (m, 3H); ^{13}C (62.9 MHz, CDCl_3) δ up: 167.6, 60.6, 31.8, 29.4, 29.2 (\times 2), 28.7, 27.5, 22.9, 22.6; down: 14.4, 14.0; IR (film) 2928, 2856, 2084, 1699, 1465, 1371, 1304, 1134, 739 cm^{-1} ; EI MS m/z (rel intensity) 167 (M^+ - N_2 , - OEt, 2), 127 (13), 99 (32), 88 (22), 81 (23), 69 (23), 55 (100).

tert-Butyl 2-Diazoundecanoate (1c). Method A: From 0.658 g of β -keto ester **5c** was obtained the α -diazo ester **1c** as a yellow oil (0.452 g, 73%). TLC R_f (5% ethyl acetate/petroleum ether) = 0.77; ^1H NMR (250 MHz, CDCl_3) δ 2.22 (t, 2H, J = 7.0 Hz), 1.45 (s, 9H), 1.24 (m, 14H), 0.85 (m, 3H); IR (film) 3428, 2957, 2856, 2078, 1694, 1456, 1368, 1133 cm^{-1} ; EI MS m/z (rel intensity) 167 (M^+ - N_2 , - $t\text{BuO}$, 6), 124 (5), 111 (2), 101 (3), 99 (24), 81 (11), 73 (7), 57 (100).

Ethyl 2-Diazo-4-methylheptanoate (1d). Method B: To a solution of the β -keto ester **5d** (1.14 g, 5.3 mmol) and *p*-nitrobenzenesulfonyl azide (2.43 g, 10.6 mmol) in 30 mL of dry CH_2Cl_2 under nitrogen at 0 °C was added dropwise DBU (1.6 mL, 10.6 mmol). The mixture was stirred for 90 min, after which 30 mL of 10% NaOH soln was added, and the mixture was warmed to rt. The crude α -diazo ester was extracted three times with CH_2Cl_2 , washed with brine, dried over Na_2SO_4 , concentrated *in vacuo*, and chromatographed to provide the α -diazo ester **1d** as a yellow oil (0.83 g, 79%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.75; ^1H NMR (250 MHz, CDCl_3) δ 4.22 (q, 2H, J = 7.2 Hz), 2.19 (ddd, 2H, J = 6.1, 14.9, 36.9 Hz), 2.10 (m, 1H), 1.31 (m, 4H), 1.27 (t, 3H, J = 7.2 Hz), 0.92 (d, 6H, J = 6.7 Hz); ^{13}C (62.9 MHz, CDCl_3) δ up: 60.7, 38.4, 30.6, 20.0;

down: 32.4, 19.1, 14.2 ($\times 2$); IR (film) 2959, 2873, 2361, 2080, 1697, 1371, 1323, 1136 cm^{-1} ; EI MS m/z (rel intensity) 170 ($\text{M}^+ - \text{N}_2$, 4), 155 (2), 141 (30), 125 (90), 113 (68), 96 (75), 82 (14), 55 (100).

Methyl (E)-2-Diazo-5-phenylpentanoate (1e). Method B: From 1.86 g of β -keto ester **5e** was obtained the α -diazo ester **1e** as a yellow oil (1.44 g, 83%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.32; ^1H NMR (250 MHz, CDCl_3) δ 7.36 (m, 5H), 6.49 (d, 1H, J = 15.8 Hz), 6.21 (dt, 1H, J = 6.8, 15.8 Hz), 3.80 (s, 3H), 3.22 (d, 2H, J = 6.8 Hz); ^{13}C (62.9 MHz, CDCl_3) δ up: 167.2, 136.5, 26.6; down: 132.6, 128.4 ($\times 2$), 127.4, 126.1 ($\times 2$), 123.8, 51.8; IR (film) 3028, 2952, 2080, 1694, 1598, 1496, 1436, 1337, 1190, 1115, 967, 913, 739, 695 cm^{-1} ; EI MS m/z (rel intensity) 188 ($\text{M}^+ - \text{N}_2$, 14), 157 (16), 129 (100), 128 (88), 117 (6), 102 (17), 77 (22), 64 (18), 51 (28).

Preparation of Methyl (Z)-2-Undecenoate (2a). To a solution of the α -diazo ester **1a** (0.208 g, 0.92 mmol) in 20 mL of dry CH_2Cl_2 at -78°C under nitrogen was added dropwise a solution of rhodium tetrakis(trifluoroacetic acid) dimer (0.0060 g, 0.01 mmol) in 0.5 mL of dry CH_2Cl_2 (precooled to 0°C). The mixture was stirred for 1 h, concentrated *in vacuo*, and immediately chromatographed to provide the (Z)- α,β -unsaturated ester **2a** as a colorless oil (0.182 g, 80%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.85; ^1H NMR (250 MHz, CDCl_3) δ 6.14 (dt, 1H, J = 7.5, 11.5 Hz), 5.68 (d, 1H, J = 11.5 Hz), 3.61 (s, 3H), 2.57 (dq, 2H, J = 1.4, 7.4 Hz), 1.35 (m, 2H), 1.18 (m, 10H), 0.80 (m, 3H); ^{13}C (62.9 MHz, CDCl_3) δ up: 166.5, 31.7, 29.3, 29.3, 29.1, 28.9, 28.8, 22.5; down: 150.7, 119.0, 50.6, 13.9; IR (film) 2926, 2856, 1728, 1645, 1436, 1197, 1175, 819 cm^{-1} ; EI MS m/z (rel intensity) 199 (M^+ , 2), 167 (7), 124 (9), 113 (100), 100 (33), 87 (47), 74 (43); HRMS (calcd for $\text{C}_{12}\text{H}_{22}\text{O}_2$) 198.1620, found 198.1613.

Ethyl (Z)-2-Undecenoate (2b). From 0.401 g of α -diazo ester **1b** was obtained the (Z)- α,β -unsaturated ester **2b** as a colorless oil (0.301, 85%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.93; ^1H NMR (250 MHz, CDCl_3) δ 6.14 (m, 1H), 5.68 (d, 1H, J = 11.5 Hz), 4.08 (q, 2H, J = 7.2 Hz), 2.00 (q, 2H, J = 7.2 Hz), 1.36 (m, 2H), 1.21 (m, 13H), 0.80 (m, 3H); ^{13}C (62.9 MHz, CDCl_3) δ up: 166.3, 59.6, 31.8, 29.3, 29.2, 29.1, 29.0, 28.9, 22.6; down: 150.4, 119.6, 14.2, 14.0; IR (film) 2108, 1722, 1644, 1465, 1134 cm^{-1} ; EI MS m/z (rel intensity) 212 (M^+ , 5), 167 (17), 127 (100), 115 (47), 101 (32), 99 (39), 88 (48), 84 (20); HRMS (calcd for $\text{C}_{13}\text{H}_{24}\text{O}_2$) 212.1776, found 212.1761. Anal. Calcd for $\text{C}_{13}\text{H}_{24}\text{O}_2$: C, 73.54%; H, 11.39%. Found: C, 73.47%; H, 11.25%.

tert-Butyl (Z)-2-Undecenoate (2c). From 0.436 g of α -diazo ester **1c** was obtained the (Z)- α,β -unsaturated ester **2c** as a colorless oil (0.353, 91%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.90; ^1H NMR (250 MHz, CDCl_3) δ 6.09 (m, 1H), 5.64 (d, 1H, J = 11.5 Hz), 2.58 (q, 2H, J = 7.2 Hz), 1.47 (s, 9H), 1.25 (m, 12H), 0.86 (m, 3H); ^{13}C (62.9 MHz, CDCl_3) δ up: 167.0, 80.8, 31.8, 29.4, 29.2, 28.7, 27.5, 22.9, 22.6; down: 148.9, 121.4, 28.3

($\times 3$), 14.0; IR (film) 2958, 2928, 2856, 2361, 1716, 1641, 1367, 1215, 1152, 820 cm^{-1} ; FAB MS m/z (rel intensity) 241 ($\text{M}^+ + 1$, 15), 221 (30), 207 (24), 185 (100), 167 (37), 147 (61); HRMS (calcd for $\text{C}_{15}\text{H}_{29}\text{O}_2$) 241.2168, found 241.2171.

Ethyl (Z)-4-Methyl-2-heptenoate (2d). From 0.2290 g of α -diazo ester **1d** was obtained the (Z)- α,β -unsaturated ester **2d** as a colorless oil (0.1612, 82%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.79; ^1H NMR (250 MHz, CDCl_3) δ 5.92 (dd, 1H, J = 10.2, 11.5 Hz), 5.68 (d, 1H, J = 11.5 Hz), 4.15 (q, 2H, J = 7.2 Hz), 3.50 (m, 1H), 1.27 (t, 3H, J = 7.1 Hz), 1.27 (m, 4H), 0.99 (d, 3H, J = 6.6 Hz), 0.88 (t, 3H, J = 7.2 Hz); ^{13}C (62.9 MHz, CDCl_3) δ up: 166.4, 59.7, 39.2, 20.7; down: 156.0, 118.2, 32.4, 20.2, 14.2, 14.1; IR (film) 2960, 2931, 1722, 1646, 1457, 1417, 1186, 1035, 824 cm^{-1} ; EI MS m/z (rel intensity) 170 (M^+ , 2), 141 (5), 125 (7), 113 (15), 95 (19), 83 (11), 82 (19), 71 (11), 67 (26), 55 (100).

Methyl (Z,E)-5-Phenyl-2-pentenoate (2e). From 0.1610 g of α -diazo ester **1e** was obtained the (Z)- α,β -unsaturated ester **2e** as a colorless oil (0.1315, 94%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.58; ^1H NMR (250 MHz, CDCl_3) δ 8.16 (dd, 1H, J = 12.4, 16.7 Hz), 7.54 (d, 2H, J = 8.1 Hz), 7.34 (m, 3H), 6.77 (q, 2H, J = 15.7 Hz), 5.74 (d, 1H, J = 11.2 Hz), 3.77 (s, 3H); ^{13}C (62.9 MHz, CDCl_3) δ up: 166.8, 136.2; down: 144.9, 141.3, 128.9, 128.6 ($\times 2$), 127.4 ($\times 2$), 124.8, 116.9, 51.1; IR (film) 3024, 2949, 2071, 1712, 1625, 1450, 1392, 1172, 1000, 959, 820 cm^{-1} ; EI MS m/z (rel intensity) 188 (M^+ , 8), 157 (10), 129 (96), 128 (100), 115 (15), 102 (25), 77 (38), 63 (34), 51 (87); HRMS (calcd for $\text{C}_{12}\text{H}_{12}\text{O}_2$) 188.0837, found 188.0829.

(Z)-3-Undecen-2-one (2f). From 0.0960 g of α -diazo ketone **1f** was obtained the (Z)- α,β -unsaturated ketone **2f** as a colorless oil (0.0757, 92%). TLC R_f (10% ethyl acetate/petroleum ether) = 0.74; ^1H NMR (250 MHz, CDCl_3) δ 6.10 (m, 2H, J = 11.7 Hz), 2.59 (q, 2H, J = 7.1 Hz), 2.19 (s, 3H), 1.39 (m, 2H), 1.26 (m, 8H), 0.86 (t, 3H, J = 5.3 Hz); ^{13}C (62.9 MHz, CDCl_3) δ up: 199.3, 31.7, 29.3, 29.2, 29.1 ($\times 2$), 29.0; down: 148.8, 127.0, 31.5, 14.0; IR (film) 2926, 2856, 2361, 1696, 1616, 1558, 1457, 1418, 1355, 1178 cm^{-1} ; EI MS m/z (rel intensity) 168 (M^+ , 2), 153 (2), 125 (3), 110 (10), 97 (100), 84 (21), 71 (19), 69 (43), 55 (44); HRMS (calcd for $\text{C}_{11}\text{H}_{20}\text{O}$) 168.1514, found 168.1517.

Acknowledgment. We thank the National Institutes of Health (GM 42056) for support for this work.

Supporting Information Available: ^1H and ^{13}C spectra for compounds **1a–e**, **2a–f**, and **5a–e** (31 pages). This material is contained in libraries on microfiche, immediately follows this article in the microfilm version of the journal, and can be ordered from the ACS; see any current masthead page for ordering information.

JO952098J