

Total Synthesis of (±)-Welwitindolinone A Isonitrile

Sarah E. Reisman, Joseph M. Ready,[‡] Atsushi Hasuoka, Catherine J. Smith, and John L. Wood*

Sterling Chemistry Laboratory, Department of Chemistry, Yale University, New Haven, Connecticut 06520-8107

Received November 9, 2005; E-mail: john.wood@yale.edu


Welwitindolinone A isonitrile (**1**) is one of several related oxindole-containing alkaloids isolated from blue-green algae by Moore and co-workers in 1994.¹ Among the known welwitindolinones, the only one comprised of a highly functionalized spirocyclobutane oxindole carbon skeleton is **1**, and consequently, it has been postulated to serve as the biosynthetic precursor to the remaining congeners (cf. welwitindolinones C and D in Scheme 1).^{1a} Although isolation and structural characterization of the welwitindolinones was driven by their biological activities (**1** is an antifungal agent), our interest in **1** was piqued by its densely packed and diverse array of synthetically challenging functional groups.² Herein we report an efficient total synthesis of (±)-**1** that exemplifies the strategic and methodological advances molecules of this complexity inspire.

From a retrosynthetic perspective (Scheme 1), we initially envisioned an approach wherein the vinyl isonitrile is derived from a ketone (**2**).³ Although this disconnection reveals a number of standard bond forming strategies, there appeared to be a paucity of methods capable of efficiently delivering the requisite C3 and C12 all-carbon quaternary centers. To address this deficiency, we developed a mild SmI₂-mediated synthesis of spiro-oxindoles (Scheme 1, boxed structures) that enables access to **2** from cyclic urethane **4** via aryl isocyanate **3** (vide infra).⁴ In addition to providing a new oxindole synthesis, these initial investigations demonstrated that known cyclohexadiene **6** could be readily converted to hydroxy-enone **5** (six steps and 56% overall yield), a compound we viewed as a potential precursor to **4**.


Having outlined a general approach and developed a method for assembling the oxindole,⁴ we began considering specific tactics for the stereocontrolled introduction of both the C12 quaternary center and the adjacent neopentyl chlorine found in key intermediate **4**. Rather than address these challenges separately, we devised an approach wherein a chloronium ion induced semi-pinacol rearrangement delivers both simultaneously (see **7** \rightarrow **8** \rightarrow **9** in Scheme 2).⁵ In this event, the C12/C13 relative stereochemistry would be dictated by mechanism (methyl migration *anti* to chloronium ion), while the overall stereocontrol would derive from diastereoface selective formation of the intermediate chloronium ion. Importantly, since both potential migrating groups of tertiary alcohol **7** are methyl, success would depend only upon chloronium ion facial selectivity. Given the rigid, bicyclic nature of **7**, diastereoface selectivity was anticipated; however, of concern was the likelihood that chlorination would occur from the least hindered convex face and result in formation of the *undesired* stereoisomer (not shown). Though somewhat speculative, we hoped to override this intrinsic bias and promote formation of the illustrated chloronium ion (**8**, Scheme 2) by installing a sufficiently large protecting group (R) on the C11 secondary hydroxyl.

Implementing this approach required a tertiary allylic alcohol substrate (**12**) which, as outlined in Scheme 3, was prepared from

Scheme 1



Scheme 2




hydroxy-enone **5** using a sequence that begins with TIPS protection followed by sequential treatment of the derived enone with LHMDs (to transiently protect the cyclic urethane as the lithium amide), L-selectride, and *N*-phenyltriflimide. The derived enol triflate (**10**) was then subjected to Pd-catalyzed CO insertion in the presence of methanol to provide enoate **11**, which was treated with excess methylmagnesium bromide/anhydrous cerium trichloride to deliver **12**. After considerable experimentation with a variety of chlorine sources, we were delighted to find that treatment of **12** with dilute aqueous sodium hypochlorite and cerium trichloride heptahydrate⁶ induces rearrangement of **12** to a *single* chloro-ketone diastereomer (**13**) in 78% isolated yield!

Having successfully implemented the semi-pinacol chemistry, we next installed the C12 vinyl moiety and advanced to cyclization precursor **4** by first desilylating **13** under mild conditions using aqueous fluorosilicic acid⁷ in warm acetonitrile. The resulting β -hydroxy ketone was reduced using tetramethylammonium triacetoxyborohydride⁸ to give a single diastereomer of diol **14**, a crystalline solid which proved suitable for single-crystal X-ray diffraction, thus providing definitive proof that the semi-pinacol rearrangement had furnished the desired relative stereochemistry.⁹

† Current address: Department of Biochemistry, The University of Texas Southwestern Medical Center at Dallas.

Scheme 3




Selective dehydration of the less-hindered C20 alcohol in **14** using Martin sulfurane¹⁰ was followed by oxidation of the remaining alcohol using the Dess-Martin periodinane¹¹ to give cyclization precursor **4** in good yield. Treatment of ketone **4** with DBU induced elimination of CO₂ to furnish the aniline, which was converted in situ (phosgene/Et₃N) to the corresponding isocyanate (**3**, Scheme 1). In accord with our preliminary studies,⁴ exposure of the crude isocyanate to a preformed mixture of SmI₂ and LiCl in THF at -78 °C delivered oxindole **2** in 75% yield with complete diastereocontrol. The stereochemical assignment was confirmed by single-crystal X-ray diffraction and is consistent with bond formation on the less-hindered convex face of the bicyclo[4.2.0]octane skeleton.⁹

With oxindole **2** in hand, we initiated efforts to install the vinyl isonitrile. Unfortunately, **2** proved to be a remarkably unreactive intermediate, and all attempts to convert it to the natural product failed.¹² As a result, we refocused our efforts on an approach wherein the C11 nitrogen would be introduced prior to oxindole formation. To this end, urethane **4** was converted to oxime **15** via a one-pot Boc-protection/CO₂-elimination sequence that was followed by treatment of the derived enone with methoxylamine hydrochloride in pyridine at 65 °C. Unfortunately, α,β -unsaturated oximes of this type proved unreactive toward the previously described SmI₂-mediated reductive cyclization, therefore necessitating development of an alternative method for accessing the oxindole. At this juncture, we recognized that a similar cyclization could potentially be applied to isocyano-isocyanate **18** by taking advantage of the known propensity of isonitriles to undergo α -deprotonation when exposed to strong base (Scheme 4).¹³ Access to the requisite cyclization substrate (**18**) was gained from **15** via sodium cyanoborohydride reduction followed by formylation, SmI₂-mediated N–O bond cleavage, Boc-deprotection, and one-pot dehydration/isocyanate generation (phosgene/Et₃N).^{14,15} Importantly, the reduction of **15** occurs exclusively from the convex face, thus furnishing a pseudoaxial C11 proton that is poised for subsequent deprotonation. With regard to the latter, we were pleased to find that exposure of crude isocyano-isocyanate **18** to LHMDS at -78 °C provided **1** as a *single diastereomer* in moderate yield.

In conclusion, we have developed an efficient synthesis of (\pm)-**1** (2.5% overall yield with an average yield of 81%). Importantly,

Scheme 4



this synthesis inspired the development of new methods for the construction of spiro-oxindoles and improved procedures for chloronium ion induced semi-pinacol rearrangement. To our knowledge, this represents the first example of a chloronium ion induced semi-pinacol rearrangement in the context of a natural product total synthesis and demonstrates the utility of this reaction for the synthesis of α -chloro-quaternary centers, such as those found in the welwitindolinones and related natural products.

Acknowledgment. Financial support was provided by Bristol-Myers Squibb, Yamanouchi, Merck, Amgen, Pfizer, and the NIH (Grant No. 1 R01 CA/GM 93591-01A). S.E.R. thanks Bristol-Myers Squibb for a graduate student fellowship. J.M.R. was the recipient of a NIH postdoctoral fellowship. In addition, we acknowledge and thank C.D. Incarvito for X-ray crystallographic analysis.

Supporting Information Available: Experimental and characterization details (PDF, CIF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (a) Stratmann, K.; Moore, R. E.; Bonjouklian, R.; Deeter, J. B.; Patterson, G. M. L.; Shaffer, S.; Smith, C. D.; Smitka, T. A. *J. Am. Chem. Soc.* **1994**, *116*, 9935–9942. (b) Jimenez, J. I.; Huber, U.; Moore, R. E.; Patterson, G. M. L. *J. Nat. Prod.* **1999**, *62*, 569–572.
- The first total synthesis of **1** was recently reported: Baran, P. S.; Richter, J. M. *J. Am. Chem. Soc.* **2005**, *127*, 15394–15396.
- Barton, D. H. R.; Bowles, T.; Husinec, S.; Forbes, J. E.; Llobera, A.; Porter, A. E. A.; Zard, S. Z. *Tetrahedron Lett.* **1988**, *29*, 3343–3346.
- Ready, J. M.; Reisman, S. E.; Hirata, M.; Weiss, M. M.; Tamaki, K.; Ovaska, T. V.; Wood, J. L. *Angew. Chem., Int. Ed.* **2004**, *43*, 1270–1272.
- Wang, B. M.; Song, Z. L.; Fan, C. A.; Tu, Y. Q.; Chen, W. M., *Synlett* **2003**, 1497–1499 and references therein.
- Moreno-Dorado, F. J.; Guerra, F. M.; Manzano, F. L.; Aladro, F. J.; Jorge, Z. D.; Massanet, G. M. *Tetrahedron Lett.* **2003**, *44*, 6691–6693.
- Pilcher, A. S.; Deshong, P. J. *Org. Chem.* **1993**, *58*, 5130–5134.
- Evans, D. A.; Chapman, K. T.; Carreira, E. M. *J. Am. Chem. Soc.* **1988**, *110*, 3560–3578.
- See Supporting Information for crystallographic data.
- Martin, J. C.; Arhart, R. J. *J. Am. Chem. Soc.* **1971**, *93*, 4327.
- Dess, D. B.; Martin, J. C. *J. Org. Chem.* **1983**, *48*, 4155–4156.
- Screening included (but was not limited to) *N*-nucleophiles, such as NH₂OH, RNHNH₂, RNH₂, RSONH₂, and NH₃, under acidic and basic conditions at ambient and elevated temperatures.
- Hoppe, D. *Angew. Chem., Int. Ed. Engl.* **1974**, *13*, 789–804.
- All attempts to reduce oxime **15** directly to the amine failed to give the desired product.
- Keck, G. E.; Wager, T. T.; McHardy, S. F. *Tetrahedron* **1999**, *55*, 11755–11772.

JA057640S