

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/221860712>

Inverse-Phosphocholine Lipids: A Remix of a Common Phospholipid

ARTICLE *in* JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · FEBRUARY 2012

Impact Factor: 12.11 · DOI: 10.1021/ja210989h · Source: PubMed

CITATIONS

15

READS

33

3 AUTHORS:



Emily K. Perttu

University of California, San Francisco

6 PUBLICATIONS 100 CITATIONS

SEE PROFILE



Aditya Kohli

University of California, San Francisco

9 PUBLICATIONS 134 CITATIONS

SEE PROFILE



Francis Szoka

University of California, San Francisco

236 PUBLICATIONS 23,436 CITATIONS

SEE PROFILE

Published in final edited form as:

J Am Chem Soc. 2012 March 14; 134(10): 4485–4488. doi:10.1021/ja210989h.

Inverse-phosphocholine lipids: A remix of a common phospholipid

Emily K. Perttu, Aditya G. Kohli, and Francis C. Szoka Jr.

Department of Bioengineering and Therapeutic Sciences, University of California, San Francisco, 513 Parnassus Avenue, San Francisco, CA 94143-0912

Abstract

Zwitterionic inverse-phosphocholine (iPC) lipids contain headgroups with an inverted charge orientation relative to phosphocholine (PC) lipids. The iPC lipid headgroup has a quaternary amine adjacent to the bilayer and a phosphate that extends into the aqueous phase. Neutral iPC lipids with ethylated phosphate groups (CPe) and anionic iPC lipids non-ethylated phosphate groups (CP) were synthesized. The surface potential of CPe liposomes remains negative across a broad pH range and in the presence of up to 10 mM Ca^{2+} . CP liposomes aggregate in the presence of Ca^{2+} , but at a slower rate than other anionic lipids above 4 mM Ca^{2+} and are susceptible to hydrolysis by alkaline phosphatase which generates a cationic lipid. CPe liposomes release encapsulated anionic carboxyfluorescein (CF) twenty times more rapidly than PC liposomes, but only release uncharged glucose twice as fast. iPC lipids afford a unique opportunity to investigate the biophysical and bioactivity-related ramifications of a charge inversion at the bilayer surface.

Zwitterionic phosphatidylcholine (PC) lipids are the major component in most bilayer membranes that compartmentalize cells. The PC headgroup is exceptionally well hydrated, maintains its zwitterionic character over a broad pH range and interacts weakly with divalent ions; all properties that make it ideal for its central role in cell membranes. In this report, we investigate how an inversion of the PC headgroup—moving the positively charged quaternary amine adjacent to the bilayer; and extending the anionic phosphate into aqueous interfacial region, alters the chemical and physical properties of the bilayer. We synthesized a family of inverse phosphocholine (iPC) lipids (Fig 1a) and characterized their response to changes in pH, interaction with calcium and permeability to the anionic water-soluble reporter, carboxyfluorescein.

Lipids have three primary regions for potential modification: the hydrophobic tails, the linker region between the tails and the head group, and the hydrophilic headgroup (Fig. 2). Changes to PC lipids at these three sites have been explored by many groups¹; however, to our knowledge, the locations of the choline and phosphate moieties have not been inverted, nor are such modifications observed in nature. Generally, naturally occurring lipids with anionic headgroups, such as phosphatidic acid (PA), phosphatidylserine (PS), phosphatidylinositol (PI), sphingosine-1-phosphate (S1P), and ceramide-1-phosphate (C1P) play more than a just a structural role. PA, PI, S1P, and C1P have been identified as key bioactive agents in a variety of signaling pathways² such as cell growth/apoptosis³, lymphocyte trafficking⁴, chemotaxis⁵, and calcium release^{3b}. Based on the biological roles of these naturally occurring lipids, it is possible that either the CP or the CPe lipids could exhibit similar activities, allowing them to adopt a therapeutic role as well as serving as a structural

Corresponding Author szoka@cgl.ucsf.edu.

Supporting Information. Synthetic protocol, materials, experimental details, DSC data, sample Ca^{2+} aggregation profiles. This material is available free of charge via the Internet at <http://pubs.acs.org>.

component in a liposome bilayer. In addition to their potential for biological activity, iPC lipids are also an exceptionally useful tool for exploring the role of charge-orientation at the bilayer-water interface.

PC lipids have hydrophobic chain length-dependent transition temperatures (T_m), that correspond to the temperature at which the bilayers they form shift from an ordered, crystalline state to a more disordered, fluid state⁶. The T_m for the three saturated chain CPe lipids (DMCPe, DPCPe, and DSCPe) were measured by differential scanning calorimetry following hydration from a thin lipid film and dispersion by brief periods of vortexing. Their T_m were found to be 21.4, 41.2, and 53.7 °C for DMCPe, DPCPe, and DSCPe respectively (See ESI, Table S1). These values are similar to the T_m of their PC-lipid analogs (DMPC, DPPC, and DSPC) which are 24.2, 41.7 and 55.3 °C respectively⁶. Interestingly, sulfo-betaine (SB)⁷ lipids and betaine-like (BL)⁸ lipids, which also have headgroups with a positively charged group adjacent to the hydrophobic region and a negatively charged group that extends away from the bilayer interface, have substantially elevated T_m relative to CP and PC lipids. SB lipids have T_m of; 46 °C (C_{14}), 59 °C (C_{16}), and 68 °C (C_{18}) and BL lipids have T_m of 49 °C (C_{14}), 58 °C (C_{16}), and 67 °C (C_{18}) when measured under the same conditions as the CP lipids. Additionally, the SB and BL lipids do not form liposomes in an aqueous solution with 150 mMNaCl, even above their T_m , whereas dry films of CPe lipids hydrated easily and formed small liposomes upon sonication above their T_m (Fig. 1b). The increased T_m and difficulty in vesicle formation of the SB and BL lipids were attributed to the ion-pairing between adjacent head-groups. The absence of these properties in the CPe lipids indicates that ion-pairing in the headgroups observed in the sulfobetaine liposomes is not present in CPe liposomes.

We investigated how the charge inversion (switching the locations of the quaternary amine and the phosphate relative to PC) of the iPC headgroups would affect the overall surface potential of the liposomes they formed (Fig. 3). The neutral, zwitterionic lipids (DOPC and DOCPe) had surface potentials that were slightly negative at high pH and became barely positive (DOPC) or neutral (DOCPe) around a pH of 2. At low pH, DOPC is a slightly more positive than DOCPe, which suggests that the pK_a for the phosphate of DOCPe may be lower than for DOPC (typically between 2–3)⁹.

The surface potentials for all three liposome solutions made from anionic lipids (POPA, DOCP, and C3-DOCP) start off highly negative and do not differ significantly above pH 5.5. The surface potential of the POPA liposomes, which do not contain a cationic moiety in the headgroup (Fig. 3a), remains negative and constant as the pH decreases. However, because the CP liposomes contain a cationic group, their surface potential should approach neutrality when significantly more than 50% of the phosphate groups have been protonated. The first protonation event in PA headgroups occurs at $pK_{a2} \sim 8$ ¹⁰, but the surface potential for the CP liposomes remains negative well below pH 8, which suggests that the pK_{a2} of the phosphate in the CP is lower than the pK_{a2} of the phosphate in PA. The second protonation event for the phosphate on the CP lipids may occur at a pH similar to the phosphate in DOCPe.

The surface potentials of the two anionic CP liposomes (DOCP and C3-DOCP) do not differ significantly for the majority of the titration, indicating that separating the cationic amine from the phosphate by an additional $-CH_2-$ unit does not change the phosphate pK_a s significantly.

The surface potential measurements revealed subtle differences between the CPe and PC headgroups, but significant differences in the charge properties of CP and PA. Lipids with anionic headgroups like PA and PS coordinate with divalent cations such as Ca^{2+} , resulting

in the aggregation and/or fusion of liposomes containing these anionic lipids¹¹. Calcium-induced aggregation occurs when Ca^{2+} ions form bridges between the outer membranes of two liposomes¹². The rate of bridge formation is affected by the rate of association/dissociation of the Ca^{2+} ions with the phosphate of the lipid headgroup, as well as the rate of dimerization of the two liposomes¹³. The association/dissociation rates and the Ca^{2+} concentration dictate the amount of Ca^{2+} bound to the liposome surface at any given time. The aggregation rate will increase as the Ca^{2+} -phosphate interaction increases and as the Ca^{2+} concentration increases. To determine how the altered electronic properties of the iPC liposomes impact their behavior in the presence of Ca^{2+} , we compared the aggregation of the CPe and CP lipids to traditional zwitterionic and anionic lipids (Fig. 4).

CP liposomes aggregate in the presence of calcium (Fig. 4a and ESI Fig. S2). The aggregation rates of DOCP liposomes differ from what is observed using DOPS or DOPA (Fig. 4a) in that the rate does not significantly increase above 4 mM Ca^{2+} . The increased aggregation rates for DOPS and POPA relative to DOCP at higher Ca^{2+} concentrations and the plateau effect for DOCP suggest some sort of saturation of the Ca^{2+} -DOCP interaction that is not observed for POPA or DOPS and could result from the presence of a formal positive charge in the DOCP headgroup.

Unlike the anionic lipids, DOCPE and DOPC liposomes did not aggregate at 10 mM Ca^{2+} (see ESI Fig. S2). To determine if Ca^{2+} interacts with DOCPE and DOPC differently, the surface potentials of the DOCPE and DOPC liposomes were measured in the presence of various concentrations of Ca^{2+} (fig. 4b). The surface potential of the DOPC liposomes becomes increasingly positive as the Ca^{2+} concentration increases, which is in agreement with previous reports¹⁴. The surface potential of the DOCPE liposomes also becomes more positive as the Ca^{2+} concentration is increased, however; there is no immediate increase between 0 mM Ca^{2+} and 0.1 mM Ca^{2+} and the overall magnitude of the increase is less. Additionally, the surface potential of the DOCPE liposomes remains negative up to 10 mM Ca^{2+} . It is possible that Ca^{2+} interacts more strongly with the surface of the DOPC liposomes due to the location of the anionic group. We hypothesize that because the phosphates in the PC headgroups are located at the bilayer interface, they are better oriented and more ordered than the phosphates in the CPe head-groups which extend into the aqueous phase. Ca^{2+} could then bind more avidly to one or two phosphates in the DOPC liposomes than in the DOCPE liposomes, where the phosphate has a greater freedom of motion. Therefore, it is not the overall surface charge that determines the extent of the Ca^{2+} interaction in DOPC and DOCPE, but the charge of the headgroup moiety adjacent to the hydrophobic bilayer surface.

The Ca^{2+} aggregation and interaction studies demonstrate that the inverted charge orientation changes how the iPC liposomes interact with ions in solution compared to PC liposomes. Monovalent ions (Na^+ and Cl^-) can also coordinate to the bilayer interface and for PC liposomes; Na^+ ions interact preferentially over Cl^- due to the electrostatic attraction between the Na^+ and the anionic phosphate¹⁵. This phenomenon, which attracts positively charged molecules or ions toward and sometimes into the bilayer, results in an increased permeability for Na^+ over Cl^- for PC liposomes¹⁶. In the case of the iPC lipids, the anionic phosphate is replaced with a positively charged quaternary amine, which should have the opposite affect; attracting anionic compounds to the surface and increasing their permeability relative to cations. Therefore, the release rate for an encapsulated anionic compound from an iPC liposome should be greater than for a PC liposome. We tested this hypothesis by measuring the release rates of an encapsulated model anionic compound, carboxyfluorescein (CF) and compared the results to the release of uncharged glucose (Fig. 5).

As predicted, the DPCPe liposomes released CF twenty times faster than DPPC liposomes at 37 °C, while releasing glucose only twice as fast. The difference between these two rate ratios suggests that DPCPe liposomes are preferentially permeability to anions showing that the charge of the headgroup moiety adjacent to the bilayer plays a significant role in the release of charged molecules. Further biophysical studies are required to learn if this is a general phenomenon.

If so, then mixed DPCPe/DPPC liposomes could be used to create a liposome with an adjustable content release profile for charged compounds (ESI Fig. S4). Adjustable release kinetics for liposomal delivery could enhance both existing therapies and aid in the development of new approaches to liposomal drug delivery.

We have synthesized a new class of zwitterionic phospholipids, iPCs, with an inverted charge orientation in the headgroup relative to traditional PC lipids. Both CP and CPe liposomes have negative surface potentials across a broad pH range, but CPe liposomes do not appreciably interact with Ca^{2+} . CP lipids have a terminal phosphate group which is similar to that found in many biologically active lipids. Furthermore, alkaline phosphatase can remove the phosphate from CP lipids (See ESI Table S2) to generate a cationic lipid. This can be exploited to make a biologically sensitive liposome which may be useful for cytoplasmic delivery of encapsulated contents. These iPC lipids provide additional opportunities to study the influence of overall headgroup charge versus headgroup structure (i.e the locations of the anionic and cationic groups) on the intrinsic biological activity of anionic signaling lipids.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We would like to acknowledge Reena Zalpuri, at the UC Berkeley Electron Microscopy Facility, for her help with the TEM experiments and Michael Motion for his assistance with the phosphatase assay.

Funding Sources This work was supported by the National Science Foundation Graduate Fellowship Research Program and by NIH grants EB003008; GM061851 and a UCSF-QB3 Pfizer grant

ABBREVIATIONS

DOPC	1,2-dioleoyl-glycero-3-phosphocholine
POPA	1-palmitoyl-2-oleoyl-glycero-3-phosphatidic acid
DOPS	1,2-dioleoyl-glycero-3-phosphoserine

REFERENCES

- (a) Ma M, Gong Y, Bong D. Lipid Membrane Adhesion and Fusion Driven by Designed, Minimally Multivalent Hydrogen-Bonding Lipids. *Journal of the American Chemical Society*. 2009; 131(46): 16919–16926. [PubMed: 19877659] (b) Rui Y, Wang S, Low PS, Thompson DH. Dipalmenylcholine–Folate Liposomes: An Efficient Vehicle for Intracellular Drug Delivery†. *Journal of the American Chemical Society*. 1998; 120(44):11213–11218. (c) Ali S, Bittman R. Mixed-chain phosphatidylcholine analogues modified in the choline moiety: preparation of isomerically pure phospholipids with bulky head groups and one acyl chain twice as long as the other. *Chemistry and Physics of Lipids*. 1989; 50(1):11–21. [PubMed: 2758523] (d) Arora A, Gupta CM. Novel thermal phase transition behavior of phosphatidylcholine analogs containing 1,2,4-butanetriol as their backbone. *Biochimica et Biophysica Acta (BBA) - Biomembranes*. 1997;

- 1324(1):61–68.(e) Anderson R, Kates M, Volcani BE. Identification of the sulfolipids in the non-photosynthetic diatom *nitzschia alba*. *Biochimica et Biophysica Acta (BBA) - Lipids and Lipid Metabolism*. 1978; 528(1):89–106.(f) Mantsch HH, Cameron DG, Tremblay PA, Kates M. Phosphatidylsulfocholine bilayers: An infrared spectroscopic characterization of the polymorphic phase behavior. *Biochimica et Biophysica Acta (BBA) - Biomembranes*. 1982; 689(1):63–72.(g) Lindh I, Stawinski J. A general method for the synthesis of glycerophospholipids and their analogs via H-phosphonate intermediates. *The Journal of Organic Chemistry*. 1989; 54(6):1338–1342.(h) Rolland J-P, Santaella C, Vierling P. Molecular packing of highly fluorinated phosphatidylcholines in monolayers. *Chemistry and Physics of Lipids*. 1996; 79(1):71–77. [PubMed: 8907245] (i) Huang Z, Szoka FC. Sterol-Modified Phospholipids: Cholesterol and Phospholipid Chimeras with Improved Biomembrane Properties. *Journal of the American Chemical Society*. 2008; 130(46): 15702–15712. [PubMed: 18950160] (j) Chandrakumar NS, Hajdu J. Syntheses of enzyme-inhibitory phospholipid analogs. Stereospecific synthesis of 2-amidophosphatidylcholines and related derivatives. *The Journal of Organic Chemistry*. 1982; 47(11):2144–2147.(k) Makiyama T, Nagasaka N, Houjyo Y, Yamaura E, Nakamura H, Koide Y, Nishida A, Murayama T. Newly synthetic ceramide-1-phosphate analogs; their uptake, intracellular localization, and roles as an inhibitor of cytosolic phospholipase A2 α and inducer of cell toxicity. *Biochemical Pharmacology*. 2010; 80(9):1396–1406. [PubMed: 20688048] (l) Lu RZ, Turcotte JG, Lin WH, Steim JM, Notter RH. Differential scanning calorimetry studies of phosphonolipid analogs of lung surfactant glycerophospholipids. *Journal of Colloid and Interface Science*. 1992; 154(1):24–34.(m) Jing B, Tokutake N, McCullough DH, Regen SL. A Quantitative Assessment of the Influence of Permanent Kinks on the Mixing Behavior of Phospholipids in Cholesterol-Rich Bilayers. *Journal of the American Chemical Society*. 2004; 126(47):15344–15345. [PubMed: 15563137] (n) McNally BA, O'Neil EJ, Nguyen A, Smith BD. Membrane Transporters for Anions That Use a Relay Mechanism. *Journal of the American Chemical Society*. 2008; 130(51):17274–17275. [PubMed: 19035637]
2. (a) Brindley DN, Pilquil C. Lipid phosphate phosphatases and signaling. *Journal of Lipid Research*. 2009; 50(Supplement):S225–S230. [PubMed: 19066402] (b) Boon JM, Lambert TN, Sisson AL, Davis AP, Smith BD. Facilitated Phosphatidylserine (PS) Flip-Flop and Thrombin Activation Using A Synthetic PS Scramblase. *Journal of the American Chemical Society*. 2003; 125(27):8195–8201. [PubMed: 12837089]
 3. (a) Deborah W, S. Stimulation of quiescent 3T3 cells by phosphatidic acid-containing liposomes. *Biochemical and Biophysical Research Communications*. 1987; 145(1):228–233. [PubMed: 3593339] (b) Desai NN, Zhang H, Olivera A, Mattie ME, Spiegel S. Sphingosine-1-phosphate, a metabolite of sphingosine, increases phosphatidic acid levels by phospholipase D activation. *Journal of Biological Chemistry*. 1992; 267(32):23122–8. [PubMed: 1429659] (c) Maceyka M, Payne SG, Milstien S, Spiegel S. Sphingosine kinase, sphingosine-1-phosphate, and apoptosis. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*. 2002; 1585(2–3):193–201.
 4. Takabe K, Paugh SW, Milstien S, Spiegel S. “Inside-Out” Signaling of Sphingosine-1-Phosphate: Therapeutic Targets. *Pharmacological Reviews*. 2008; 60(2):181–195. [PubMed: 18552276]
 5. Chalfant, C.; Poeta, MD. Sphingolipids as signaling and regulatory molecules. Springer Science +Business Media; 2010.
 6. Huang C, Wang Z.-q, Lin H.-n, Brumbaugh EE, Li S. Interconversion of bilayer phase transition temperatures between phosphatidylethanolamines and phosphatidylcholines. *Biochimica et Biophysica Acta (BBA) -Biomembranes*. 1994; 1189(1):7–12.
 7. Perttu EK, Szoka FC. Zwitterionic sulfobetaine lipids that form vesicles with salt-dependent thermotropic properties. *Chemical Communications*. 2011
 8. Kohli, A. G. a. F. C. S., Jr. Synthesis and characterization of zwitterionic betaine-like diacyl lipids with transition temperatures similar to homologous phosphatidylethanolamines. In preparation. 2011
 9. Köhler U, Mantsch HH, Casal HL. Infrared spectroscopic characterization of the interaction of ascorbyl palmitate with phospholipid bilayers. *Canadian Journal of Chemistry*. 1988; 66(4):983–988.
 10. Hauser H. Mechanism of spontaneous vesiculation. *Proceedings of the National Academy of Sciences*. 1989; 86(14):5351–5355.
 11. (a) Wilschut J, Duzgunes N, Fraley R, Papahadjopoulos D. Studies on the mechanism of membrane fusion: kinetics of calcium ion induced fusion of phosphatidylserine vesicles followed

- by a new assay for mixing of aqueous vesicle contents. *Biochemistry*. 1980; 19(26):6011–6021. [PubMed: 7470445] (b) Papahadjopoulos D, Nir S, Düzgünes N. Molecular mechanisms of calcium-induced membrane fusion. *Journal of Bioenergetics and Biomembranes*. 1990; 22(2):157–179. [PubMed: 2139437] (c) Leventis R, Gagne J, Fuller N, Rand R, Silvius J. Divalent Cation Induced Fusion and Lipid Lateral Segregation in Phosphatidylcholine-Phosphatidic Acid Vesicles. *Biochemistry*. 1986; 25(22):6978–6987. [PubMed: 3801406] (d) Chauhan A, Chauhan VPS, Bockerhoff H. Effect of cholesterol on calcium-induced aggregation of liposomes and calcium diphosphatide membrane traversal. *Biochemistry*. 1986; 25(7):1569–1573. [PubMed: 3707894]
12. Portis A, Newton C, Pangborn W, Papahadjopoulos D. Studies on the mechanism of membrane fusion: evidence for an intermembrane calcium(2+) ion-phospholipid complex, synergism with magnesium(2+) ion, and inhibition by spectrin. *Biochemistry*. 1979; 18(5):780–790. [PubMed: 420815]
 13. Lansman J, Haynes DH. Kinetics of a Ca²⁺-triggered membrane aggregation reaction of phospholipid membranes. *Biochimica et Biophysica Acta (BBA) - Biomembranes*. 1975; 394(3):335–347.
 14. Wan P, Zhao Y, Tong H, Yang Z, Zhu Z, Shen X, Hu J. The inducing effect of lecithin liposome organic template on the nucleation and crystal growth of calcium carbonate. *Materials Science and Engineering: C*. 2009; 29(1):222–227.
 15. Hauser H, Oldani D, Phillips MC. Mechanism of ion escape from phosphatidylcholine and phosphatidylserine single bilayer vesicles. *Biochemistry*. 1973; 12(22):4507–4517. [PubMed: 4796045]
 16. Khavrutskii IV, Gorfe AA, Lu B, McCammon JA. Free Energy for the Permeation of Na⁺ and Cl[–] Ions and Their Ion-Pair through a Zwitterionic Dimyristoyl Phosphatidylcholine Lipid Bilayer by Umbrella Integration with Harmonic Fourier Beads. *Journal of the American Chemical Society*. 2009; 131(5):1706–1716. [PubMed: 19146415]




Figure 1.

(a) General structure of PC, CP and CPe lipids (b) TEM image of DPCPe/Cholesterol (6:4) liposomes 65 nm in diameter by light scattering, scale bar = 50 nm.




Figure 2.
Generic lipid and lipid vesicle (liposome) structures





Figure 3.

(a) Illustration of the charge orientation and net charge of naturally occurring phospholipid headgroups (PC, PS, PG, PA) and the CP lipid headgroups (CPe and CP). * The neutral block of the PS headgroup contains a primary amine and carboxylate that are zwitter-neutral at physiological pH. (b) Liposome zeta-potential as a function of solution pH.

**Figure 4.**

(a) Calcium-induced liposome aggregation rates. (b) Shift in zeta potential in the presence of Ca^{2+} .




Figure 5.
CF and glucose release profiles from DPPC and DPCPe liposomes.