

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/6876273>

# Characterization of the Cereulide NRPS $\alpha$ -Hydroxy Acid Specifying Modules: Activation of $\alpha$ -Keto Acids and Chiral Reduction on the Assembly Line

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · SEPTEMBER 2006

Impact Factor: 12.11 · DOI: 10.1021/ja0640187 · Source: PubMed

---

CITATIONS

63

---

READS

27

## 3 AUTHORS:



[Nathan Magarvey](#)

McMaster University

45 PUBLICATIONS 861 CITATIONS

SEE PROFILE



[Monika Ehling-Schulz](#)

University of Veterinary Medicine in Vienna

73 PUBLICATIONS 1,983 CITATIONS

SEE PROFILE



[Christopher Walsh](#)

Harvard Medical School

307 PUBLICATIONS 17,564 CITATIONS

SEE PROFILE

## Characterization of the Cereulide NRPS $\alpha$ -Hydroxy Acid Specifying Modules: Activation of $\alpha$ -Keto Acids and Chiral Reduction on the Assembly Line

Nathan A. Magarvey,<sup>†</sup> Monika Ehling-Schulz,<sup>‡</sup> and Christopher T. Walsh<sup>\*,†</sup>

Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, 240 Longwood Avenue, Boston, Massachusetts 02115, and Department of Biosciences, Technische Universitat Munchen, WZW, D-85354 Freising, Germany

Received June 7, 2006; E-mail: christopher\_walsh@hms.harvard.edu

Several nonribosomal depsipeptide natural products are composites of  $\alpha$ -hydroxy acid and  $\alpha$ -amino acid monomers.<sup>1</sup> Cereulide (ces) (1), the emetic toxin from the human pathogen *Bacillus cereus*, and valinomycin (vlm) (2), from *Streptomyces* spp., are closely related macrocyclic K<sup>+</sup> ionophores.<sup>2,3</sup> The macrocyclic core of each natural product contains alternating peptide (six) and ester (six) bonds and their cyclododecdepsipeptide structures consist of a tetradepsipeptide unit repeated three times. For ces (1) the tetradepsipeptide is D- $\alpha$ -hydroxyisocaproyl-D-alanyl-L- $\alpha$ -hydroxyisovaleryl-L-valyl; for vlm (2), the repeat unit is D- $\alpha$ -hydroxyisovaleryl-D-valyl-L-lactyl-L-valyl (Figure 1).  $\alpha$ -Hydroxy acids of ces (1) and vlm (2) are known to originate from their respective  $\alpha$ -amino acid pools, but it has been unclear if hydroxy acids or amino acids are the monomer species that are selected and elongated by the nonribosomal depsipeptide enzymatic machinery.<sup>4,5</sup> In fact, as we show in this work it, is the corresponding  $\alpha$ -keto acids that get selected, tethered as pantetheinyl thioesters and reduced to  $\alpha$ -hydroxyacyl-S-pantetheinyl carrier protein intermediates in cis reductase domains.

Recently, the *ces* and *vlm* biosynthetic gene clusters have been cloned and sequenced.<sup>6,7</sup> Both of the *ces* NRPSs CesaA (3391 aa) and CesB (2681 aa) begin with a module of the following domain arrangement, adenylation (A)-reductase/dehydrogenase (KR in Figure 1) peptidyl carrier protein (PCP).<sup>6</sup> The Vlm1 NRPS predicted from the *S. tsusimaensis* ATCC 15141 *vlm* biosynthetic gene cluster is annotated as having a module with the following domains: A-transaminase (TA)-dehydrogenase-PCP.<sup>7</sup> We have independently sequenced a second *vlm* cluster from *S. levoris* A-9<sup>9</sup> (accession # DQ640825) (Figure S1, Supporting Information) and no such TA domain was found. Comparison of CesaA, *S. levoris* A-9 Vlm1, and the *S. tsusimaensis* Vlm1 revealed them to be similar in size, sequence, and overall content (Figure S2). From our reanalysis, no TA domain exists within *S. tsusimaensis*, but a stretch of residues preceding the reductase/dehydrogenase domains of CesaA/B and Vlm1/2 from *S. levoris* A-9 bears similarity to proposed "spacer regions" of melithiazol and myxothiazol type I polyketide synthases (PKSs).<sup>8</sup> Thus, both CesaA/B NRPSs and Vlm1/2 NRPSs have similar domain organization (Figure 1). Most notable is the presence of predicted KR domains which are similar to  $\beta$ -ketoacyl reductase domains based on sequence analysis to KR domains found in PKS assembly lines, and further, these KR domains are inserted into the A domains (Figure S3). To characterize the CesaA and CesB modules for monomer recognition and establish how the *ces* NRPS incorporates  $\alpha$ -hydroxy acids, DNA fragments encoding the first CesaA and B modules (A-KR-PCP) were obtained by PCR from *B. cereus* F4810/72 genomic DNA and cloned into *E. coli* expression vectors. The *cesA* fragment was cloned into pTrcHis-TOPO TA vector (Invitrogen), and the *cesB* fragment was cloned





Figure 1. Cereulide and valinomycin and their NRPSs.

into the *NcoI* and *XhoI* sites of pET28b (Novagen). The CesaA module was heterologously expressed in *E. coli* BL21 (DE3) as a doubly His-tagged (C and N-termini) protein, and the CesB module was expressed as an N-terminally His-tagged protein. Both proteins were purified to over 90% purity using Ni-NTA affinity and gel exclusion chromatographies (Figure S4). To determine the substrate of the A domains contained within the His-tagged CesaA and CesB (A-KR-PCP) modules, the radioactive ATP-PPi exchange assay was used as a measure of reversible acyl-AMP formation. Test substrates for CesaA were L- and D-isoleucine (Ile), L- and D- $\alpha$ -hydroxy isocaproic acid (L- and D-HIC), and  $\alpha$ -ketoisocaproic acid (KIC). For CesB, test substrates were L- and D-valine (Val), both enantiomers of  $\alpha$ -hydroxy isovaleric acid (L- and D-HIV), as well as  $\alpha$ -ketoisovaleric acid (KIV). Both CesaA and CesB were found to preferentially activate  $\alpha$ -keto acids over their corresponding  $\alpha$ -hydroxy acids and  $\alpha$ -amino acids, establishing them as novel  $\alpha$ -keto acid activating A domains (Figure 2, the CesaA and B predicted NRPS codes are shown in Table S1). A trichloro- $\alpha$ -ketoisocaproic acid activating A domain was recently described in a barbamide NRPS.<sup>10</sup>

Next, the ability of CesaA and CesB A domains to transacylate  $\alpha$ -keto acyl-adenylates onto their adjacent PCPs was tested. CesaA and CesB were converted from their purified apo-forms to their respective holo-forms by CesP, a phosphopantetheinyl transferase encoded by the *ces* cluster. CesP was also expressed and purified from *E. coli* as a His-tagged variant. Radio-labeled KIC and KIV were prepared from [<sup>14</sup>C]-labeled Ile and Val, respectively, using snake venom L-amino acid oxidase. In each case the [<sup>14</sup>C]- $\alpha$ -keto acids were loaded onto their cognate holo-PCPs by both the CesaA and B A domains as assayed by radioactive incorporation into proteins via TCA-precipitation and subsequent liquid scintillation counting. The fate of the  $\alpha$ -ketoacyl-S-PCPs was then examined.

<sup>†</sup> Harvard Medical School.

<sup>‡</sup> Technische Universitat Munchen.



**Figure 2.** (A) ATP-PPi exchange assay with CesA A domain substrates; (B) radio-TLC of CesA products (lane 1,  $^{14}\text{C}$  labeled  $\alpha$ -KIC; lane 2, D-, L-HIC standard; lane 3, product of CesA KR domain); (C) chiral radio-HPLC of CesA KR domain product (green) with cold D-HIC and L-HIC (black); (D) ATP-PPi exchange assay with CesB A domain; (E) radio-TLC of CesB products (lane 1,  $^{14}\text{C}$  labeled  $\alpha$ -KIV; lane 2, D-, L-HIV standard; lane 3, product of CesB KR domain); (F) chiral radio-HPLC of CesB KR domain product (blue) and D-HIV and L-HIV (black).

In Figure 1 (and also Figure S3) we have noted the prediction of KR domains within CesA and B  $\alpha$ -keto acid A domains. These are distinct from the KR domains found in PKS modules in two ways. First the PKS domains are  $\beta$ -ketoacyl-S-carrier protein reductases ( $\beta$ -KRs) while the CesA/B (and presumably VlmA/B) KR domains would be  $\alpha$ -ketoacyl-S-carrier protein reductases ( $\alpha$ -KRs). Second, while PKS  $\beta$ -KR domains are downstream of acyltransferase (AT) domains in modules where they are found, the reductase domains present in CesA and CesB are embedded within the  $\alpha$ -keto acid A domains between A domain motifs A8 and A9 (schematized in Figure 1 and shown in Figure S3). The A domain region spanning sequence motifs A8–A9 is a flexible loop,<sup>11</sup> which must serve as a “stuffer region”, accommodating diverse catalytic domains such as *N*- and *C*-methyltransferases, decarboxylase domains, and now  $\alpha$ -KR domains. To establish  $\alpha$ -KR domain function, [ $^{14}\text{C}$ ]-labeled  $\alpha$ -keto acyl-S-PCPs were generated as above on the CesA and CesB PCP domains, followed by the addition of NADPH. The thioester-bound products were liberated by TycF, the tyrocidine type II thioesterase and analyzed by radio-TLC versus KIV, KIC, L,D-HIV and L,D-HIC standards. The product obtained from the CesA module had an  $R_f$  identical to HIC, whereas the CesB product had an  $R_f$  equivalent to HIV (Figure 2). Chirality of  $\alpha$ -hydroxy acids was established by radio-HPLC using chiral column chromatography of underivatized acids. The CesA  $\alpha$ -hydroxy acid was identified as D-HIC, whereas L-HIV was the sole enantiomer formed by the CesB KR domain (Figure 2). Product chirality corresponds with the stereochemistry (D-HIC and L-HIV) of such residues within 1. Thus the  $\alpha$ -KR domain within CesA is a D-reductase and within CesB is an L-reductase. Structural analyses will be warranted to evaluate chirality and make comparisons to the D- and L-specific (more commonly referred to as *R*- and *S*-specific)  $\beta$ -KR domains in PKSs. Every other bond within the *ces* tetradepsipeptide unit (i.e., D- $\alpha$ -hydroxyisocaprolyl-D-alanyl-L- $\alpha$ -hydroxyisovaleryl-L-valyl) is an ester. The elongation module for the L- $\alpha$ -hydroxyisovaleryl monomer is the CesB A–KR–PCP and the condensation (C) domain found in trans at the C-terminus of CesA (Figure 1). We propose that the terminal CesA C domain is an ester synthase rather than an amide synthase working in trans with the HIC-D-Ala peptide as donor and the  $\alpha$ -hydroxy group of the CesB thioester-bound L-HIV as nucleophilic acceptor. Elongation to the D-HIC-D-Ala-

L-HIV-L-Val chain is presumably followed by transfer to the TE domain, while a second tetradepsipeptide builds up on the adjacent PCP of CesB. We anticipate the TE domain then acts like the enterobactin synthetase TE:<sup>12</sup> condensing two tetradepsipeptides to an octadepsipetidyl-O-TE and then a dodecadepsipetidyl-O-TE prior to macrolactonization to the cyclic twelve residue product cereulide.

The  $\alpha$ -hydroxy acid incorporation strategy of  $\alpha$ -keto acid selection, activation, tethering, and in situ chiral reduction should also hold for the *vlm* NRPSs (Figure 1), with the telltale  $\alpha$ -KR embedded in the A domain. Two strategies appear to be utilized for  $\alpha$ -hydroxy acid monomer incorporation into natural products via NRPS assembly lines: (1) The bacterial NRPS logic exemplified by the *ces* NRPSs and (2) the proposed fungal NRPS logic (e.g., enniatin and PF1022A)<sup>13a,b</sup> where an A domain activates and directly tethers the  $\alpha$ -hydroxy acid generated from a nonassembly line associated  $\alpha$ -keto acid dehydrogenase.<sup>13c</sup> As yet there is no evidence for a third variant where an aminoacyl-S-carrier protein would be oxidatively deaminated and then reduced to a hydroxy acyl moiety.

It will be of interest to see how many other nonribosomal products with alternating ester and amide bonds are generated by this coupled  $\alpha$ -keto acid activation, tethering, and reduction logic. Three of the domains in the *ces* and *vlm* NRPSs are worthy of detailed study: the A domains for their ability to generate  $\alpha$ -keto acyl-AMPs, the  $\alpha$ -KRs that do chiral reduction of  $\alpha$ -ketoacyl-S-carrier proteins as opposed to the  $\beta$ -ketoacyl-S-carrier proteins in PKS action, and the C domains that are chiral ester synthases rather than amide synthases.

**Acknowledgment.** Support by NIH Grant GM049338 (C.T.W.).

**Note Added after ASAP Publication.** After this paper was published ASAP on July 28, 2006, further changes were introduced to clarify sentence 3 of paragraph 2 and to correct the Vlm notation in sentences 5 and 6 of paragraph 2. The corrected version was published on the same date.

**Supporting Information Available:** Details of protein preparation, enzymatic, and chemical analysis. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) Ballard, C. E.; Yu, H.; Wang, B. *Curr. Med. Chem.* **2002**, *9*, 471–498 and references therein.
- (2) (a) Agata, N.; Mori, M.; Ohta, M.; Suwan, S.; Ohtani, I.; Isobe, M. *FEMS Microbiol. Lett.* **1994**, *121*, 31–34. (b) Agata, N.; Ohta, M.; Mori, M.; Isobe, M. *FEMS Microbiol. Lett.* **1995**, *129*, 17–20.
- (3) (a) Brockmann, H.; Geeren, H. *Justus Liebigs Ann. Chem.* **1957**, *603*, 216–232. (b) Ogata, E.; Rasmussen, H. *Biochemistry*, **1966**, *5*, 57–66.
- (4) Kuse, M.; Franz, T.; Koga, K.; Suwan, S.; Isobe, M.; Agata, N.; Ohta, M. *Bioorg. Med. Chem. Lett.* **2000**, *10*, 735–739.
- (5) (a) MacDonald, J. C.; Slater, G. P. *Can. J. Biochem.* **1968**, *46*, 573–578. (b) Ristow, H.; Salnikow, J.; Kleinkauf, H. *FEBS Lett.* **1974**, *42*, 127–130. (c) Anke, T.; Lipmann, F. *FEBS Lett.* **1977**, *82*, 337–340.
- (6) Cheng, Y. Q. *ChemBiochem* **2006**, *7*, 471–477.
- (7) (a) Ehling-Schulz, M.; Vukov, N.; Schulz, A.; Shaheen, R.; Andersson, M.; Martlbauer, E.; Scherer, S. *Appl. Environ. Microbiol.* **2005**, *71*, 105–113. (b) Ehling-Schulz, M.; Fricker, M.; Grallert, H.; Riek, P.; Wagner, M.; Scherer, S. *BMC Microbiol.* **2006**, *6*, 20.
- (8) (a) Silakowski, B.; Schairer, H. U.; Ehret, H.; Kunze, B.; Weinig, S.; Nordsiek, G.; Brandt, P.; Blocker, H.; Hoffe, G.; Beyer, S.; and Muller, R. *J. Biol. Chem.* **1999**, *274*, 37391–37399. (b) Weinig, S.; Hecht, H. J.; Mahmud, T.; and Muller, R. *Chem. Biol.* **2003**, *10*, 939–952.
- (9) Perkins, J. B.; Guterman, S. K.; Howitt, C. L.; Williams, V. E., II.; Pero, J. *Bacteriol.* **1990**, *172*, 3108–3116.
- (10) Flatt, P. M.; O'Connell, S. J.; McPhail, K. L.; Zeller, G.; Willis, C. L.; Sherman, D. H.; Gerwick, W. H. *J. Nat. Prod.* **2006**, *69*, 938–944.
- (11) Conti, E.; Stachelhaus, T.; Marahiel, M. A.; Brick, P. *EMBO J.* **1997**, *14*, 4174–4183.
- (12) Shaw-Reid, C. A.; Kelleher, N. L.; Losey, H. C.; Gehring, A. M.; Berg, C.; Walsh, C. T. *Chem. Biol.* **1999**, *6*, 385–400.
- (13) (a) Gliniski, M.; Urbanke, C.; Hornbogen, T.; Zocher, R. *Arch. Microbiol.* **2002**, *178*, 267–273. (b) Weckwerth, W.; Miyamoto, K.; Iinuma, K.; Krause, M.; Gliniski, M.; Storm, T.; Bonse, G.; Kleinkauf, H.; Zocher, R. *J. Biol. Chem.* **2000**, *275*, 17909–17915. (c) Lee, C.; Gorisch, H.; Kleinkauf, H.; Zocher, R. *J. Biol. Chem.* **1992**, *267*, 11741–11744.

JA0640187