Am Chem Soc. Author manuscript; available in PMC 2010 August 5.

Published in final edited form as:

J Am Chem Soc. 2009 August 5; 131(30): 10366–10367. doi:10.1021/ja904124b.

# All Carbon Quaternary Centers *via* Ruthenium-Catalyzed Hydroxymethylation of 2-Substituted Butadienes Mediated by Formaldehyde: Beyond Hydroformylation

Tomas Smejkal<sup>a,b,c</sup>, Hoon Han<sup>c</sup>, Bernhard Breit<sup>a,b,\*</sup>, and Michael J. Krische<sup>b,c,\*</sup>

- <sup>a</sup> Albert-Ludwigs-Universität Freiburg, Institut für Organische Chemie und Biochemie, 79104 Freiburg, Germany
- <sup>b</sup> Albert-Ludwigs-Universität Freiburg, Freiburg Institute for Advanced Studies (FRIAS), 79104 Freiburg, Germany
- <sup>c</sup> University of Texas at Austin, Department of Chemistry and Biochemistry, Austin, TX 78712

### **Abstract**

Ruthenium catalyzed transfer hydrogenation of 2-substituted dienes **1a-1i** in the presence of paraformaldehyde results in reductive coupling at the 2-position to furnish products of hydroxymethylation **3a-3i**, which embody all carbon quaternary centers. Reductive coupling of diene **1g** to paraformaldehyde under standard conditions, but employing either *deuterio*-paraformaldehyde or *d*<sub>8</sub>-isopropanol, or both *deuterio*-paraformaldehyde corroborate a catalytic mechanism involving rapid and reversible diene hydrometallation with incomplete regioselectivity in advance of C-C coupling. These present method provides an alternative to the hydroformylation of conjugated dienes, for which efficient, regioselective catalytic systems remain undeveloped.

Hydroformylation is the largest volume application of homogenous metal catalysis and the prototypical C-C bond forming hydrogenation. Whereas alkene hydroformylation is well developed, the hydroformylation of conjugated dienes has proven especially challenging. As part of a broad program aimed at the development of hydrogen-mediated C-C bond formations beyond hydroformylation, one of the present authors reported ruthenium catalyzed reductive couplings of carbonyl compounds to various unsaturates, including dienes, allenes, alkynes, and enynes. In lieu of efficient protocols for diene hydroformylation, the ruthenium catalyzed reductive coupling of dienes to paraformaldehyde, an abundant C1-

mkrische@mail.utexas.edu; bernhard.breit@chemie.unifreiburg.de.

feedstock, was investigated. Here, we report that ruthenium catalyzed transfer hydrogenation of 2-substituted dienes in the presence of paraformaldehyde delivers products of reductive C-C coupling in good yield. Remarkably, and in contrast to prior work on diene-carbonyl reductive coupling,  $^{4-8}$  conditions that promote interconversion of  $\pi$ -allyl **A** to the isomeric  $\pi$ -allyl **B** were identified,  $^{9}$  enabling C-C coupling at the 2-position of the diene to furnish products incorporating all carbon quaternary centers.

$$\begin{array}{c} \text{Me} \\ \text{Me} \\ \text{Me} \\ \text{Me} \\ \text{Ne} \\ \text{Ho} \\$$

Initial studies focused on the reductive coupling of myrcene  ${\bf 1a}$  to paraformaldehyde. Upon an assay of our previously disclosed conditions,  $^{4a,b}$  the catalyst prepared in situ from RuHCl(CO) (PPh<sub>3</sub>)<sub>3</sub> and *rac*-BINAP was most effective, providing an 18% isolated yield of C-C coupling product. Surprisingly, this product appeared as an equimolar mixture of the anticipated adduct  ${\bf 2a}$  and its regioisomer  ${\bf 3a}$ , wherein coupling occurs at the substituted position of the diene to furnish the all carbon quaternary center. It was postulated that product  ${\bf 3a}$  forms through isomerization of  $\pi$ -allyl isomer  ${\bf A}$  to  $\pi$ -allyl  ${\bf B}$  by way of reversible  $\beta$ -hydride elimination-diene hydrometallation. Based on this hypothesis, ruthenium catalysts that embody greater cationic character were assayed, as coordinative unsaturation should promote  $\beta$ -hydride elimination, potentially accelerating isomerization. Indeed, upon an assay of counterions, it was found that RuH(O<sub>2</sub>CC<sub>7</sub>F<sub>15</sub>)(CO)(dppb)(PPh<sub>3</sub>), which is prepared in situ from RuH<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>3</sub> and HO<sub>2</sub>CC<sub>7</sub>F<sub>15</sub>; <sup>10</sup> provides a 76% isolated yield of C-C coupling product as a 1:4 mixture of isomers  ${\bf 2a}$  and  ${\bf 3a}$ , respectively, in the presence of dppb.

It was hypothesized that the relative energies of the competing transition structures for carbonyl addition dictate the distribution of products **2** and **3**. If one assumes intervention of a chair-like transition structure, the path to isomers 2 mandates pseudo-axial orientation of the diene 2-substituent (Scheme 1). Hence, a larger 2-substituent should disfavor formation of isomers **2**. Indeed, exposure of the cyclohexyl-substituted diene **1b** to the aforementioned reaction conditions results in formation of the primary neopentyl alcohol **3b** as a single regioisomer. Branching directly adjacent to the 2-position is not required, as illustrated by formation of adducts **3c** and **3e**. However, sterically demanding groups are required at oxygen and nitrogen, respectively, to maintain complete levels of regioselectivity. To probe potential for substrate-induced diastereoselectivity, dienes **1d** and **1f**, which possess a preexisting stereogenic center, were subjected to standard reaction conditions. However, the resulting neopentyl alcohols were formed as equimolar mixtures of diastereomers. Finally, as demonstrated by formation of adducts **3g-3i**, <sup>11</sup> 2-aryl-1,3-butadienes are subject to highly regioselective hydroxymethylation (Table 1).

To gain further mechanistic insight, isotopic labeling studies were undertaken. Diene 1g was subjected to three separate experiments employing *deuterio*-paraformaldehyde,  $d_8$ -isopropanol and both *deuterio*-paraformaldehyde and  $d_8$ - isopropanol under otherwise standard conditions (Table 2). The observed patterns of deuterium incorporation exclude pathways involving ruthenium catalyzed hydroformylation,  $^{12}$  potentially enabled through decomposition of paraformaldehyde to form syngas (CO/H<sub>2</sub>). Rather, these data are consistent with a scenario involving diene hydrometallation- $\beta$ -hydride elimination at different positions of the diene by way of intermediates **A–D**. Formaldehyde addition from the primary  $\sigma$ -allyl haptomer derived from **B** through a chair-like transition structure is postulated to provide isomers **3** (Scheme 1).

As previously discussed, strain associated with pseudo-axial orientation of large diene 2-substituents appears to disfavor formation of isomers 2. In contrast, the transition structure en route to isomers 3 involves pseudo-equatorial orientation of the diene 2-substituents and projection of these groups into open volumes of space.

Formaldehyde hemiacetals mediate reductive coupling in competition with isopropanol.  $^{1}$ H NMR analysis of crude reaction mixtures reveal both acetone and isopropyl formate. Additionally, in the absence of isopropanol, but under otherwise standard conditions, diene 1g is converted to formate esters 2g-formate and 3g-formate in 42% isolated yield as a 1:4 ratio of regioisomers, respectively. The difference in crystallinity and, hence, reactivity between paraformaldehyde and *deuterio*-paraformaldehyde may account for the observed drop in deuterium incorporation for "14" upon use of *deuterio*-paraformaldehyde and 18-isopropanol compared to paraformaldehyde and 18-isopropanol.

$$R^{1} \xrightarrow{\text{LnRu-H}} R^{1} \xrightarrow{\text{RuLn}} R^{1} \xrightarrow{\text{RuLn}} (CH_{2}O)_{n} \xrightarrow{\text{64\% Yield}} (GH_{2}OH) \xrightarrow{\text{Me}} ($$

In summary, ruthenium catalyzed transfer hydrogenation of 2-substituted dienes in the presence of paraformaldehyde results in reductive coupling at the 2-position to furnish products of hydroxymethylation that contain all carbon quaternary centers. This process represents an alternative to 1,3-diene hydroformylation, for which efficient regioselective catalytic systems remain undeveloped.

# **Supplementary Material**

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

Acknowledgment is made to the Robert A. Welch Foundation, NIH-NIGMS (RO1-GM069445) and the Freiburg Institute for Advanced Studies (FRIAS) for partial support of this research.

# References

- For selected reviews on hydroformylation, see: (a)Weissermel K, Arpe H-J. Industrial Organic Chemistry Wiley-VCHWeinheim2003:127–144.144(b)van Leeuwen PWNM, Claver C. Rhodium catalyzed hydroformylation. KluverDordrecht2000(c)Breit B, Seiche W. Synthesis 2001:1–36.36
- 2. Hydroformylation of conjugated dienes typically occurs in low yield to provide complex mixtures, see: (a)Clement WH, Orchin M. Ind Eng Chem Prod Res Dev 1965;4:283.(b)Fell B, Bahrmann H. J Mol Catal 1977;2:211.(c)Bahrmann H, Fell B. J Mol Catal 1980;8:329.(d)Botteghi C, Branca M, Saba A. J Organomet Chem 1980;184:C17.(e)van Leeuwen PWNM, Roobeek CF. J Mol Catal 1985;31:345. (f)Chalchat JC, Garry RPh, Lecomte E, Michet A. Flavour Fragrance J 1991;6:178.(g)Bertozzi S, Campigli N, Vitulli G, Lazzaroni R, Salvadori PJ. Organomet Chem 1995;487:41.(h)Horiuchi T, Ohta T, Nozaki K, Takaya H. Chem Commun 1996:155.(i)Horiuchi T, Ohta T, Shirakawa E, Nozaki K, Takaya H. Tetrahedron 1997;53:7795.(j)Barros HJV, Hanson BE, dos Santos EN, Gusevskaya EV. Appl Catal, A 2004;278:57.(k)Barros HJV, da Silva JG, Guimarães CC, dos Santos EN, Gusevskaya EV. Organometallics 2008;27:4523.
- 3. For selected reviews on C-C bond forming hydrogenation and transfer hydrogenation, see: (a)Ngai MY, Kong JR, Krische MJ. J Org Chem 2007;72:1063. [PubMed: 17288361](b)Skucas E, Ngai MY, Komanduri V, Krische MJ. Acc Chem Res 2007;40:1394. [PubMed: 17784728](c)Bower JF, Kim IS, Patman RL, Krische MJ. Angew Chem Int Ed 2009;48:34.

4. For ruthenium catalyzed C-C bond forming transfer hydrogenation, see: (a) Dienes: Shibahara F, Bower JF, Krische MJ. J Am Chem Soc 2008;130:6338. [PubMed: 18444617](b)Shibahara F, Bower JF, Krische MJ. J Am Chem Soc 2008;130:14120. [PubMed: 18841895](c) Allenes: Ngai M-Y, Skucas E, Krische MJ. Org Lett 2008;10:2705. [PubMed: 18533665](d)Skucas E, Zbieg JR, Krische MJ. J Am Chem Soc 2009;131:5054. [PubMed: 19317402](e) Alkynes: Patman RL, Chaulagain MR, Williams VM, Krische MJ. J Am Chem Soc 2009;131:2066. [PubMed: 19173651](f)Williams VM, Leung JC, Patman RL, Krische MJ. Tetrahedron 2009;65:5024.(g) Enynes: Patman RL, Williams VM, Bower JF, Krische MJ. Angew Chem Int Ed 2008;47:5220.

- 5. For related catalytic C-C couplings that occur by way of nucleophilic ruthenium π-allyls, see: (a)Tsuji Y, Mukai T, Kondo T, Watanabe Y. J Organomet Chem 1989;369:C51.(b)Kondo T, Ono H, Satake N, Mitsudo T-a, Watanabe Y. Organometallics 1995;14:1945.(c)Kondo T, Hiraishi N, Morisaki Y, Wada K, Watanabe Y, Mitsudo T-a. Organometallics 1998;17:2131.(d)Yu CM, Lee S, Hong YT, Yoon SK. Tetrahedron Lett 2004;45:6557.(e)Omura S, Fukuyama T, Horiguchi J, Murakami Y, Ryu I. J Am Chem Soc 2008;130:14094. [PubMed: 18841894]
- 6. For selected reviews of ruthenium catalyzed C-C coupling, see: (a)Trost BM, Toste FD, Pinkerton AB. Chem Rev 2001;101:2067. [PubMed: 11710241](b)Kondo T, Mitsudo T-a. Curr Org Chem 2002;6:1163.(c)Dérien S, Monnier F, Dixneuf PH. Top Organomet Chem 2004;11:1.
- 7. For catalytic intermolecular dienealdehyde reductive coupling, see: (a)Kimura M, Ezoe A, Shibata K, Tamaru Y. J Am Chem Soc 1998;120:4033.(b)Takimoto M, Hiraga Y, Sato Y, Mori M. Tetrahedron Lett 1998;39:4543.(c)Kimura M, Fujimatsu H, Ezoe A, Shibata K, Shimizu M, Matsumoto S, Tamaru Y. Angew Chem Int Ed 1999;38:397.(d)Kimura M, Shibata K, Koudahashi Y, Tamaru Y. Tetrahedron Lett 2000;41:6789.(e)Kimura M, Ezoe A, Tanaka S, Tamaru Y. Angew Chem Int Ed 2001;40:3600. (f)Loh TP, Song HY, Zhou Y. Org Lett 2002;4:2715. [PubMed: 12153217](g)Sato Y, Sawaki R, Saito N, Mori M. J Org Chem 2002;67:656. [PubMed: 11856003](h)Jang HY, Huddleston RR, Krische MJ. Angew Chem Int Ed 2003;42:4074.(i)Bareille L, Le Gendre P, Moïse C. Chem Commun 2005:775. (j)Kimura M, Ezoe A, Mori M, Iwata K, Tamaru Y. J Am Chem Soc 2006;128:8559. [PubMed: 16802822](k)Yang Y, Zhu SF, Duan HF, Zhou CY, Wang LX, Zhou QL. J Am Chem Soc 2007;129:2248. [PubMed: 17269780](l)Sato Y, Hinata Y, Seki R, Oonishi Y, Saito N. Org Lett 2007;9:5597. [PubMed: 18020355]
- 8. For a recent review encompassing nickel-catalyzed dienealdehyde reductive coupling, see: Kimuara M, Tamaru Y. Top Curr Chem 2007;279:173.
- 9. For isomerization of ruthenium  $\pi$ -allyls, see: Xue P, Bi S, Sung HHY, Williams ID, Lin Z, Jia G. Organometallics 2004;23:4735. and reference 5e.
- 10. Dobson A, Robinson SR, Uttley MF. J Chem Soc, Dalton Trans 1974:370.
- 11. Optically enriched 3g was previously prepared in 10 steps via enzymatic resolution: Fadel A, Vandromme L. Tetrahedron: Asymmetry 1999;10:1153.. Using (R)-CatASium T2 as ligand instead of dppb, optically enriched 3g (57% ee) is accessible in only 2 steps.
- 12. For a review of ruthenium catalyzed alkene hydroformylation, see: Kalck P, Peres Y, Jenck J. Adv Organomet Chem 1991;32:121.

**Scheme 1.** Plausible catalytic mechanism accounting for the results of isotopic labelling.

### Table 1

Ruthenium catalyzed reductive coupling of 2-substituted dienes **1a-1i** to paraformaldehyde via transfer hydrogenation.<sup>a</sup>

(CH<sub>2</sub>O)<sub>n</sub> (300 mol%) **1a-1i** (100 mol%) RuH<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>3</sub> (5 HO<sub>2</sub>CC<sub>7</sub>F<sub>15</sub> (5 mo DPPB (5 mol%)

*i*-PrOH-Me<sub>2</sub>CO (1M, or *i*-PrOH (1M) 80-90 °C, 20 h

1a, R = (CH<sub>2</sub>)<sub>2</sub>CH=CMe<sub>2</sub>

1d, R = CHMeOTIPS


1g, R = p-MeOPH

1b, R = cyclohexyl

1e,  $R = CH_2N(Bn)Ts$ 

1h, R = m-MeOPH

62% Yield, **3b** 1:≥20, **2b:3b**, 90 °C<sup>C</sup>



(CH<sub>2</sub>O)<sub>n</sub> (300 mol%) **1a-1i** (100 mol%)

RuH<sub>2</sub>(CO)(PPh<sub>3</sub>)<sub>3</sub> (5 HO<sub>2</sub>CC<sub>7</sub>F<sub>15</sub> (5 mo DPPB (5 mol%)

*i*-PrOH-Me<sub>2</sub>CO (1M, or *i*-PrOH (1M) 80-90 °C, 20 h

 $1a, R = (CH_2)_2CH=CMe_2$ 

1d, R = CHMeOTIPS

1g, R = p-MeOPH

1b, R = cyclohexyl1e,  $R = CH_2N(Bn)Ts$ 

1h, R = m-MeOPH

TIPSO Me OH Me 
$$64\%$$
 Yield,  $3\mathbf{d}$ ,  $1:1$  dr  $1:\geq 20$ ,  $2\mathbf{d}:3\mathbf{d}$ ,  $80$  °C  $^{b,d}$ 

MeO OH 72% Yield, 
$$3g$$
 1:220,  $2g:3g$ , 80 °C $^b$ 

MeO MeO OH 74% Yeild, 
$$3\mathbf{h}$$
  $1:\geq 20, \mathbf{2h}: \mathbf{3h}, 80 \, ^{\circ}\mathbf{C}^b$ 

 $<sup>^{</sup>a}\mathrm{In}$  all cases, cited yields are of isolated material and represent the average of two runs.

 $<sup>^</sup>b{\rm Isopropanol\text{-}Me2CO}\,(1~{\rm M},9:\!1)$  was used as solvent.

<sup>&</sup>lt;sup>c</sup>Isopropanol (1 M) was used as solvent.

 $d_{\mbox{\footnotesize Reaction time}}$  was extended to 40 h. See Supporting Information for detailed experimental procedures.

$(\mathrm{CD_2O})_\mathrm{n}, i\text{-PrOH}$	$(\mathrm{CH_2O})_\mathrm{n}, d_8$ -i-PrOH	$(\mathrm{CD_2O})_\mathrm{n}, d_{8}$ -i-PrOH
$H_a$ (5% $^2$ H)	$H_a$ (51.5% $^2$ H)	$H_a (17\%^2 H)$
$H_b (100\%^2 H)$	$H_b$ (0% $^2$ H)	$H_{\rm b}  (100\%^{ 2}{\rm H})$
$H_c (16.5\% ^2H)$	$H_c$ (46% $^2$ H)	$H_{c}$ (76.5% $^{2}H$ )
$H_d (12\% ^2H)$	$H_d$ (51% $^2H$ )	$H_d$ (58.5% $^2H$ )
$H_e$ (14% $^2$ H)	$H_e$ (50% $^2$ H)	$H_e (57.5\%^2 H)$

 $<sup>^</sup>a$ The extent of  $^2$ H incorporation was determined using  $^1$ H and  $^2$ H NMR. The indicated values represent the average of two runs.