

Amine-Catalyzed Coupling of Allenic Esters to α,β -Unsaturated Carbonyls


Catherine A. Evans and Scott J. Miller*

Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467-3860

Received August 6, 2003; E-mail: scott.miller.1@bc.edu

Organic reactions that lead to increases in molecular complexity are of great value in both target-oriented and diversity-oriented synthesis. In this context, the phosphine-catalyzed cycloaddition of allenic esters to enones represents a process that has received substantial attention in recent years (Scheme 1).¹ Thus, allene **1** and enone **2** undergo a reaction to give cycloadduct **3** in the presence of substoichiometric quantities of triphenyl phosphine. From a mechanistic point of view, the reaction is often represented as proceeding through a zwitterionic intermediate that serves as a dipole for a [3 + 2]-cycloaddition.² In the process of investigating whether nitrogen-based nucleophiles could induce similar reactivity, we found that a very different path is followed. Rather than a P-catalyzed cycloaddition, an N-catalyzed conjugate addition is observed. In this case, building blocks **1** and **2**, upon exposure to a catalytic quantity (10 mol %) of quinuclidine, afford product **4** in excellent yield (87%).

Scheme 1



The reaction proceeds under very mild conditions (10 mol % quinuclidine, PhCH_3 , 20 °C), and is efficient for a range of α,β -unsaturated carbonyl compounds.³ Table 1 presents the substrate scope we have charted to date. Tetralone derivative **2** participates in the reaction, affording adduct **4** in 87% yield within 24 h (entry 1). Indanone analogue **5** participates in a faster reaction, delivering **6** in 81% isolated yield after 12 h (entry 2). Heterocyclic versions also participate; compound **7** undergoes reaction to give allenic ester **8** in 78% yield (entry 3); substituted enone **9** delivers a comparable yield of **10** (78%, entry 4) as a 1.8:1 mixture of diastereomers (*trans/cis*). Benzosuberone analogue **11** participated in a slower reaction, delivering **12** in 65% yield after 36 h (entry 5). Likewise, *p*-methoxy-substituted tetralone **13** delivers **14** in 50% yield after 48 h (entry 6). Acyclic enones such as ethyl vinyl ketone (**15**) deliver the linear substituted allenolate **16** in 70% yield. Imide **17** is an excellent substrate for the process, affording compound **18** in 98% isolated yield (entry 8). Ester **19** is also a good substrate for the reaction as product **20** is obtained in 85% yield (entry 9). Of note is that a number of substrates have been found to be unreactive under the reaction conditions. For example, while acyclic enone **15** reacts efficiently under the reaction conditions, enone **21** is unreactive (entry 10). Despite prolonged reaction times, and the subjection of the reagents to elevated temperatures, product **22** was not observed under the conditions that were explored.


Baylis–Hillman adducts such as **23** are also converted to product **24** under the standard reaction conditions (eq 1). Yet, MOM-ether

Table 1. Substrate Scope for Coupling of Ethyl Allenolate to α,β -Unsaturated Carbonyl Compounds^a

Entry	Substrate	Product	Time	Yield ^b
1			24 h	87%
2			12 h	81%
3			20 h	78%
4			20 h	78% (1.8:1, <i>trans/cis</i>)
5			36 h	65%
6			48 h	50%
7			24 h	70%
8			24 h	98%
9			36 h	85%
10			No Reaction	No Reaction

^a All reactions were conducted at room temperature in PhCH_3 in the presence of quinuclidine (10 mol %). ^b Yields refer to isolated yield after silica gel chromatography.

25 does not produce the corresponding allenolate-coupled product **26** under the reaction conditions.⁴



The fact that Baylis–Hillman adducts participate in this quinuclidine-catalyzed reaction prompted us to explore whether a Baylis–Hillman reaction might be coupled to the allenolate conjugate addition since both processes are in fact now known to be catalyzed by nucleophilic amines.⁵ Such a process would constitute a three-




Table 2. Substrate Scope for Three-Component Coupling of Ethyl Allenolate, Aldehydes and Acrylate **19**^a

Entry	Aldehyde	Product	Time	Yield ^b
1	$R_1 = \text{PhCH}_2\text{CH}_2$	27	18 h	88 %
2	$R_2 = \text{MeCH}_2\text{CH}_2$	28	18 h	80 %
3	$R_3 = i\text{-Bu}$	29	18 h	60 %
4	$R_4 = \text{Et}$	30	18 h	65 %


^a All reactions were conducted at room temperature in CH_3CN in the presence of quinuclidine (20 mol %). ^b Yields refer to isolated yield after silica gel chromatography.

component coupling (TCC)⁶ between an α,β -unsaturated carbonyl compound (e.g., acrylate **19**), an aldehyde, and the allenolate **1**. Indeed, when **19** (1.0 equiv) is subjected to a Baylis–Hillman reaction (4.0 equiv of aldehyde)⁷ and allenolate **1** (4.0 equiv) is introduced into the same flask after an appropriate interval (12 h), TCC products may be isolated in good yield (eq 2, Table 2).⁸ For example, with hydrocinnamaldehyde, product **27** is obtained in 88% yield (entry 1). Saturated aldehydes represent good substrates for the reaction, with *n*-butyraldehyde affording product **28** in 80% yield (entry 2); 3-methyl-butyraldehyde delivers **29** in 60% yield (entry 3). Propionaldehyde also affords comparable reaction efficiency, wherein allene **30** is obtained in 65% isolated yield. In each case, products **27–30** are obtained as predominantly single diastereomers (>20:1 trans/cis by ¹H NMR spectroscopic analysis).^{4,9}

The divergent reactivity exhibited by the starting materials in the presense of different nucleophiles represents an example of conditionally dependent reactivity (Scheme 2). In the case of the phosphine-catalyzed reaction, the catalyst may react with the allene to generate zwitterionic enolate **31**. This species may then undergo cycloaddition with **2** to deliver spirocyclic ylide **32**. Facile 1,2-proton transfer then yields **3**, and regeneration of catalyst. This mechanism, as proposed by others,¹⁰ benefits from the ability of phosphorus to support the ylide-like structure **32**. In contrast, the amine-catalyzed pathway does not benefit from analogous stabilization. Rather, zwitterionic enolate **33** may undergo 1,4-addition to **2** to deliver **34**. This newly generated enolate does not undergo a second C–C bond-forming step to form the spirocycle because the ammonium ion-substituted adduct cannot similarly stabilize the ylide. Thus, proton transfer and tautomerization ensues, yielding adduct **4** with concomitant regeneration of the amine catalyst. While this mechanistic hypothesis remains to be verified, it does account for the conditionally dependent reactivity of **1** and **2** in the presence of different nucleophilic catalysts.

In conclusion, we have presented an efficient, amine-catalyzed coupling of allenolate esters and α,β -unsaturated carbonyl com-

Scheme 2



pounds. The scope of the reaction has been defined under a very mild set of conditions. Significantly, the implications of the discovery have been extended to include a three-component coupling reaction where two distinct catalytic C–C bond-forming events unite three building blocks under the influence of a unique catalyst.

Acknowledgment. This research is supported by the National Science Foundation (CHE-0236591). In addition, we are grateful to the Merck Chemistry Council and Pfizer Global Research for support. S.J.M. is a Fellow of the Alfred P. Sloan Foundation and a Camille Dreyfus Teacher-Scholar.

Supporting Information Available: Experimental procedures and product characterization for all new compounds synthesized (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (a) Zhang, C.; Lu, X. *J. Org. Chem.* **1995**, *60*, 2906. (b) Zhu, G.; Chen, Z.; Jiang, Q.; Xiao, D.; Cao, P.; Zhang, X. *J. Am. Chem. Soc.* **1997**, *119*, 3836. (c) Xu, Z.; Lu, X. *Tetrahedron Lett.* **1999**, *40*, 549. (d) Du, Y.; Lu, X.; Yu, Y. *J. Org. Chem.* **2002**, *67*, 8901.
- For several key references to nucleophilic catalysis via conjugate addition of P-nucleophiles, see: (a) Trost, B. M.; Kazmaier, U. *J. Am. Chem. Soc.* **1992**, *114*, 7933. (b) Guo, C.; Lu, X. *J. Chem. Soc., Perkin Trans. 1* **1993**, 1921. (c) Trost, B. M.; Dake, G. R. *J. Am. Chem. Soc.* **1997**, *119*, 7595. (d) Trost, B. M.; Dake, G. R. *J. Org. Chem.* **1997**, *62*, 5670. (e) Wang, L.-C.; Luiz, A.-L.; Agapiou, K.; Jang, H.-Y.; Krische, M. J. *J. Am. Chem. Soc.* **2002**, *124*, 2402. (f) Frank, S. A.; Mergott, D. J.; Roush, W. R. *J. Am. Chem. Soc.* **2002**, *124*, 2404. (g) Wang, J.-C.; Ng, S.-S.; Krische, M. J. *J. Am. Chem. Soc.* **2003**, *125*, 3682.
- DABCO is also an effective catalyst for the reaction.
- The stereochemical assignment is in the Supporting Information.
- (a) Basavaiah, D.; Rao, A. J.; Satyanarayana, T. *Chem. Rev.* **2003**, *103*, 811. (b) Aggarwal, V. K.; Emme, I.; Fulford, S. Y. *J. Org. Chem.* **2003**, *68*, 692.
- For some recent examples of catalytic multicomponent coupling reactions, see: (a) Montgomery, J. *Acc. Chem. Res.* **2000**, *33*, 467. (b) Wender, P. A.; Gamber, G. G.; Hubbard, R. D.; Zhang, L. *J. Am. Chem. Soc.* **2002**, *124*, 2876. (c) Patel, J. S.; Jamison, T. F. *Angew. Chem., Int. Ed.* **2003**, *42*, 1364. (d) Jeganmohan, M.; Shanmugasundaram, M.; Cheng, C.-H. *Org. Lett.* **2003**, *5*, 881. (e) Bossharth, E.; Desbordes, P.; Monteiro, N.; Balme, G. *Org. Lett.* **2003**, *5*, 2441.
- Lee, W. D.; Yang, K.-S.; Chen, K. *Chem. Commun.* **2001**, 1612.
- A byproduct of the liberated naphthol involves conjugate addition to scavenge the excess allenolate; this product is readily removed by chromatography.
- For previous conjugate additions of 5-methylene-1,3-dioxan-4-ones, see: (a) Bulliard, M.; Zehnder, M.; Giese, B. *Helv. Chim. Acta* **1991**, *74*, 1600. (b) Piber, M.; Leahy, J. W. *Tetrahedron Lett.* **1998**, *39*, 2043.
- Lu, X.; Zhang, C.; Xu, Z. *Acc. Chem. Res.* **2001**, *34*, 535.

JA0377798