

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/6541688>

Giant Heterometallic Cu₁₇ Mn₂₈ Cluster with T_d Symmetry and High-Spin Ground State

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · MARCH 2007

Impact Factor: 12.11 · DOI: 10.1021/ja065707l · Source: PubMed

CITATIONS

128

READS

71

8 AUTHORS, INCLUDING:



Wei-Xiong Zhang

Sun Yat-Sen University

83 PUBLICATIONS 3,792 CITATIONS

SEE PROFILE



Ming-Liang Tong

Sun Yat-Sen University

280 PUBLICATIONS 12,143 CITATIONS

SEE PROFILE



Motohiro Nakano

Osaka University

174 PUBLICATIONS 4,369 CITATIONS

SEE PROFILE



David Hendrickson

University of California, San Diego

599 PUBLICATIONS 25,889 CITATIONS

SEE PROFILE

Giant Heterometallic $\text{Cu}_{17}\text{Mn}_{28}$ Cluster with T_d Symmetry and High-Spin Ground StateWen-Guo Wang,[†] Ai-Ju Zhou,[†] Wei-Xiong Zhang,[†] Ming-Liang Tong,^{*,†} Xiao-Ming Chen,[†] Motohiro Nakano,[§] Christopher C. Beedle,[‡] and David N. Hendrickson^{*,‡}

MOE Key Laboratory of Bioinorganic and Synthetic Chemistry, School of Chemistry and Chemical Engineering, Sun Yat-Sen University, Guangzhou 510275, People's Republic of China, Department of Applied Chemistry, Osaka University, Suita 565-0871, Osaka, Japan, and Department of Chemistry and Biochemistry-0358, University of California at San Diego, La Jolla, California 92093-0358

Received August 7, 2006; E-mail: tongml@mail.sysu.edu.cn; dhendrickson@ucsd.edu

Since the 1990s, manganese clusters with large spin ground states have received increasing attention mainly due to their single-molecule magnet behavior.^{1,2} A number of high nuclearity homometallic manganese complexes from Mn_{16} to Mn_{84} have been explored to date.^{3–11} However, the rational synthesis of high nuclear heterometallic manganese clusters remains a challenge. Triethanolamine (teaH_3) is a potentially versatile ligand, and several polynuclear complexes have been recently reported.¹² Enlightened by the facile reactions of zerovalent copper with $\text{Co}^{\text{II}}/\text{Zn}^{\text{II}}/\text{Cd}^{\text{II}}/\text{Pd}^{\text{II}}$ salts and aminoalcohols,¹³ we tried to extend such reactions into a new synthetic route to heterometallic manganese clusters. We report herein the synthesis, structure, and magnetic properties of a giant heterometallic $\text{Cu}_{14}\text{Cu}^{\text{II}}_{13}\text{Mn}^{\text{II}}_{12}\text{Mn}^{\text{III}}_{12}\text{Mn}^{\text{IV}}_{12}$ cluster, $[\text{Cu}_{17}\text{Mn}_{28}\text{O}_{40}(\text{tea})_{12}(\text{HCO}_2)_6(\text{H}_2\text{O})_4] \cdot 36\text{H}_2\text{O}$ (**1**). To our knowledge, **1** is the largest manganese teaH_3 cluster and unprecedented with five metal oxidation states.

The reaction of copper powder, $\text{Mn}(\text{OAc})_2 \cdot 4\text{H}_2\text{O}$, and teaH_3 in a 1:2:1 ratio in *N,N'*-dimethylformamide (DMF) (30 mL) at 85 °C gave a dark green solution from which dark block crystals of **1** were slowly deposited after several months. The in situ generated formate ligand arises from hydrolysis of DMF which has been previously documented.¹⁴

The crystal structure of **1**¹⁵ contains a giant $[\text{Cu}_{17}\text{Mn}_{28}\text{O}_{40}]^{42+}$ core of T_d symmetry with 4 Cu^{I} ions, 13 Cu^{II} ions, 4 Mn^{II} ions, 12 Mn^{III} ions, 12 Mn^{IV} ions, 28 $\mu_4\text{-O}^{2-}$, and 12 $\mu_3\text{-O}^{2-}$ ions. Each $\mu_4\text{-O}^{2-}$ ion joins three Mn ions and one Cu ion ($2\text{Mn}^{\text{III}} + \text{Mn}^{\text{II}} + \text{Cu}^{\text{II}}$ or $2\text{Mn}^{\text{IV}} + \text{Mn}^{\text{III}} + \text{Cu}^{\text{I}}$ or $3\text{Mn}^{\text{IV}} + \text{Cu}^{\text{II}}$), while each $\mu_3\text{-O}^{2-}$ ion joins one Mn^{IV} ion and two Mn^{III} ions. There are five metal oxidation states of Mn^{2+} , Mn^{3+} , Mn^{4+} , Cu^+ , and Cu^{2+} in **1**, which were established by consideration of bond lengths, bond valence sum (BVS) calculations,¹⁶ charge considerations, and the observation of Jahn–Teller (JT) distortions for the Mn^{III} centers. The $[\text{Cu}_{17}\text{Mn}_{28}\text{O}_{40}]^{42+}$ core is surrounded by 12 tea^{3-} groups, 6 HCO_2^- groups, and 4 aqua ligands into a neutral cluster (Figure 1).

Each Cu^{I} ion is coordinated in a tetrahedral geometry by three $\mu_4\text{-O}^{2-}$ ions and one aqua ligand. The Cu^{II} ions have two different coordination environments: the central Cu^{I} ion is coordinated in a perfect tetrahedral environment by 4 $\mu_4\text{-O}^{2-}$ ions ($\text{Cu}–\text{O} = 1.975(11)$ Å), while each of the outer 12 symmetry-related Cu^{II} ions is coordinated in a distorted trigonal bipyramidal environment by three alkoxo O atoms in the plane ($\text{O}–\text{Cu}^{\text{II}}–\text{O} = 102.9(3)–127.58(17)^\circ$) and one amino N atom and one $\mu_4\text{-O}^{2-}$ ion at the axial positions ($\text{N1}–\text{Cu}^{\text{II}}–\text{O4} = 172.6(3)^\circ$). Each tea^{3-} ligand acts in a $\mu_4:\eta^1:\eta^2:\eta^2:\eta^2$ mode to cap an outer Cu^{II} atom and bridge one Mn^{II} and two




Figure 1. Molecular structure (a) and coordination environments (b) of **1**. H atoms and lattice water molecules have been omitted for clarity. Symmetry: (a) $-y, x, -z$; (b) $y, -x, -z$; (c) $-x, -y, z$; (d) $-y, -x, z$; (e) $-z, -x, y$; (f) x, z, y ; (g) $-x, y, -z$; (h) y, z, x ; (i) z, x, y ; (j) y, x, z ; (k) $-y, z, -x$.




Figure 2. The adamantane-like Mn_{28} cluster containing six $[\text{Mn}^{\text{IV}}_2\text{Mn}^{\text{III}}_2\text{O}_4]$ and four $[\text{Mn}^{\text{IV}}_3\text{Mn}^{\text{II}}\text{O}_4]$ cubanes via sharing all Mn^{IV} atoms. Atom color code: Mn^{IV} , dark yellow; Mn^{III} , navy blue; Mn^{II} , magenta; O, red.

Mn^{III} ions. There are three oxidation states for the manganese ions, which is uncommon for the manganese clusters.^{8,10,17} All Mn centers are six-coordinate (Figure S1), through interactions with five $\mu_4\text{-O}^{2-}$ ions and one $\mu_3\text{-O}^{2-}$ ion for Mn1 in an essentially octahedral geometry (average $\text{Mn1}–\text{O} = 1.904$ Å), one $\mu_4\text{-O}^{2-}$ ion, two $\mu_3\text{-O}^{2-}$ ions, two formate O atoms, and one alkoxide O atom for Mn2 in a tetragonally elongated octahedral geometry ($\text{Mn2}–\text{O2}$ and $\text{Mn2}–\text{O7} = 2.221(6)$ and $2.143(8)$ Å constituting the JT axis), and three $\mu_4\text{-O}^{2-}$ ions and three alkoxide O atoms for Mn3 forming a distorted trigonal prism (average $\text{Mn3}–\text{O} = 2.190$ Å), respectively.

The most striking structural feature is that all the 28 Mn ions are bridged by the 28 $\mu_4\text{-O}^{2-}$ and 12 $\mu_3\text{-O}^{2-}$ ions into 6 $\text{Mn}^{\text{III}}_2\text{Mn}^{\text{IV}}_2\text{O}_4$ cubanes ($\text{Mn}–\text{O}–\text{Mn} = 93.6(2)–102.5(2)^\circ$) and 4 $\text{Mn}^{\text{II}}\text{Mn}^{\text{IV}}_3\text{O}_4$ cubanes ($\text{Mn}^{\text{II}}–\text{O}–\text{Mn}^{\text{IV}}$, $\text{Mn}^{\text{IV}}–\text{O}–\text{Mn}^{\text{IV}} = 97.5(3)–99.9(3)^\circ$), which are further joined to be a cubane-based manganese cage via sharing all the 12 Mn^{IV} ions (Figure 2). Such a structure is unprecedented in the known high nuclearity Mn clusters. Finally, the connectivity of the $[\text{Cu}_{17}\text{Mn}_{28}\text{O}_{40}]^{42+}$ core can be precisely

[†] Sun Yat-Sen University.[§] Osaka University.[‡] University of California at San Diego.




Figure 3. Plot of reduced magnetization ($M/N\beta$) versus H/T for **1** in the temperature range of 1.8–4.0 K (5 kG (orange), 10 kG (dark green), 20 kG (brown), 30 kG (dark red), 40 kG (teal), 50 kG (dark blue), 60 kG (green), 70 kG (red)). Solid lines represent best theoretical fit. Inset illustrates $\chi_m T$ versus T plot for complex **1** from 1.8 to 300 K (1 kG (black), 10 kG (red), 50 kG (blue)).




Figure 4. AC magnetic measurements for **1** at $H_{ac} = 5$ G and $H_{dc} = 0$.

described by noting that the central Cu^{I} ion connects four $\text{Mn}^{\text{II}}\text{Mn}^{\text{IV}}_3\text{O}_4$ cubanes into a tetrahedral array. Six $\text{Mn}^{\text{III}}_2\text{Mn}^{\text{IV}}_2\text{O}_4$ cubanes are located on the midlines of the six edges of the tetrahedron and connect the four $\text{Mn}^{\text{II}}\text{Mn}^{\text{IV}}_3\text{O}_4$ cubanes via the 12 Mn^{IV} corners into an adamantane-like cage. Each of the four $[\text{Cu}^{\text{I}}(\text{H}_2\text{O})]$ groups (on 3-fold axes) connects three $\text{Mn}^{\text{III}}_2\text{Mn}^{\text{IV}}_2\text{O}_4$ cubanes, and the outer 12 Cu^{II} ions connect the Mn_{28} cluster via the remaining $\mu_4\text{-O}^{2-}$ sites.

DC magnetic susceptibility data were collected in fields ranging from 70 to 1 kG in the temperature range of 1.8–300 K (Figure 3). The reduced magnetization (M) versus H/T data cannot be fit to one state experiencing axial zero-field splitting $D\hat{S}_z^2$ (in fact, T_d symmetry does not allow this). The χT versus T data (inset Figure 3) taken at fields of less than 1 kG show a maximum of $\sim 325 \text{ cm}^3 \text{ mol}^{-1} \text{ K}$. This suggests a $S = 5/2$ ground state in small fields, which is reasonable given that the $\text{Mn}^{\text{III}}\cdots\text{Mn}^{\text{III}}$, $\text{Mn}^{\text{IV}}\cdots\text{Mn}^{\text{IV}}$, and $\text{Mn}^{\text{III}}\cdots\text{Mn}^{\text{IV}}$ interactions within the cubane units are expected to be ferromagnetically coupled, while the $\text{Mn}^{\text{II}}\cdots\text{Mn}^{\text{IV}}$, $\text{Cu}^{\text{II}}\cdots\text{Mn}^{\text{III}}$, and $\text{Cu}^{\text{II}}\cdots\text{Mn}^{\text{IV}}$ interactions are antiferromagnetic which would give $S = 12S_{\text{Mn(IV)}} + 12S_{\text{Mn(III)}} - 4S_{\text{Mn(II)}} - 13S_{\text{Cu(II)}} = 5/2$. The non-superimposability of isofields in the reduced magnetization versus H/T plot usually indicates the presence of axial zero-field interactions. However, we have fit all the data (solid lines in Figure 3) assuming that there is a $S = 63/2$ excited state lying 5 K above the $S = 5/2$ ground state. The fit is quite good, and it is important to emphasize that the model has no zero-field splitting for either of the states. This is consistent with the T_d symmetry that precludes the $D\hat{S}_z^2$ term in the spin Hamiltonian. High-frequency EPR studies have been initiated to verify the two spin state model.

AC magnetic susceptibility data for **1** (Figure 4) do show a frequency-dependent out-of-phase signal; however, only the onset

of the signal is seen. Lower temperature AC data or micro-SQUID data are needed to establish whether this interesting complex **1** functions as a single-molecule magnet.

In summary, we have established an approach to generate a giant heterometallic mixed-valent $\text{Cu}^{\text{I}}/\text{Cu}^{\text{II}}/\text{Mn}^{\text{II}}/\text{Mn}^{\text{III}}/\text{Mn}^{\text{IV}}$ cluster which contains an unprecedented adamantane-like Mn_{28} core connected wholly by Mn_4O_4 cubane units and exhibits unusual magnetic properties.

Acknowledgment. This work was supported by the NSFC (Nos. 20525102 & 20471069), the FANEDD of China (200122), the Scientific and Technological Project of Guangdong Province (04205405), and the NSF (U.S.A.).

Supporting Information Available: Synthesis details, additional plots, and an X-ray crystallographic file in CIF format for the structural determination of **1**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Gatteschi, D.; Sessoli, R. *Angew. Chem., Int. Ed.* **2003**, *42*, 268–297. (b) Aromí, G.; Brechin, E. K. *Struct. Bonding* **2006**, *122*, 1–67.
- (2) Sessoli, R.; Tsai, H.-L.; Schake, A. R.; Wang, S.; Vincent, J. B.; Folting, K.; Gatteschi, D.; Christou, G.; Hendrickson, D. N. *J. Am. Chem. Soc.* **1993**, *115*, 1804–1816.
- (3) Price, J. P.; Batten, S. R.; Moubaraki, B.; Murray, K. S. *Chem. Commun.* **2002**, 762–763.
- (4) Brechin, E. K.; Sanudo, E. C.; Wernsdorfer, W.; Boskovic, C.; Yoo, J.; Hendrickson, D. N.; Yamaguchi, A.; Ishimoto, H.; Concolino, T. E.; Rheingold, X. A. L.; Christou, G. *Inorg. Chem.* **2005**, *44*, 502–511.
- (5) Dendrinou-Samara, C.; Alexiou, M.; Zaleski, C. M.; Kampf, J. W.; Krik, M. L.; Kessissoglou, D. P.; Pecoraro, V. L. *Angew. Chem., Int. Ed.* **2003**, *42*, 3763–3766.
- (6) Brockman, J. T.; Huffman, J. C.; Christou, G. *Angew. Chem., Int. Ed.* **2002**, *41*, 2506–2508.
- (7) Murugesu, M.; Raftery, J.; Wernsdorfer, W.; Christou, G.; Brechin, E. K. *Inorg. Chem.* **2004**, *43*, 4203–4209.
- (8) Murugesu, M.; Habrych, M.; Wernsdorfer, W.; Abboud, K. A.; Christou, G. *J. Am. Chem. Soc.* **2004**, *126*, 4766–4767.
- (9) (a) Jones, L. F.; Brechin, E. K.; Collison, D.; Harrison, A.; Teat, S. J.; Wernsdorfer, W. *Chem. Commun.* **2002**, 2974–2975. (b) Jones, L. F.; Raftery, J.; Teat, S. J.; Collison, D.; Brechin, E. K. *Polyhedron* **2005**, *24*, 2443–2449.
- (10) Soler, M.; Wernsdorfer, W.; Folting, K.; Pink, M.; Christou, G. *J. Am. Chem. Soc.* **2004**, *126*, 2156–2165.
- (11) Tasiopoulos, A. J.; Vinslava, A.; Wernsdorfer, W.; Abboud, K. A.; Christou, G. *Angew. Chem., Int. Ed.* **2004**, *43*, 2117–2121.
- (12) (a) Khan, M. I.; Tabussum, S.; Doedens, R. J. *Chem. Commun.* **2003**, 532. (b) Wittick, L. M.; Murray, K. S.; Moubaraki, B.; Batten, S. R.; Spiccia, L.; Berry, K. J. *Dalton Trans.* **2004**, 1003–1011. (c) Murugesu, M.; Wernsdorfer, W.; Abboud, K. A.; Christou, G. *Angew. Chem., Int. Ed.* **2005**, *44*, 892–896. (d) Rajaraman, G.; Murugesu, M.; Sanudo, E. C.; Soler, M.; Wernsdorfer, W.; Helliwell, M.; Muryn, C.; Raftery, J.; Teat, S. J.; Christou, G.; Brechin, E. K. *J. Am. Chem. Soc.* **2004**, *126*, 15445–15457. (e) Wittick, L. M.; Jones, L. F.; Jensen, P.; Moubaraki, B.; Spiccia, L.; Berry, K. L.; Murray, K. S. *Dalton Trans.* **2006**, 1534–1543.
- (13) (a) Makhankova, V. G.; Vassilyeva, O. Y.; Kokozyay, V. N.; Skelton, B. W.; Reedijk, J.; Van Albada, G. A.; Sorace, S.; Gatteschi, D. *New J. Chem.* **2001**, *25*, 685–689. (b) Makhankova, V. G.; Vassilyeva, O. Y.; Kokozyay, V. N.; Reedijk, J.; Van Albada, G. A.; Jezierska, J.; Skelton, B. W. *Eur. J. Inorg. Chem.* **2002**, 2163–2169.
- (14) (a) Perrin, D. D.; Armarego, W. L. F. *Purification of Laboratory Chemicals*, 3rd ed.; Pergamon: Oxford, 1988. (b) Murthy, N. N.; Mahroof-Tahir, M.; Karlin, K. D. *J. Am. Chem. Soc.* **1993**, *115*, 10404–10405.
- (15) Crystal data for **1**: $\text{C}_{78}\text{H}_{230}\text{Cu}_{17}\text{Mn}_{28}\text{N}_{12}\text{O}_{128}$, 6003.24 g mol⁻¹, cubic $I43m$, $a = 20.3441(9) \text{ \AA}$, $V = 8420.1(6) \text{ \AA}^3$, $Z = 2$, $\rho = 2.368 \text{ g cm}^{-3}$, $T = 123 \text{ K}$, $\mu = 4.229 \text{ mm}^{-1}$, $F(000) = 5998$, $2\theta_{\text{max}} = 56.0^\circ$, $S = 1.086$. The structure was solved by direct methods and refined on F^2 to $R_1(wR_2)$ 0.0456 (0.1177) using 1758 reflections with $I > 2\sigma(I)$.
- (16) (a) Bond valence sum calculations for Mn^{II} , Mn^{III} , and Mn^{IV} ions gave oxidation state values of 2.193, 3.350, and 4.413, respectively, and Cu^{I} and Cu^{II} ions gave oxidation state values of 1.284 and 1.797–1.951, respectively. (b) Liu, W.; Thorp, H. H. *Inorg. Chem.* **1993**, *32*, 4102–4105.
- (17) Sun, Z.; Gantzel, P. K.; Hendrickson, D. N. *Inorg. Chem.* **1996**, *35*, 6640–6641.

JA065707L