

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/5772736>

ChemInform Abstract: Enantioselective, Organocatalytic Oxy-Michael Addition to γ/δ -Hydroxy- α,β -enones: Boronate-Amine Complexes as Chiral Hydroxide Synthons

ARTICLE *in* JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · FEBRUARY 2008

Impact Factor: 12.11 · DOI: 10.1021/ja076802c · Source: PubMed

CITATIONS

80

READS

33

3 AUTHORS, INCLUDING:



John R Falck

University of Texas Southwestern Medical Ce...

920 PUBLICATIONS 31,746 CITATIONS

SEE PROFILE

Published in final edited form as:

J Am Chem Soc. 2008 January 9; 130(1): 46–48. doi:10.1021/ja076802c.

Enantioselective, Organocatalytic Oxy-Michael Addition to γ/δ -Hydroxy- α,β -enones: Boronate-Amine Complexes as Chiral Hydroxide Synthons


De Run Li, Andiappan Murugan, and J. R. Falck*

Departments of Biochemistry and Pharmacology, University of Texas Southwestern Medical Center, Dallas, Texas 75390 USA

Abstract

An organocatalytic, enantioselective oxy-Michael addition to achiral γ - and δ -hydroxy- α,β -enones was developed. The key transformation is an unprecedented, asymmetric conjugate addition triggered by complexation between an *in situ* generated boronic acid hemiester and a chiral amine catalyst. Functionally, the intermediate amine-boronate complex acts as a chiral hydroxide surrogate or synthon. The resultant chiral β -hydroxy-ketones are obtained in good to excellent yields and high ee following mild oxidative removal of the cyclic boronate. Natural products (*R*,12*Z*,15*Z*)-2-hydroxy-4-oxohexacos-12,15-dienyl acetate and (+)-(*S*)-Streptenol A were synthesized to demonstrate the utility of this reaction.


Structural motif **1** is present in a wide range of natural products and synthetic intermediates.¹ While Michael additions of hydroxide or synthetic equivalents to α,β -unsaturated carbonyls represent an attractive approach to this moiety,² the strong basicity of the former and generally poor nucleophilicity or lability of the latter often render this option problematic. In its place, the intramolecular oxy-Michael addition of hemiacetal/hemiketal-derived alkoxides has emerged as a popular alternative strategy,³ although the resultant cyclic acetals/ketals can be difficult to remove. In some instances, satisfactory diastereoselectivity has been attained via exploitation of adjacent secondary hydroxy or amino stereocenters.^{4,5} In 2001, Watanabe *et al.*⁶ introduced an asymmetric version of the oxy-Michael addition utilizing chiral hemiketals derived from **D**-glucose and **D**-fructose in the more challenging case of achiral γ/δ -hydroxy- α,β -enones. Herein, we reported an unprecedented organocatalytic, enantioselective oxy-Michael addition to achiral γ/δ -hydroxy- α,β -enones and its use in the preparation of **1** (eq 1).⁷ The key transformation is the asymmetric conjugate addition triggered by complexation between boronic acid hemiester **3**, generated *in situ* from γ/δ -hydroxy- α,β -enones, and a chiral amine catalyst. Functionally, the intermediate amine-boronate complex acts as a *chiral hydroxide surrogate or synthon*. Mild, oxidative removal of the boronate moiety from the dioxaborolane ($n = 0$) or dioxaborinane ($n = 1$) adduct **2** furnishes **1** in good to excellent overall yield and % ee.



Recently, this⁸ and other laboratories⁹ have highlighted the nucleophilic properties of organoboronic acids and the unique stereospecific reactions of their borate complexes. Despite

expectations, when model compound (*E*)-4-hydroxy-1-phenyl-2-buten-1-one (**4**) was mixed with equimolar phenylboronic acid and activated 4Å molecular sieves in CH₂Cl₂ (Table 1, entry 1), no intramolecular Michael addition was observed, even though the hemiester (**3**; R=Ph, n = 0) could be detected by ¹H and ¹³C NMR. Inclusion of some common bases, viz., bicarbonate (entry 2), carbonate (entry 3), and pyridine (entry 4), likewise disappointed. However, a catalytic amount of Et₃N (20 mol%, entry 5) gave rise to the desired dioxaborolane **5** from which diol **6** was secured in 42% overall yield (50% recovered **4** after 48 h) following workup with basic H₂O₂. We attribute this dramatic difference to the in situ formation of a nucleophilic, pyramidized quaternary boronate complex.^{10,11} DABCO, under otherwise identical conditions, significantly improved the overall conversion (entry 6) whereas diisopropylamine (20 mol%, entry 7) worked even better and delivered **6** in 86% overall yield in just 16 h. The reaction rate for the latter was enhanced even further in toluene (entry 8), but not in DME (entry 9); the yield showed only a modest solvent dependency. DBU (entry 10) and the highly hindered 1,2,2,6,6-pentamethylpiperidine (entry 11) offered no advantage.

Inspired by the pronounced success of push/pull-type bifunctional organocatalysts,¹² we pursued an asymmetric version of this intramolecular oxy-Michael addition (Figure 1). Coordination of the carbonyl by the thiourea (the pull) and complexation of the tertiary nitrogen to boron (the push) were expected to simultaneously enhance the nucleophilicity of the boronate oxygen as well as envelope the enone in a chiral environment.¹³ Indeed, Michael addition to **4** in CH₂Cl₂ mediated by catalyst **7**^{12b} was complete in 16 h (Table 2, entry 1), > 3 times faster than a similar Et₃N catalyzed addition. More significantly, **12** was obtained in 91% yield and 91% ee after basic H₂O₂ workup.^{14,15} A comparable reaction in toluene proceeded still faster (8 h), but with significantly reduced enantioselectivity (65% ee), whereas in DME the rate was slower (24 h) and the % ee improved modestly to 94%. The less expensive quinine-based catalyst **8**^{12b} was virtually equivalent to **7** in all respects (24 h, 95% ee). Notably, catalysts **9**^{12b} and **10**^{12a} in DME (entry 2) provided access to the opposite enantiomeric diol, **13**, in synthetically useful yields and enantioselectivities (40 h/89% ee and 48 h/91% ee, respectively). In sharp contrast to the accelerated rate seen with *i*Pr₂NH, catalyst **11**, which also contains a secondary amine, proved surprisingly sluggish and was not pursued further.




The scope of the oxy-Michael was further explored using catalyst **8** and a representative sampling of γ -hydroxy- α,β -enones (Table 2). Predictably, arylketones bearing strong electron withdrawing substituents (entry 3) reacted faster than electron rich systems (entries 4 and 5), although the enantioselectivities of the latter were better.

Aliphatic ketones (entries 6 and 7), regardless of steric congestion adjacent to the carbonyl (**20** vs. **22**), had retarded reaction rates, yet still afforded excellent overall yields. The conversion of **24** into **25** (entry 8) thus appears anomalous for its comparatively rapid rate and may reflect an unanticipated coordination by the terminal oxygen substituent. The survival of the labile silyl (TES) ether also testifies to the mildness of the reaction conditions. Importantly,

additional substitution at the olefin (entry 9) or carbinol (entry 10) was well tolerated and adds to the level of structural complexity that can be achieved.

To validate the applicability of the foregoing methodology in natural products total synthesis, acetate **33**,¹⁶ an extraordinarily potent antifungal/hepatic protective agent isolated from avocado, was prepared by a biomimetic route (Scheme 1). Addition of lithium dimethyl methylphosphonate to methyl linoleate (**30**) and condensation of the adduct with glycoaldehyde furnished enone **31** which was subjected to oxy-Michael addition catalyzed by **8**. The product (*R*)-diol **32** (90% yield, 91% ee¹⁴) was selectively acetylated to give **33**.¹⁷

With δ -hydroxy- α,β -enones, oxy-Michael addition proceeded quite slowly in all solvents, although toluene was generally the best. Increasing the catalyst loading to 20 mol% and the temperature to 50°C, however, allowed the reaction to proceed at an acceptable rate and enantioselectivity for aromatic enones (Table 3: entries 1–3). For the more recalcitrant aliphatic enones (entries 4–6), these conditions were not sufficient. We, thus, screened a panel of commercial arylboronic acids to identify 3,4,5-trimethoxyphenylboronic acid as a more efficacious nucleophilic partner, which furnished aliphatic diols in good to excellent enantioselectivities at suitable rates. Diol **44** was identical in all respects with (+)-(*S*)-streptenol A, one of four known streptenols produced by *Streptomyces luteogriseus* that has attracted attention as an immunostimulant as well as an inhibitor of cholesterol biosynthesis and tumor cells.¹⁹



In contrast to carboxylic acids, boronic acids and their chiral complexes have not been well explored as nucleophilic reagents in organic synthesis. Furthermore, the often idiosyncratic reactivity of boronates offers unique opportunities for stereoselective manipulations. As an illustration, the oxy-Michael adduct formed in situ from **4** and phenylboronic acid under catalysis by **8** acted as a template for the stereoselective addition of allenylboronic acid to the carbonyl, possibly via intermediate **45** (eq 2). Diol **46** was generated, without isolation of intermediates, in good overall yield and diastereoselectivity.^{20,21} Further developments including diastereoselective and intermolecular oxy-Michael additions are under investigation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


Acknowledgments

¹¹B NMR measured by Dr. RenSheng Luo (UMSL). X-ray analysis by Radha Akella (UTSW). Financial support provide by the Robert A. Welch Foundation and NIH (GM31278, DK38226).

REFERENCES

- Recent review: Schetter B, Mahrwald R. *Angew. Chem., Int. Ed* 2006;45:7506–7525.
- For examples of asymmetric Michael additions of oxygen-centered nucleophiles, see: Kano T, Tanaka Y, Maruoka K. *alcohols. Tetrahedron* 2007;63:8658–8664.(b) Bertelsen S, Diner P, Johansen RL, Jorgensen KA. *oximes. J. Am. Chem. Soc* 2007;129:1536–1537. [PubMed: 17284000]Vanderwal CD, Jacobsen EN. *J. Am. Chem. Soc* 2004;126:14724–14725. [PubMed: 15535689] (c) Govender T, Hojabri L, Moghaddam FM, Arvidsson PI. *phenols. Tetrahedron: Asymmetry* 2006;17:1763–1767.
- Chandrasekhar S, Rambabu C, Shyamsunder T. *Tetrahedron Lett* 2007;48:4683–4685.

4. Evans DA, Gauchet-Prunet JA. *J. Org. Chem* 1993;58:2446–2453.
5. Chen YK, Yoshida M, MacMillan DWC. *J. Am. Chem. Soc* 2006;128:9328–9329. [PubMed: 16848457]
6. Watanabe H, Machida K, Itoh D, Nagatsuka H, Kitahara T. *Chirality* 2001;13:379–385. [PubMed: 11400192]
7. For a review of asymmetric organocatalytic oxy-Michael additions, see:
Berkessel AGroergerHAsymmetric Organocatalysis2005WeinheimWiley-VCHChapter 4
8. Falck JR, Bondlela M, Venkataraman SK, Srinivas D. *J. Org. Chem* 2001;66:7148–7150. [PubMed: 11597243]
9. (a) Yamaoka Y, Miyabe H, Takemoto Y. *J. Am. Chem. Soc* 2007;129:6686–6687. [PubMed: 17488015] (b) Oshima K, Aoyama Y. *J. Am. Chem. Soc* 1999;121:2315–2316. (c) Matteson DS. *Pure App. Chem* 2003;75:1249–1253.
10. Amine-borate complexes are well-known: Gillis EP, Burke MD. *J. Am. Chem. Soc* 2007;129:6716–6717. [PubMed: 17488084]
11. Rehybridization of the boron center from sp^2 to sp^3 via coordination with trivalent nitrogen was unambiguously supported by the upfield chemical shift in the ^{11}B NMR as described in Supporting Information section and ref. ^{9b}.
12. (a) Okino T, Hoashi Y, Takemoto Y. *J. Am. Chem. Soc* 2003;125:12672–12673. [PubMed: 14558791] (b) Vakulya B, Varga S, Csámpai A, Soos T. *Org. Lett* 2005;7:1967–1969. [PubMed: 15876031]
13. The critical role played by thiourea hydrogen bonding in the activation of the enone is evident from a control experiment in which the thiourea catalyst **8** was replaced with 20 mol% O-benzoylquinine (Shi M, Lei Z-Y, Zhao M-X, Shi J-W. *Tetrahedron Lett* 2007;48:5743–5746.) Diol **12** was obtained in <5% yield and the majority of starting **4** was recovered unchanged.
14. Enantiomeric ratio measured by chiral HPLC.
15. Absolute configuration assigned by comparison with literature optical rotation data: Ticozzi C, Zanarotti A. *Tetrahedron Lett* 1994;35:7421–7424.
16. MacLeod JK, Schäffeler L. *Synthesis. J. Nat. Prod* 1995;58:1270–1273. Interestingly, the (*S*)-enantiomer is inactive.
17. Ishihara K, Kurihara H, Yamamoto H. *J. Org. Chem* 1993;58:3791–3793.
18. Selective primary silylation of **40** (TBDPSCl, DCM, DMAP, rt, 10 h) followed by *p*-methoxybenzylation (PMB-O(NH)CCl₃) gave compound (*i*): $[\alpha]_{\text{D}}^{20} = +5.9$ (*c* 0.85, CHCl₃); lit. $[\alpha]_{\text{D}}^{20} = +4.8$ (*c* 1.49, CHCl₃). See, Li DR, Zhang D-H, Sun C-Y, Zhang J-W, Yang L, Chen J, Liu B, Su C, Zhou W-S, Lin G-Q. *Chem. Eur. J* 2006;12:1185–1204.



19. Dollt H, Hammann P, Blechert S. *Helv. Chim. Acta* 1999;82:1111–1122. and references cited therein.
20. A similar diastereoselective addition of allenylboronic acid to β -hydroxy ketones has been reported: Ikeda N, Omori K, Yamamoto H. *Tetrahedron Lett* 1986;27:1175–1178.
21. The absolute configuration of **46** was confirmed by x-ray analysis (see Supporting Information section).






Figure 1.
Proposed asymmetric catalysis

**Scheme 1.**

Biomimetic Synthesis of Antifungal/Hepatic Protective Agent from Avocadoa

^aReagents and conditions: (a) H₃CP(O)(OMe)₂, LDA, THF, -78 °C, 2 h, 90%; (b) [HC(O)CH₂OH]₂, LiCl, DIPEA, rt, CH₃CN, 2 h, 51%; (c) (i) PhB(OH)₂ (1.2 equiv), **8** (10 mol%), 4 Å MS, CH₂Cl₂, rt, 56 h; (ii) H₂O₂, Na₂CO₃, rt, 15 min; (d) AcCl (1.2 equiv), collidine, CH₂Cl₂, -78°C, 10 h.

Table 1





















Base catalyzed oxy-Michael of **4**^a


entry	base	solvent	time(h)	yield ^b (%)
1	none	CH ₂ Cl ₂	48	0
2	NaHCO ₃	CH ₂ Cl ₂	48	0
3	Na ₂ CO ₃	CH ₂ Cl ₂	48	0
4	Pyridine	CH ₂ Cl ₂	48	0
5	Et ₃ N	CH ₂ Cl ₂	48	42
6	DABCO	CH ₂ Cl ₂	48	70
7	iPr ₂ NH	CH ₂ Cl ₂	16	86
8	iPr ₂ NH	PhCH ₃	8	87
9	iPr ₂ NH	DME	48	70
10	DBU	CH ₂ Cl ₂	48	30
11	PMP ^c	PhCH ₃	80	82

^aRxn conditions: (i) PhB(OH)₂ (1.2 equiv), base (20 mol%), 4 Å MS, rt; (ii) H₂O₂, Na₂CO₃, rt, 15 min.^bOverall for two steps from enone **4** to diol **6**.^cPMP = 1,2,2,6,6-pentamethylpiperidine.

Table 2

Asymmetric oxy-Michael of γ -hydroxy- α,β -enones

entry	enone	diol	solvent	time(h)	Yield ^a (%)	ee ^b (%)
1			CH ₂ Cl ₂ ^c	16	91	91
			PhCH ₃ ^c	8	87	65
			DME ^c	24	89	94
2			DME ^d	24	90	95
			DME ^e	40	84	89
			DME ^f	48	84	91
3			DME ^d	14	92	90
4			DME ^d	27	85	97
5			DME ^d	22	83	97
6			CH ₂ Cl ₂ ^d	72	78	89
7			CH ₂ Cl ₂ ^d	72	95	87
8			CH ₂ Cl ₂ ^d	27	94	92
9			PhCH ₃ ^d	36	78	98
10			DME ^d	28	71	99

^a Isolated yield.

^b Determined by chiral HPLC; absolute configuration assigned in analogy with **12** and **33**.

^c Rxn conditions: (i) PhB(OH)₂ (1.2 equiv), **7** (10 mol%), 4 Å MS, rt; (ii) H₂O₂, Na₂CO₃, rt, 15 min.



















^d Catalyst **8** (10 mol%) was used.

^e Catalyst **9** (10 mol%) was used.

^f Catalyst **10** (10 mol%) was used.

Table 3

Oxy-Michael of δ -hydroxy- α,β -enones

entry	enone	ArB(OH) ₂	diol	time(h)	yield ^d (%)	ee ^b (%)
1				12 ^c	80	91
2				26 ^d 27 ^e	72 71	83 86
3				18 ^c	86	87
4				24 ^c	73	84
5				50 ^c	82	96
6				37 ^c	75	88

^a Isolated yield.

^b Determined by chiral HPLC; absolute configuration assigned in analogy with natural **40** and chemical correlation of **40** with a known intermediate (see reference 18).

^c Rxn conditions: (i) ArB(OH)₂ (1.2 equiv), **8** (20 mol%), 4 Å MS, toluene, 50 °C; (ii) H₂O₂, Na₂CO₃, rt, 15 min.

^d Catalyst **9** (20 mol%) was used.

^e Catalyst **10** (20 mol%) was used.