

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/265419353>

# Femtosecond Transient Absorption Spectroscopy of the Medicinal Agent Curcumin in Diamide Linked $\gamma$ -Cyclodextrin Dimers

ARTICLE in THE JOURNAL OF PHYSICAL CHEMISTRY B · SEPTEMBER 2014

Impact Factor: 3.3 · DOI: 10.1021/jp507272f · Source: PubMed

READS

37

5 AUTHORS, INCLUDING:



**Takaaki Harada**

Okinawa Institute of Science and Technology

10 PUBLICATIONS 49 CITATIONS

SEE PROFILE



**Duc-Truc Pham**

University of Adelaide

27 PUBLICATIONS 136 CITATIONS

SEE PROFILE



**Stephen F. Lincoln**

University of Adelaide

336 PUBLICATIONS 4,041 CITATIONS

SEE PROFILE



**Tak W Kee**

University of Adelaide

44 PUBLICATIONS 1,276 CITATIONS


SEE PROFILE

Femtosecond Transient Absorption Spectroscopy of the Medicinal Agent Curcumin in Diamide Linked  $\gamma$ -Cyclodextrin Dimers

Takaaki Harada, Hamish L. McTernan, Duc-Truc Pham, Stephen F. Lincoln, and Tak W. Kee\*

Department of Chemistry, The University of Adelaide, Adelaide, South Australia 5005, Australia

**ABSTRACT:** Curcumin is a biologically active polyphenol and a yellow pigment extracted from turmeric. Our previous study has shown effective encapsulation of curcumin using diamide linked  $\gamma$ -cyclodextrin dimers, namely 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur, through cooperative 1:1 host-guest complexation. In this study, the excited-state dynamics of curcumin complexed with either 66 $\gamma$ CD<sub>2</sub>su or 66 $\gamma$ CD<sub>2</sub>ur in water are investigated using femtosecond transient absorption spectroscopy. Both 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes in water show only an excited-state absorption (ESA) band at 530 nm without any stimulated emission (SE) signals, indicating non-radiative decays as the major relaxation pathways. The ESA dynamics of 66 $\gamma$ CD<sub>2</sub>su-curcumin are similar to those of 66 $\gamma$ CD<sub>2</sub>ur-curcumin, consisting of a rapid growth component and three decay components. The growth component, which has a time constant of 0.25–0.41 ps, is assigned to solvent reorganization. The relatively fast decay components with time constants of 9.3–21.8 ps show significant deuterium isotope effect, consistent with the presence of excited-state intramolecular hydrogen atom transfer (ESIHT) of curcumin. The small-amplitude and slow decay components may be attributed to the dynamics of complexed curcumin and molecular motions due to flexibility of 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur. In addition, transient absorption anisotropy measurements reveal slow rotational motions of 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes. The overall results show that complexation in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur has pronounced effects on the photophysics of curcumin.



## ■ INTRODUCTION

Curcumin (Figure 1a) is a naturally occurring yellow polyphenol present in the rhizomes of the spice plant *Curcuma longa*, commonly known as turmeric.<sup>1,2</sup> It constitutes 77% of curcuminoids, which are composed of a group of curcumin analogues, in company with demethoxycurcumin (17%) and bisdemethoxycurcumin (3%).<sup>3</sup> Another curcuminoid, cyclocurcumin, which is also present but at a much lower level,<sup>4</sup> was recently studied.<sup>5</sup> The dominant conformation of curcumin is the keto-enol form in polar solvents (Figure 1a).<sup>1,2</sup> Recent research on curcumin has shown its medicinal effects, including anticancer,<sup>6–10</sup> anti-Alzheimer's disease,<sup>11–13</sup> anti-cystic fibrosis,<sup>14,15</sup> and anti-inflammation properties.<sup>16</sup> However, the poor aqueous stability and solubility of curcumin limit bioavailability and hence hinder applications of curcumin as an effective therapeutic drug.<sup>17–19</sup> Therefore, molecular assemblies as delivery systems, which include micelles,<sup>20–23</sup> globular proteins,<sup>24,25</sup> polymer nanoparticles,<sup>26,27</sup> micelle-like aggregates and hydrogels,<sup>28</sup> and cyclodextrins,<sup>29,30</sup> have been developed and investigated to improve the availability of curcumin in vivo.

Cyclodextrins (CDs) are naturally occurring cyclic oligosaccharides with either 6 ( $\alpha$ ), 7 ( $\beta$ ), or 8 ( $\gamma$ ) glucopyranoside units. Figure 1b shows the structure of  $\gamma$ -CD. The hydrophobic interior of CDs has the ability to encapsulate a hydrophobic molecular species, while the hydrophilic exterior allows the CD-drug host-guest complexes to be suspended in water, which is attractive for delivering a hydrophobic drug.<sup>31,32</sup> However, this approach is limited to small molecule drugs due to the limited


cavity size of CDs.<sup>33</sup> In order to further improve the CD-based drug delivery systems, a diamide linker between two CDs has been introduced to promote cooperative binding to large molecular guests.<sup>34–36</sup> Our previous study has shown a strong 1:1 cooperative binding of diamide linked  $\gamma$ -CD dimers, namely 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur, to curcumin (Figures 1c and 1d), which results in a remarkable aqueous stability of curcumin under physiological conditions.<sup>29</sup>

Our recent study has also reported the use of 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur as curcumin delivery systems to cancer cells,<sup>6</sup> and other studies have shown that curcumin has a range of phototherapeutic effects on melanoma cells.<sup>37–39</sup> There is a growing interest in understanding the photophysical processes of curcumin.<sup>40–47</sup> Thus far, femtosecond transient absorption spectroscopy and fluorescence upconversion spectroscopy have shown that the dominant relaxation pathway of curcumin is excited-state intramolecular hydrogen atom transfer (ESIHT).<sup>40–45</sup> The ESIHT process is influenced by polar solvents including methanol, DMSO, and acetone due to solvent-curcumin interactions. Although there have been reports of a relatively fast ESIHT of curcumin in nonpolar

**Special Issue:** Photoinduced Proton Transfer in Chemistry and Biology Symposium

**Received:** July 21, 2014

**Revised:** August 25, 2014



**Figure 1.** Structures of (a) curcumin, (b)  $\gamma$ -CD, and curcumin complexed with (c) 66 $\gamma$ CD<sub>2</sub>su and (d) 66 $\gamma$ CD<sub>2</sub>ur.

solvent systems,<sup>43,44,47</sup> it is still highly unlikely that the observed ESIHT occurs in a direct manner, possibly due to the presence of trace quantities of water in these solvents. Previous work on 3-hydroxyflavone showed that trace amounts of hydrogen-bonding substances can have a significant effect on ESIHT.<sup>48</sup> Furthermore, Schwartz et al. showed that even with rigorous purification of methylcyclohexane traces of hydrogen-bonding impurities (e.g., H<sub>2</sub>O) are still present, affecting ESIHT of 3-hydroxyflavone.<sup>49</sup> Nevertheless, the study by Schwartz et al. revealed that ESIHT of 3-hydroxyflavone occurs with a time constant of 240 fs in a highly purified and dried environment.<sup>49</sup> This work provided the first insight into the ultrafast timescale for a direct ESIHT. Interestingly, a very recent study by Mohammed et al. showed that ESIHT of 1,8-dihydroxy-9,10-anthraquinone (DHAQ) involves a direct hydrogen atom transfer from an enol to a keto group in a wide range of solvents.<sup>50</sup> The results suggest that ESIHT of DHAQ occurs within the first 150 fs of the dynamics. Therefore, it follows that a direct ESIHT of curcumin should occur with a similar time constant to those of 3-hydroxyflavone and DHAQ. In addition to the timescale of ESIHT, there have been some debates on whether the excited-state hydrogen atom transfer reaction of curcumin is intramolecular or intermolecular in nature in polar protic solvents.<sup>41,42,47</sup> While Adhikary et al. proposed an intramolecular pathway,<sup>41,42</sup> Ghosh et al. suggested intermolecular hydrogen-bonding of curcumin with

the polar protic solvent as an excited-state hydrogen atom transfer mechanism.<sup>47</sup> Although it appears that there are two different pathways, these two proposals are in fact very similar, if not identical. This is because the intramolecular pathway proposed by Adhikary et al. actually involves interactions of curcumin with the polar protic solvent, which is consistent with the conclusions drawn by Kasha and Maroncelli on the ESIHT reactions of 3-hydroxyflavone and 7-azaindole, respectively.<sup>48,51</sup> Therefore, rather than having two different mechanisms for excited-state hydrogen atom transfer, there is in fact a consensus in the field that interactions with polar protic solvents play an important role in ESIHT of curcumin.

Here, we report the excited-state photodynamics of curcumin complexed by the cyclodextrin dimers, 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur, for the first time. The steady-state absorption and fluorescence spectra show that curcumin is nonfluorescent when it is complexed by 66 $\gamma$ CD<sub>2</sub>su or 66 $\gamma$ CD<sub>2</sub>ur. Femtosecond transient absorption spectroscopy reveals excited-state absorption of curcumin complexed by 66 $\gamma$ CD<sub>2</sub>su or 66 $\gamma$ CD<sub>2</sub>ur in water, without any stimulated emission signals. The non-radiative decay processes involve ESIHT of curcumin, reorganization of water molecules within the cavity of 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur, and other processes including dynamics of complexed curcumin and molecular motions due to flexibility of the  $\gamma$ -CD moieties of 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur. Time-resolved anisotropy results show that the rotational motions of curcumin complexed by 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur occur on a significantly longer timescale than that of curcumin in a polar organic solvent. Overall, the femtosecond transient absorption results provide insight into the photophysical properties of curcumin when it is complexed by 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur.

## MATERIALS AND METHODS

**Materials.** Curcumin (1,7-bis(4-hydroxy-3-methoxyphenyl)hepta-1,6-diene-3,5-dione) was obtained from LKT Laboratories (purity >98%). A previous study showed that curcumin with this purity was still comparable to that with a 70% purity,<sup>42</sup> and hence, the 2% impurities were unlikely to affect our spectroscopic measurements. Methanol (MeOH, HPLC grade, 99.7%) from Scharlau was used as received, and deuterated water (D<sub>2</sub>O) and methanol (MeOD-*d*<sub>4</sub>) were purchased from Cambridge Isotope Laboratories (D, 99.9%). Water (H<sub>2</sub>O) was obtained from a Millipore Milli-Q NANO-pure water system. The pH of H<sub>2</sub>O and pD of D<sub>2</sub>O were approximately 7. The C6<sup>A</sup>-to-C6<sup>A</sup> diamide linked  $\gamma$ -cyclodextrin dimers, *N,N'*-bis(6<sup>A</sup>-deoxy- $\gamma$ -cyclodextrin-6<sup>A</sup>-yl) succinamide (66 $\gamma$ CD<sub>2</sub>su), and *N,N'*-bis(6<sup>A</sup>-deoxy- $\gamma$ -cyclodextrin-6<sup>A</sup>-yl) urea (66 $\gamma$ CD<sub>2</sub>ur), were synthesized using literature methods.<sup>36</sup> Briefly, the native  $\gamma$ -CDs were substituted with 4-toluenesulfonyl chloride for activation at the C6<sup>A</sup> position, which yielded 6<sup>A</sup>-O-(4-methylbenzenesulfonyl)- $\gamma$ -cyclodextrin (6 $\gamma$ CDTs). For the synthesis of 66 $\gamma$ CD<sub>2</sub>su, the reaction between 6 $\gamma$ CDTs and ammonium bicarbonate produced 6<sup>A</sup>-amino-6<sup>A</sup>-deoxy- $\gamma$ -cyclodextrin, 6 $\gamma$ CDNH<sub>2</sub>, which was then dimerized by the reaction with bis(4-nitrophenyl) succinate as the linker. For the synthesis of 66 $\gamma$ CD<sub>2</sub>ur, the reaction between 6 $\gamma$ CDTs and sodium azide produced 6<sup>A</sup>-azido-6<sup>A</sup>-deoxy- $\gamma$ -cyclodextrin, 6 $\gamma$ CDN<sub>3</sub>, which was then dimerized by reaction with carbon dioxide as the linker.

**Steady-State UV-Visible Absorption and Fluorescence Spectroscopic Studies.** UV-visible absorption spectra of 1  $\mu$ M curcumin complexed with either 66 $\gamma$ CD<sub>2</sub>su or 66 $\gamma$ CD<sub>2</sub>ur in

a 1:1 ratio in H<sub>2</sub>O or D<sub>2</sub>O in a 1-cm quartz cuvette were recorded from 300 to 700 nm using a Cary 5000 UV-Vis/NIR spectrophotometer (Varian). Similarly, UV-visible absorption spectra of 1  $\mu$ M curcumin in either MeOH or MeOD-*d*<sub>4</sub> were acquired from 300 to 700 nm. Fluorescence spectra of these solutions were subsequently recorded from 415 to 700 nm using a Cary Eclipse Fluorescence spectrophotometer (Varian) with the excitation and emission slit widths set at 5 nm. The excitation wavelength for these experiments was set at 400 nm. The reported spectra were averaged over 5 scans at a scan rate of 600 nm/min.


**Femtosecond Transient Absorption Spectroscopic Studies.** Solutions of 80  $\mu$ M curcumin complexed with either 66 $\gamma$ CD<sub>2</sub>su or 66 $\gamma$ CD<sub>2</sub>ur in a 1:1 ratio in H<sub>2</sub>O or D<sub>2</sub>O were used in the transient absorption spectroscopic studies. Similarly, a curcumin concentration of 80  $\mu$ M was used in MeOH or MeOD-*d*<sub>4</sub>. All the measurements were acquired using a quartz cuvette with a 2-mm path length. Less than 10% of curcumin photodegradation was observed after each set of data acquisition.

The laser system used for the femtosecond transient absorption experiments consisted of a Ti:sapphire mode-locked oscillator (Spectra-Physics, Tsunami), which seeded a Ti:sapphire regenerative amplifier (Spectra-Physics, Spitfire Pro XP) pumped by a 20W Q-switched Nd:YLF laser (Spectra-Physics, Empower). The output of the amplifier was centered at 800 nm with a repetition rate of 1 kHz and pulse duration of 100 fs, which was then split into pump and probe beamlines. The 400-nm pump pulses were generated using a type-I BBO crystal (Eksma Optics), which was modulated at 500 Hz and then focused onto the sample with a spot size of 710  $\mu$ m and pulse energy of 720 nJ. The probe beam passed through a delay stage and was used to generate a white light continuum in a 2-mm sapphire crystal. The probe passed through a beam splitter to produce the sample and reference beams with a spot size of 50  $\mu$ m at the sample position. The sample and reference beams were then directed into complementary CMOS detectors for detection in the visible region. The probe polarization was oriented at magic angle (54.7°) with respect to the pump polarization. In all the measurements, a full width at half maximum (fwhm) of 100 fs was used for data analysis.

The same laser system was utilized for the femtosecond transient absorption anisotropy study. The measurement involved collecting two transient absorption signals with the probe polarization at 0° (parallel,  $I_{||}$ ) and 90° (perpendicular,  $I_{\perp}$ ) with respect to the pump polarization, in order to determine the time-dependent anisotropy  $r(t)$  (see eq 1). In this study, a concentration of 160  $\mu$ M was used for 66 $\gamma$ CD<sub>2</sub>su or 66 $\gamma$ CD<sub>2</sub>ur while the concentration of curcumin was kept at 80  $\mu$ M curcumin to minimize the probability of multiple curcumin molecules binding to a single diamide linked  $\gamma$ -CD dimer.

## RESULTS AND DISCUSSION

**Steady-State Absorption and Fluorescence Spectra of Curcumin Complexed in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur.** The steady-state absorption and fluorescence spectra of the 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes are displayed in Figures 2a and 2b, respectively. The spectral envelopes of the curcumin complexes in H<sub>2</sub>O and D<sub>2</sub>O are very similar except for a minor blue shift for the D<sub>2</sub>O solution. In comparison with H<sub>2</sub>O and D<sub>2</sub>O solvent systems, Figure 2c shows the steady-state UV–visible absorption and fluorescence



**Figure 2.** Steady-state absorption and fluorescence spectra of curcumin complexed with (a) 66 $\gamma$ CD<sub>2</sub>su and (b) 66 $\gamma$ CD<sub>2</sub>ur in H<sub>2</sub>O (red) and D<sub>2</sub>O (blue), and (c) curcumin in MeOH (red) and MeOD-*d*<sub>4</sub> (blue), respectively. The gray dashed spectra in (a) and (b) represent steady-state absorption and fluorescence spectra of curcumin in MeOH as a reference. The fluorescence spectra of 66 $\gamma$ CD<sub>2</sub>su-curcumin or 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes in H<sub>2</sub>O or D<sub>2</sub>O were multiplied by ten for illustration purposes.

spectra of curcumin in MeOH (red) and MeOD-*d*<sub>4</sub> (blue). The absorption peak around 430 nm corresponds to the  $\pi$ – $\pi^*$  transition of the keto-enol tautomer of curcumin (Figure 1a).<sup>20,42</sup> The absorption spectrum of curcumin in MeOD-*d*<sub>4</sub> exhibits a very small blue shift without observable spectral changes.<sup>52</sup> The absorption peak around 420 nm and spectral envelope of 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin are similar to those of curcumin in MeOH, which are shown as a gray dashed spectrum in Figures 2a and 2b. This result indicates that the keto-enol tautomer of curcumin is present in the 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes, as shown in Figures 1c and 1d.<sup>29</sup>


Additionally, the absorption spectra of both 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes show no 350-nm spectral shoulder, indicating a lack of significant water-curcumin interactions.<sup>20,24,29</sup> However, the water-curcumin interactions are still sufficiently strong such that the 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes exhibit very weak fluorescence signals (Figures 2a and 2b).<sup>29</sup> At the same curcumin concentration, the fluorescence spectrum of curcumin in MeOD-*d*<sub>4</sub> exhibits a higher intensity than that of curcumin in MeOH, while their spectral envelopes remain similar (Figure 2c).<sup>52</sup> The fluorescence intensities of 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin in D<sub>2</sub>O are also higher than those in H<sub>2</sub>O. It is clear that the fluorescence intensity and hence the fluorescence quantum yield of curcumin is sensitive to the hydrogen/deuterium (H/D) exchange of curcumin.

It has been shown previously that the enolic hydrogen of curcumin undergoes H/D exchange in deuterated protic solvents and changes the fluorescence quantum yields of



curcumin.<sup>52</sup> A previous time-resolved fluorescence spectroscopic study has concluded that ESIHT is a major photophysical process of curcumin in polar protic solvents.<sup>42</sup> The higher fluorescence quantum yield of curcumin in MeOD-*d*<sub>4</sub> than that in MeOH is due to a longer fluorescence lifetime as a result of the deuterium isotope effect.<sup>42,52</sup> Although time-resolved fluorescence spectroscopy has been very useful in previous work, it is inapplicable to 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin due to their nonfluorescent nature.<sup>29</sup> Hence, in this study, transient absorption spectroscopy was used to offer insight into the excited-state dynamics of the 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes.

**Femtosecond Transient Absorption of Curcumin Complexed in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur.** Femtosecond transient absorption spectroscopy has been employed for investigations of the excited-state dynamics of curcumin in different systems.<sup>40,47</sup> Here we report the first femtosecond transient absorption study on curcumin solely complexed in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur, in a 1:1 host:guest ratio in the aqueous environment. Figures 3a and 3b show the transient absorption spectra of the 66 $\gamma$ CD<sub>2</sub>su-curcumin complex in H<sub>2</sub>O and D<sub>2</sub>O, respectively, with several probe delay times, ranging from 0.1 to 200 ps. The excited-state absorption (ESA) band of curcumin complexed in 66 $\gamma$ CD<sub>2</sub>su in H<sub>2</sub>O at 0.1 ps shows a 500-nm peak, but the 530-nm ESA signal becomes pronounced




**Figure 3.** Transient absorption spectra of 66 $\gamma$ CD<sub>2</sub>su-curcumin complex in (a) H<sub>2</sub>O and (b) D<sub>2</sub>O, those of 66 $\gamma$ CD<sub>2</sub>ur-curcumin complex in (c) H<sub>2</sub>O and (d) D<sub>2</sub>O, and those of curcumin in (e) MeOH and (f) MeOD-*d*<sub>4</sub>, at different time delays. Each set of spectra is normalized to the maximum  $\Delta$ OD signal of the sample. The gray dashed spectrum in (a–d) represents transient absorption spectra of curcumin in either (e) MeOH or (f) MeOD-*d*<sub>4</sub> at 30 ps as a reference.

at >1 ps. Figures 3a and 3b show that the transient absorption dynamics of 66 $\gamma$ CD<sub>2</sub>su-curcumin in H<sub>2</sub>O are faster than those in D<sub>2</sub>O, which is discussed below. A similar behavior in the transient absorption spectra are also present when curcumin is complexed with 66 $\gamma$ CD<sub>2</sub>ur in H<sub>2</sub>O or D<sub>2</sub>O, as shown in Figures 3c and 3d, respectively.

Figures 3e and 3f show the transient absorption spectra of curcumin in MeOH and MeOD-*d*<sub>4</sub>, respectively, with several probe delay times ranging from 0.1 to 200 ps. Curcumin in MeOH exhibits a rapid ESA peak around 600 nm at 0.1 ps, which is followed by the appearance of another ESA peak around 500 nm and a stimulated emission (SE) signal around 540 nm at 1 ps. These early time ESA and SE peaks of curcumin in MeOH were also observed in a previous study.<sup>40</sup> The ESA at 600 nm rapidly evolves into SE between 10 and 200 ps. The SE signal observed at 540 nm becomes a part of the ESA band centered at 500 nm around 10 ps. This band reaches a maximum  $\Delta$ OD value around 30 ps and decreases thereafter.

There are three notable differences between the transient absorption spectra of *complexed* and *free* curcumin. First, the transient absorption spectra of 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes only exhibit ESA signals within the time window of investigation. The transient absorption spectrum of curcumin in MeOH and MeOD-*d*<sub>4</sub> at 30 ps are shown in Figures 3a–3d as gray dashed spectra to highlight the absence of SE signals of curcumin complexed in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur. The absence of SE signals for the 66 $\gamma$ CD<sub>2</sub>su-curcumin or 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes is consistent with the very weak fluorescence signals, as shown in Figures 2a and 2b. Second, there is an apparent difference in the maximum ESA peak position between complexed and free curcumin. Curcumin complexed in either 66 $\gamma$ CD<sub>2</sub>su or 66 $\gamma$ CD<sub>2</sub>ur only shows an ESA peak at 530 nm. In contrast, the ESA maximum of free curcumin appears at 500 nm. However, the ESA and SE signals of free curcumin overlap between 450 and 700 nm such that the overall ESA maximum of free curcumin appears at 500 nm. Finally, the ESA signals of the 66 $\gamma$ CD<sub>2</sub>su-curcumin or 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes evolve faster than that of curcumin in methanol, as is discussed below.

**Excited-State Dynamics of Curcumin Complexed in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur.** Figures 4a and 4b show the 500-nm ESA signals of curcumin complexed with 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur, respectively, in H<sub>2</sub>O (red) and D<sub>2</sub>O (blue) as a function of pump-probe delay time. The 500-nm ESA signals of 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes were fitted with a quad-exponential function consisting of a growth component ( $\tau_1$ ) and three decay components ( $\tau_2$ ,  $\tau_3$ , and  $\tau_4$ ), which are summarized in Table 1. The ESA signals of 66 $\gamma$ CD<sub>2</sub>su-curcumin complex in H<sub>2</sub>O and D<sub>2</sub>O show a rapid growth component with a time constant ( $\tau_1$ ) of 0.25 ps with an amplitude of approximately ~40% of the maximum signal amplitude. Comparable results were obtained for the 66 $\gamma$ CD<sub>2</sub>ur-curcumin complex (Table 1). There are three decay components characterized by time constants,  $\tau_2$ ,  $\tau_3$ , and  $\tau_4$ , for the ESA signals of both the 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes (Table 1). The time constants  $\tau_2$  are approximately 9.3 ps (39%) and 14.6 ps (34%) for the 66 $\gamma$ CD<sub>2</sub>su-curcumin complex in H<sub>2</sub>O and D<sub>2</sub>O, respectively. For the 66 $\gamma$ CD<sub>2</sub>ur-curcumin complex in H<sub>2</sub>O and D<sub>2</sub>O solvent systems, the time constants  $\tau_2$  are 12.7 ps (55%) and 21.8 ps (53%), respectively. The time constants  $\tau_3$  and  $\tau_4$  are fixed to 109 and 1200 ps, respectively,<sup>53</sup> which are discussed below.



**Figure 4.** Normalized excited-state absorption signals of (a)  $66\gamma\text{CD}_2\text{su}$ -curcumin and (b)  $66\gamma\text{CD}_2\text{ur}$ -curcumin complexes in  $\text{H}_2\text{O}$  (red) and  $\text{D}_2\text{O}$  (blue), and (c) those of curcumin in  $\text{MeOH}$  (red) and  $\text{MeOD-}d_4$  (blue), at 500 nm. The insets show the signals at later times. Note the timescales of (a) and (b) are different from that of (c).

The rapid growth components  $\tau_1$  of the ESA signal for both  $66\gamma\text{CD}_2\text{su}$ -curcumin and  $66\gamma\text{CD}_2\text{ur}$ -curcumin complexes are assigned to reorganization of  $\text{H}_2\text{O}$  and  $\text{D}_2\text{O}$ ,<sup>41,53,54</sup> which is consistent with the Stokes shift observed in the steady-state spectra (Figures 2a and 2b). Vajda et al. have investigated the dynamics of water molecules inside the  $\gamma$ -CD cavity, and their results indicate the presence of significant solute-solvent interactions inside the  $\gamma$ -CD cavity, occurring within 1 ps.<sup>53</sup> Therefore, the interactions between curcumin and water molecules inside and cavities are expected. The presence of water-curcumin interactions is consistent with the non-fluorescent nature of  $66\gamma\text{CD}_2\text{su}$ -curcumin and  $66\gamma\text{CD}_2\text{ur}$ -curcumin complexes,<sup>29</sup> and the results from a previous

molecular dynamics study on these complexes.<sup>55</sup> The process with a time constant of  $\tau_2$  is the dominant relaxation process and shows a deuterium isotope effect of 1.6 for  $66\gamma\text{CD}_2\text{su}$ -curcumin and 1.7 for  $66\gamma\text{CD}_2\text{ur}$ -curcumin. This relaxation is attributable to ESIHT, which is in agreement with previous studies on curcumin in protic solvents including methanol.<sup>41,42</sup> The substantial contribution of the decay component  $\tau_2$  (approximately 35–55%) indicates that ESIHT is an important photophysical process of excited-state curcumin complexed with  $66\gamma\text{CD}_2\text{su}$  and  $66\gamma\text{CD}_2\text{ur}$ . However, the resultant deuterium isotope effect of  $\tau_2$  is less than 2, indicating that other relaxation processes may occur at similar timescales to ESIHT. The relative amplitude of the ESIHT component is lower for  $66\gamma\text{CD}_2\text{su}$ -curcumin (37%) than  $66\gamma\text{CD}_2\text{ur}$ -curcumin (54%), as shown in Table 1. The difference in relative amplitude is likely to be related to the difference in the hydrogen-bonding interactions of curcumin in these assemblies. A recent computational study suggested a significant level of hydrogen bonding between curcumin and the diamide linker of  $66\gamma\text{CD}_2\text{ur}$ .<sup>55</sup> However, this interaction is absent between curcumin and  $66\gamma\text{CD}_2\text{su}$  due to the high flexibility of the diamide linker. The long decay time constants  $\tau_3$  and  $\tau_4$  are present in the ESA decay without showing any deuterium isotope effect. These small-amplitude and slow relaxation processes may involve motions of complexed curcumin and the  $\gamma$ -CD moieties of  $66\gamma\text{CD}_2\text{su}$  and  $66\gamma\text{CD}_2\text{ur}$  due to their flexibility. A previous study has shown that the limited cavity size of  $\gamma$ -CD leads to the appearance of small-amplitude and slow relaxation processes, including diffusive motions of the guest molecule, reorientation of highly constrained water molecules within the  $\gamma$ -CD cavity, or fluctuations of the  $\gamma$ -CD molecule.<sup>53</sup> In the same study, the time constants of 109 and 1200 ps were assigned to these molecular motions.<sup>53</sup> Here, we fixed  $\tau_3$  and  $\tau_4$  to these values as these molecular motions are expected to be present in both  $66\gamma\text{CD}_2\text{su}$ -curcumin and  $66\gamma\text{CD}_2\text{ur}$ -curcumin complexes. Similar results have also been reported in other studies.<sup>56,57</sup> Moreover, a recent computational study has shown the presence of water molecules and diffusive motions of curcumin within the  $\gamma$ -CD cavities of  $66\gamma\text{CD}_2\text{su}$  and  $66\gamma\text{CD}_2\text{ur}$ .<sup>55</sup> In addition, our previous 2D NOESY  $^1\text{H}$  NMR spectroscopic results show the intermolecular alkyl hydrogen-hydrogen interactions between curcumin and the  $\gamma$ -CD moieties of  $66\gamma\text{CD}_2\text{su}$  and  $66\gamma\text{CD}_2\text{ur}$ .<sup>29</sup> In short, the slow relaxation dynamics of  $66\gamma\text{CD}_2\text{su}$ -curcumin and  $66\gamma\text{CD}_2\text{ur}$ -curcumin complexes may be derived from fluctuations of the  $\gamma$ -CD moieties, reorientation of constrained water molecules, diffusive motions of curcumin within the  $\gamma$ -CD

**Table 1.** Transient Absorption Kinetic Parameters of Curcumin in Different Systems at 500 nm<sup>a</sup>

| host                           | solvent              | $a_1^b$ | $\tau_1$ (ps) | $a_2^b$ | $\tau_2$ (ps) | $a_3$ | $\tau_3^c$ (ps) | $\langle\tau\rangle^d$ (ps) |
|--------------------------------|----------------------|---------|---------------|---------|---------------|-------|-----------------|-----------------------------|
| $66\gamma\text{CD}_2\text{su}$ | $\text{H}_2\text{O}$ | −0.42   | 0.25          | 0.39    | 9.3           | 0.14  | 109             | 126                         |
| $66\gamma\text{CD}_2\text{su}$ | $\text{D}_2\text{O}$ | −0.44   | 0.25          | 0.34    | 14.6          | 0.14  | 109             | 210                         |
| $66\gamma\text{CD}_2\text{ur}$ | $\text{H}_2\text{O}$ | −0.38   | 0.33          | 0.55    | 12.7          | 0.04  | 109             | 83                          |
| $66\gamma\text{CD}_2\text{ur}$ | $\text{D}_2\text{O}$ | −0.36   | 0.41          | 0.53    | 21.8          | 0.08  | 109             | 103                         |
| —                              | $\text{MeOH}$        | −0.06   | 1.17          | −0.43   | 9.4           | 0.51  | 116             | 116                         |
| —                              | $\text{MeOD-}d_4$    | −0.07   | 1.81          | −0.42   | 10.6          | 0.51  | 231             | 231                         |

<sup>a</sup>The transient absorption kinetics were fitted with the multiexponential function,  $f(t) = \sum_{i=1}^n a_i \exp(-t/\tau_i)$ , where  $\sum_{i=1}^n |a_i| = 1$ , with the minimum number of exponential terms ( $n$ ). The relative error values of the parameters are approximately 10% from three independent measurements. <sup>b</sup>The negative amplitude signifies a growth component. <sup>c</sup>The  $\tau_3$  and  $\tau_4$  values are fixed to 109 and 1200 ps, respectively, for curcumin complexed in  $66\gamma\text{CD}_2\text{su}$  and  $66\gamma\text{CD}_2\text{ur}$ .<sup>53</sup> See text for details. <sup>d</sup> $\langle\tau\rangle$  is defined as  $(\sum_{i=1}^k a_i \tau_i) / \sum_{i=1}^k a_i$  for the decay components ( $k$ ).

cavities, and/or alkyl hydrogen-hydrogen vibrational motions between curcumin and the  $\gamma$ -CD moieties.

Figure 4c shows the time-dependent 500-nm ESA signals of free curcumin in MeOH (red) and MeOD- $d_4$  (blue), which decay slower than those of curcumin complexed in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur. The 500-nm ESA signals of curcumin in MeOH and MeOD- $d_4$  are fitted with a triexponential function with two growth components and a single decay component, which are shown in Table 1. The two growth components, with time constants  $\tau_1$  of 1.17 ps (6%) and  $\tau_2$  of 9.4 ps (43%), are in agreement with previous studies.<sup>40,47</sup> The growth components in the ESA signals of curcumin in MeOD- $d_4$  result in time constants  $\tau_1$  of 1.81 ps (7%) and  $\tau_2$  of 10.6 ps (42%), which are similar to those for curcumin in MeOH. These time constants for curcumin in either MeOH or MeOD- $d_4$  are assigned to reorganization of solvent molecules. In addition to the growth components, the ESA signals of curcumin in MeOH show a single decay component with a time constant  $\tau_3$  of 116 ps (51%), as previously reported.<sup>40,47</sup> The decay component in the ESA signal of curcumin in MeOD- $d_4$  results in a time constant  $\tau_3$  of 231 ps, showing a significant deuterium isotope effect of 2.0. This decay component is related to ESIHT, as previously investigated using femtosecond fluorescence upconversion spectroscopy.<sup>41,42</sup> Furthermore, the ESIHT process is influenced by polar solvent molecules, including methanol, due to significant solvent-curcumin interactions, resulting in a longer time constant of free curcumin in MeOH than that of curcumin complexed in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur.

Overall, the non-radiative relaxation processes of curcumin complexed in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur are determined to be solvent reorganization, ESIHT, and other slow dynamics. For free curcumin in MeOH and MeOD- $d_4$ , solvent reorganization and ESIHT are the non-radiative relaxation processes. In all cases, the ESIHT is a major relaxation process and it plays an important role in the photophysical properties of curcumin.

**Anisotropy of Curcumin Complexed in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur.** To gain further insight into the dynamics of curcumin, the rotational motions of curcumin complexed in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur were investigated and compared with that of free curcumin in methanol. Transient absorption anisotropy involves measurements of two transient absorption signals with the probe polarization at 0° (parallel,  $I_{\parallel}$ ) and 90° (perpendicular,  $I_{\perp}$ ) with respect to the pump polarization. The polarization anisotropy function,  $r(t)$ , is defined as follows.


$$r(t) = \frac{I_{\parallel}(t) - I_{\perp}(t)}{I_{\parallel}(t) + 2I_{\perp}(t)} \quad (1)$$

The anisotropy decay,  $r(t)$ , is fitted using a single exponential function,

$$r(t) = r_0 \exp(-t/\tau^{(r)}) \quad (2)$$

where  $r_0$  and  $\tau^{(r)}$  are the initial anisotropy value and molecular rotation time constant, respectively.

Figures 5a and 5b show the anisotropy decays of 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin in H<sub>2</sub>O, respectively. The 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin in H<sub>2</sub>O show  $r_0 = 0.33 \pm 0.01$  and  $0.32 \pm 0.01$ , respectively. The deviation from the theoretical limit of 0.4 in both cases may be due to rapid energy transfer from curcumin to either 66 $\gamma$ CD<sub>2</sub>su or 66 $\gamma$ CD<sub>2</sub>ur and/or to the surrounding solvent molecules. It is clear from Figures 5a and 5b that  $\tau^{(r)}$  of 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin in H<sub>2</sub>O are significantly longer than



**Figure 5.** Transient absorption anisotropy decays of (a) 66 $\gamma$ CD<sub>2</sub>su-curcumin in H<sub>2</sub>O, (b) 66 $\gamma$ CD<sub>2</sub>ur-curcumin in H<sub>2</sub>O, and (c) curcumin in MeOH, at 500 nm. Note the timescales of (a) and (b) are different from that of (c). The insets show the anisotropy variations in the first 175 ps. Refer to eq 1 for the definitions of  $I_{\parallel}(t)$  and  $I_{\perp}(t)$ .

the time window available for polarization anisotropy measurements. The  $\tau^{(r)}$  value of >1200 ps produces a reasonable fit to  $r(t)$  for these species. In contrast, Figure 5c shows the anisotropy decay of curcumin in MeOH, with  $r_0 = 0.34 \pm 0.01$  ps and  $\tau^{(r)} = 249 \pm 13$  ps. These values are in good agreement with those from a previous study.<sup>42</sup> It is clear that rotational motions are present for curcumin in MeOH within the experimental time window. The slow anisotropy decays of 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin indicate substantially slower rotational motions than that of curcumin in MeOH, as summarized in Table 2.

To understand the difference between the  $\tau^{(r)}$  of curcumin in MeOH and in the diamide linked  $\gamma$ -CD dimers, we turn to the Debye-Stokes-Einstein relation to estimate the rotational time constant.

**Table 2. Time-Resolved Transient Absorption Anisotropy Decay Parameters of Curcumin in Different Systems at 500 nm<sup>a</sup>**

| host                           | solvent          | $r_0$           | $\tau^{(r)}$ (ps) | $\tau_{\text{DSE}}^{(r)}$ (ps) |
|--------------------------------|------------------|-----------------|-------------------|--------------------------------|
| 66 $\gamma$ CD <sub>2</sub> su | H <sub>2</sub> O | $0.33 \pm 0.01$ | >1200             | 869–1189                       |
| 66 $\gamma$ CD <sub>2</sub> ur | H <sub>2</sub> O | $0.32 \pm 0.01$ | >1200             | 869–998                        |
| –                              | MeOH             | $0.34 \pm 0.01$ | $249 \pm 13$      | 375                            |

<sup>a</sup>The transient absorption anisotropy decays were fitted to  $r(t) = r_0 \exp(-t/\tau^{(r)})$ . <sup>b</sup>Debye-Stokes-Einstein equation at 293 K (refer to eq 3).



$$\tau_{\text{DSE}}^{(r)} = V\eta/k_{\text{B}}T \quad (3)$$

The symbols  $V$ ,  $\eta$ ,  $k_{\text{B}}$ , and  $T$  are the hydrodynamic volume of the species under investigation, the viscosity of the medium, the Boltzmann constant, and the temperature in Kelvin, respectively. With the use of  $\eta_{\text{MeOH}} = 5.90 \times 10^{-4}$  Pa s at 293 K and a hydrodynamic radius of 0.85 nm for curcumin,<sup>58</sup>  $\tau_{\text{DSE}}^{(r)}$  of curcumin in MeOH is 375 ps, which is in general agreement with the experimentally determined  $\tau^{(r)}$  of  $\sim 250$  ps. Similarly, with the use of the lengths of the succinamide and urea linkers and the dimensions of  $\gamma$ -CD,<sup>33</sup> the hydrodynamic volumes of 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur are estimated to be  $(3.50\text{--}4.79) \times 10^{-27}$  m<sup>3</sup> and  $(3.50\text{--}4.02) \times 10^{-27}$  m<sup>3</sup>, respectively, depending on the distances between the  $\gamma$ -CD moieties owing to flexibility of the diamide linkers. Therefore, given that  $\eta_{\text{H}_2\text{O}} = 1.01 \times 10^{-3}$  Pa s at 293 K, the range of  $\tau_{\text{DSE}}^{(r)}$  values for curcumin complexed in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur are 869–1189 and 869–998 ps, respectively. These  $\tau_{\text{DSE}}^{(r)}$  values are also in general agreement with the experimentally determined  $\tau^{(r)}$  values of  $>1200$  ps. These results indicate that the substantially longer  $\tau^{(r)}$  of 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin complexes than that of curcumin in MeOH is due to the significantly larger hydrodynamic volumes of these complexes.

## CONCLUSIONS

The excited-state dynamics of curcumin complexed by either 66 $\gamma$ CD<sub>2</sub>su or 66 $\gamma$ CD<sub>2</sub>ur in water are investigated for the first time, using femtosecond transient absorption spectroscopy. Comparisons are made with respect to *free* curcumin in methanol to gain insight into the effects of complexation in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur. Steady-state absorption and emission spectroscopic studies indicate significant reduction of water-curcumin interactions as a result of complexation in 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur. Both 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin in water only show an ESA signal at 530 nm without any SE signals, indicating the presence of significant non-radiative relaxation. The ESA dynamics of 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin are similar, having a rapid growth component and three decay components. The growth component of 0.25–0.41 ps is assigned to rapid reorganization of solvent molecules. The relatively fast decay components are attributed to relaxation due to ESIHT (9.3–21.8 ps). The small-amplitude and slow decay components are likely to be due to dynamics of complexed curcumin and molecular motions due to flexibility of the  $\gamma$ -CD moieties of 66 $\gamma$ CD<sub>2</sub>su and 66 $\gamma$ CD<sub>2</sub>ur. Finally, the transient absorption anisotropy results reveal that the slow rotational motions of 66 $\gamma$ CD<sub>2</sub>su-curcumin and 66 $\gamma$ CD<sub>2</sub>ur-curcumin in comparison with that of curcumin in MeOH are due to their large hydrodynamic volumes.

## AUTHOR INFORMATION

### Corresponding Author

\*E-mail: tak.kee@adelaide.edu.au. Tel: +61 (0)8 8313 5039. Fax: +61 (0)8 8313 4358.

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

This work was supported in part by research grants from the Australian Research Council (DP0878100, LE0989747, DP110103177, 20774028, and 20774030). The authors

acknowledge Ms. Mandy H. M. Leung and Mr. Scott N. Clifton at the Department of Chemistry, University of Adelaide, for assistance with the femtosecond transient absorption measurements.

## REFERENCES

- (1) Leung, M. H. M.; Harada, T.; Kee, T. W. Delivery of Curcumin and Medicinal Effects of the Copper(II)-Curcumin Complexes. *Curr. Pharm. Des.* **2013**, *19*, 2070–2083.
- (2) Goel, A.; Kunnumakkara, A. B.; Aggarwal, B. B. Curcumin as “Curecumin”: From Kitchen to Clinic. *Biochem. Pharmacol.* **2008**, *75*, 787–809.
- (3) Anand, P.; Thomas, S. G.; Kunnumakkara, A. B.; Sundaram, C.; Harikumar, K. B.; Sung, B.; Tharakan, S. T.; Misra, K.; Priyadarsini, I. K.; Rajasekharan, K. N.; et al. Biological Activities of Curcumin and Its Analogues (Congeners) Made by Man and Mother Nature. *Biochem. Pharmacol.* **2008**, *76*, 1590–1611.
- (4) Kiuchi, F.; Goto, Y.; Sugimoto, N.; Akao, N.; Kondo, K.; Tsuda, Y. Nematocidal Activity of Turmeric: Synergistic Action of Curcuminoids. *Chem. Pharm. Bull.* **1993**, *41*, 1640–1643.
- (5) Adhikary, R.; Barnes, C. A.; Trampel, R. L.; Wallace, S. J.; Kee, T. W.; Petrich, J. W. Photoinduced trans-to-cis Isomerization of Cyclocurcumin. *J. Phys. Chem. B* **2011**, *115*, 10707–10714.
- (6) Harada, T.; Giorgio, L.; Harris, T. J.; Pham, D.-T.; Ngo, H. T.; Need, E. F.; Coventry, B. J.; Lincoln, S. F.; Easton, C. J.; Buchanan, G.; et al. Diamide Linked  $\gamma$ -Cyclodextrin Dimers as Molecular-Scale Delivery Systems for the Medicinal Pigment Curcumin to Prostate Cancer Cells. *Mol. Pharmaceutics* **2013**, *10*, 4481–4490.
- (7) Aggarwal, B.; Kumar, A.; Bharti, A. Anticancer Potential of Curcumin: Preclinical and Clinical Studies. *Anticancer Res.* **2003**, *23*, 363–398.
- (8) Duvoix, A.; Blasius, R.; Delhalle, S.; Schnekenburger, M.; Morceau, F.; Henry, E.; Dicato, M.; Diederich, M. Chemopreventive and Therapeutic Effects of Curcumin. *Cancer Lett.* **2005**, *223*, 181–190.
- (9) Dorai, T.; Aggarwal, B. Role of Chemopreventive Agents in Cancer Therapy. *Cancer Lett.* **2004**, *215*, 129–140.
- (10) Mehta, K.; Pantazis, P.; McQueen, T.; Aggarwal, B. Antiproliferative Effect of Curcumin (Diferuloylmethane) against Human Breast Tumor Cell Lines. *Anticancer Drugs* **1997**, *8*, 470–481.
- (11) Ono, K.; Hasegawa, K.; Naiki, H.; Yamada, M. Curcumin Has Potent Anti-amyloidogenic Effects for Alzheimer's  $\beta$ -Amyloid Fibrils In Vitro. *J. Neurosci. Res.* **2004**, *75*, 742–750.
- (12) Baum, L.; Ng, A. Curcumin Interaction with Copper and Iron Suggests One Possible Mechanism of Action in Alzheimer's Disease Animal Models. *J. Alzheimer's Dis.* **2004**, *6*, 367–377.
- (13) Begum, A. N.; Jones, M. R.; Lim, G. P.; Morihara, T.; Kim, P.; Heath, D. D.; Rock, C. L.; Pruitt, M. A.; Yang, F.; Hudspeth, B.; et al. Curcumin Structure-Function, Bioavailability, and Efficacy in Models of Neuroinflammation and Alzheimer's Disease. *J. Pharmacol. Exp. Ther.* **2008**, *326*, 196–208.
- (14) Egan, M.; Pearson, M.; Weiner, S.; Rajendran, V.; Rubin, D.; Glockner-Pagel, J.; Canny, S.; Du, K.; Lukacs, G.; Caplan, M. Curcumin, a Major Constituent of Turmeric, Corrects Cystic Fibrosis Defects. *Science* **2004**, *304*, 600–602.
- (15) Cartiera, M. S.; Ferreira, E. C.; Caputo, C.; Egan, M. E.; Caplan, M. J.; Saltzman, W. M. Partial Correction of Cystic Fibrosis Defects with PLGA Nanoparticles Encapsulating Curcumin. *Mol. Pharmaceutics* **2010**, *7*, 86–93.
- (16) Lantz, R.; Chen, G.; Solyom, A.; Jolad, S.; Timmermann, B. The Effect of Turmeric Extracts on Inflammatory Mediator Production. *Phytomedicine* **2005**, *12*, 445–452.
- (17) Anand, P.; Kunnumakkara, A. B.; Newman, R. A.; Aggarwal, B. B. Bioavailability of Curcumin: Problems and Promises. *Mol. Pharmaceutics* **2007**, *4*, 807–818.
- (18) Letchford, K.; Liggins, R.; Burt, H. Pharmaceutics, Preformulation and Drug Delivery. *J. Pharm. Sci.* **2008**, *97*, 1179–1190.



- (19) Kaminaga, Y.; Nagatsu, A.; Akiyama, T.; Sugimoto, N.; Yamazaki, T.; Maitani, T.; Mizukami, H. Production of Unnatural Glucosides of Curcumin with Drastically Enhanced Water Solubility by Cell Suspension Cultures of *Catharanthus Roseus*. *FEBS Lett.* **2003**, *555*, 311–316.
- (20) Wang, Z.; Leung, M. H. M.; Kee, T. W.; English, D. S. The Role of Charge in the Surfactant-Assisted Stabilization of the Natural Product Curcumin. *Langmuir* **2010**, *26*, 5520–5526.
- (21) Leung, M. H. M.; Colangelo, H.; Kee, T. W. Encapsulation of Curcumin in Cationic Micelles Suppresses Alkaline Hydrolysis. *Langmuir* **2008**, *24*, 5672–5675.
- (22) Tønnesen, H. H. Solubility, Chemical and Photochemical Stability of Curcumin in Surfactant Solutions: Studies of Curcumin and Curcuminoids, XXVIII. *Pharmazie* **2002**, *57*, 820–824.
- (23) Banerjee, C.; Ghosh, S.; Mandal, S.; Kuchlyan, J.; Kundu, N.; Sarkar, N. Exploring the Photophysics of Curcumin in Zwitterionic Micellar System: An Approach to Control ESIPT Process in the Presence of Room Temperature Ionic Liquids (RTILs) and Anionic Surfactant. *J. Phys. Chem. B* **2014**, *118*, 3669–3681.
- (24) Leung, M. H. M.; Kee, T. W. Effective Stabilization of Curcumin by Association to Plasma Proteins: Human Serum Albumin and Fibrinogen. *Langmuir* **2009**, *25*, 5773–5777.
- (25) Sahu, A.; Kasoju, N.; Bora, U. Fluorescence Study of the Curcumin-Casein Micelle Complexation and Its Application as a Drug Nanocarrier to Cancer Cells. *Biomacromolecules* **2008**, *9*, 2905–2912.
- (26) Mohanty, C.; Acharya, S.; Mohanty, A. K.; Dilnawaz, F.; Sahoo, S. K. Curcumin-Encapsulated MePEG/PCL Diblock Copolymeric Micelles: A Novel Controlled Delivery Vehicle for Cancer Therapy. *Nanomedicine* **2010**, *5*, 433–449.
- (27) Shaikh, J.; Ankola, D. D.; Beniwal, V.; Singh, D.; Kumar, M. N. V. R. Nanoparticle Encapsulation Improves Oral Bioavailability of Curcumin by at Least 9-Fold When Compared to Curcumin Administered with Piperine as Absorption Enhancer. *Eur. J. Pharm. Sci.* **2009**, *37*, 223–230.
- (28) Harada, T.; Pham, D.-T.; Lincoln, S. F.; Kee, T. W. The Capture and Stabilization of Curcumin Using Hydrophobically Modified Polyacrylate Aggregates and Hydrogels. *J. Phys. Chem. B* **2014**, *118*, 9515–9523.
- (29) Harada, T.; Pham, D.-T.; Leung, M. H. M.; Ngo, H. T.; Lincoln, S. F.; Easton, C. J.; Kee, T. W. Cooperative Binding and Stabilization of the Medicinal Pigment Curcumin by Diamide Linked  $\gamma$ -Cyclodextrin Dimers: A Spectroscopic Characterization. *J. Phys. Chem. B* **2011**, *115*, 1268–1274.
- (30) Tønnesen, H.; Måsson, M.; Loftsson, T. Studies of Curcumin and Curcuminoids. XXVII. Cyclodextrin Complexation: Solubility, Chemical and Photochemical Stability. *Int. J. Pharm.* **2002**, *244*, 127–135.
- (31) Uekama, K.; Hirayama, F.; Irie, T. Cyclodextrin Drug Carrier Systems. *Chem. Rev.* **1998**, *98*, 2045–2076.
- (32) Davis, M.; Brewster, M. Cyclodextrin-Based Pharmaceuticals: Past, Present and Future. *Nat. Rev. Drug Discovery* **2004**, *3*, 1023–1035.
- (33) Szejtli, J. Introduction and General Overview of Cyclodextrin Chemistry. *Chem. Rev.* **1998**, *98*, 1743–1753.
- (34) Pham, D.-T.; Clements, P.; Easton, C. J.; Papageorgiou, J.; May, B. L.; Lincoln, S. F. Complexation of 6-(4'-(toluidinyl)naphthalene-2-sulfonate) by  $\beta$ -Cyclodextrin and Linked  $\beta$ -Cyclodextrin Dimers. *New J. Chem.* **2008**, *32*, 712–718.
- (35) Pham, D.-T.; Clements, P.; Easton, C. J.; Papageorgiou, J.; May, B. L.; Lincoln, S. F. Dimerisation and Complexation of 6-(4'-t-Butylphenylamino)naphthalene-2-sulphonate by  $\beta$ -Cyclodextrin and Linked  $\beta$ -Cyclodextrin Dimers. *Supramol. Chem.* **2009**, *21*, 510–519.
- (36) Pham, D.-T.; Ngo, H. T.; Lincoln, S. F.; May, B. L.; Easton, C. J. Synthesis of C6(A)-to-C6(A) and C3(A)-to-C3(A) Diamide Linked  $\gamma$ -Cyclodextrin Dimers. *Tetrahedron* **2010**, *66*, 2895–2898.
- (37) Chan, W.-H.; Wu, H.-J. Anti-Apoptotic Effects of Curcumin on Photosensitized Human Epidermal Carcinoma A431 Cells. *J. Cell. Biochem.* **2004**, *92*, 200–212.
- (38) Koon, H. K.; Leung, A. W. N.; Yue, K. K. M.; Mak, N. K. Photodynamic Effect of Curcumin on NPC/CNE2 Cells. *J. Environ. Pathol. Toxicol. Oncol.* **2006**, *25*, 205–216.
- (39) Park, K.; Lee, J.-H. Photosensitizer Effect of Curcumin on UVB-Irradiated HaCaT Cells through Activation of Caspase Pathways. *Oncol. Rep.* **2007**, *17*, 537–540.
- (40) Leung, M. H. M.; Pham, D.-T.; Lincoln, S. F.; Kee, T. W. Femtosecond Transient Absorption Spectroscopy of Copper(II)-Curcumin Complexes. *Phys. Chem. Chem. Phys.* **2012**, *14*, 13580–13587.
- (41) Adhikary, R.; Carlson, P. J.; Kee, T. W.; Petrich, J. W. Excited-State Intramolecular Hydrogen Atom Transfer of Curcumin in Surfactant Micelles. *J. Phys. Chem. B* **2010**, *114*, 2997–3004.
- (42) Adhikary, R.; Mukherjee, P.; Kee, T. W.; Petrich, J. W. Excited-State Intramolecular Hydrogen Atom Transfer and Solvation Dynamics of the Medicinal Pigment Curcumin. *J. Phys. Chem. B* **2009**, *113*, 5255–5261.
- (43) Nardo, L.; Paderno, R.; Andreoni, A.; Måsson, M.; Haukvik, T.; Tønnesen, H. H. Role of H-Bond Formation in the Photoreactivity of Curcumin. *Spectroscopy* **2008**, *22*, 187–198.
- (44) Nardo, L.; Andreoni, A.; Bondani, M.; Måsson, M.; Tønnesen, H. H. Studies on Curcumin and Curcuminoids. XXXIV. Photophysical Properties of a Symmetrical, Nonsubstituted Curcumin Analogue. *J. Photochem. Photobiol. B* **2009**, *97*, 77–86.
- (45) Mandal, S.; Ghosh, S.; Banik, D.; Banerjee, C.; Kuchlyan, J.; Sarkar, N. An Investigation into the Effect of the Structure of Bile Salt Aggregates on the Binding Interactions and ESIHT Dynamics of Curcumin: A Photophysical Approach to Probe Bile Salt Aggregates as a Potential Drug Carrier. *J. Phys. Chem. B* **2013**, *117*, 13795–13807.
- (46) Khopde, S. M.; Indira Priyadarsini, K.; Palit, D. K.; Mukherjee, T. Effect of Solvent on the Excited-State Photophysical Properties of Curcumin. *Photochem. Photobiol.* **2000**, *72*, 625–631.
- (47) Ghosh, R.; Mondal, J. A.; Palit, D. K. Ultrafast Dynamics of the Excited States of Curcumin in Solution. *J. Phys. Chem. B* **2010**, *114*, 12129–12143.
- (48) McMorrow, D.; Kasha, M. Intramolecular Excited-State Proton Transfer in 3-Hydroxyflavone. Hydrogen-Bonding Solvent Perturbations. *J. Phys. Chem.* **1984**, *88*, 2235–2243.
- (49) Schwartz, B. J.; Peteanu, L. A.; Harris, C. B. Direct Observation of Fast Proton Transfer: Femtosecond Photophysics of 3-Hydroxyflavone. *J. Phys. Chem.* **1992**, *96*, 3591–3598.
- (50) Mohammed, O. F.; Xiao, D.; Batista, V. S.; Nibbering, E. T. J. Excited-State Intramolecular Hydrogen Transfer (ESIHT) of 1,8-Dihydroxy-9,10-anthraquinone (DHAQ) Characterized by Ultrafast Electronic and Vibrational Spectroscopy and Computational Modeling. *J. Phys. Chem. A* **2014**, *118*, 3090–3099.
- (51) Moog, R. S.; Maroncelli, M. 7-Azaindole in Alcohols: Solvation Dynamics and Proton Transfer. *J. Phys. Chem.* **1991**, *95*, 10359–10369.
- (52) Barik, A.; Goel, N.; Priyadarsini, K.; Mohan, H. Effect of Deuterated Solvents on the Excited State Photophysical Properties of Curcumin. *J. Photosci.* **2004**, *11*, 95–99.
- (53) Vajda, S.; Jimenez, R.; Rosenthal, S.; Fidler, V.; Fleming, G.; Castner, E. Femtosecond to Nanosecond Solvation Dynamics in Pure Water and Inside the  $\gamma$ -Cyclodextrin Cavity. *J. Chem. Soc., Faraday Trans.* **1995**, *91*, 867–873.
- (54) Jimenez, R.; Fleming, G. R.; Kumar, P.; Maroncelli, M. Femtosecond Solvation Dynamics of Water. *Nature* **1994**, *369*, 471–473.
- (55) Wallace, S. J.; Kee, T. W.; Huang, D. M. Molecular Basis of Binding and Stability of Curcumin in Diamide-Linked  $\gamma$ -Cyclodextrin Dimers. *J. Phys. Chem. B* **2013**, *117*, 12375–12382.
- (56) Roy, D.; Mondal, S. K.; Sahu, K.; Ghosh, S.; Sen, P.; Bhattacharyya, K. Temperature Dependence of Anisotropy Decay and Solvation Dynamics of Coumarin 153 in  $\gamma$ -Cyclodextrin Aggregates. *J. Phys. Chem. A* **2005**, *109*, 7359–7364.
- (57) Sen, P.; Roy, D.; Mondal, S. K.; Sahu, K.; Ghosh, S.; Bhattacharyya, K. Fluorescence Anisotropy Decay and Solvation

Dynamics in a Nanocavity: Coumarin 153 in Methyl  $\beta$ -Cyclodextrins.

*J. Phys. Chem. A* **2005**, *109*, 9716–9722.

(58) Zsila, F.; Bikadi, Z.; Simonyi, M. Circular Dichroism Spectroscopic Studies Reveal pH Dependent Binding of Curcumin in the Minor Groove of Natural and Synthetic Nucleic Acids. *Org. Biomol. Chem.* **2004**, *2*, 2902–2910.