

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231667513>

Anisotropic Interactions in Protein Mixtures: Self Assembly and Phase Behavior in Aqueous Solution

ARTICLE *in* JOURNAL OF PHYSICAL CHEMISTRY LETTERS · FEBRUARY 2012

Impact Factor: 7.46 · DOI: 10.1021/jz201680m

CITATIONS

12

READS

22

5 AUTHORS, INCLUDING:



Anil Kurut

University of Copenhagen

7 PUBLICATIONS 44 CITATIONS

SEE PROFILE



Björn Persson

Lund University

9 PUBLICATIONS 119 CITATIONS

SEE PROFILE



Jan Forsman

Lund University

89 PUBLICATIONS 1,056 CITATIONS

SEE PROFILE

Anisotropic Interactions in Protein Mixtures: Self Assembly and Phase Behavior in Aqueous Solution

Anil Kurut,^{*} Björn A. Persson, Torbjörn Åkesson, Jan Forsman, and Mikael Lund^{*}

Department of Theoretical Chemistry, Lund University, P.O.B. 124 SE-22100 Lund, Sweden


E-mail: anil.kurut@teokem.lu.se; mikael.lund@teokem.lu.se

^{*}To whom correspondence should be addressed

Abstract

Recent experimental studies show that oppositely charged proteins can self-assemble to form seemingly stable micro-spheres in aqueous salt solutions. We here use parallel tempering Monte Carlo simulations to study protein phase separation of lysozyme/ α -lactalbumin mixtures and show that anisotropic electrostatic interactions are important for driving protein self-assembly. In both dilute and concentrated protein phases, the proteins strongly align according to their charge distribution. While this alignment can be greatly diminished by a *single* point mutation, phase separation is completely suppressed when neglecting electrostatic anisotropy. The results highlight the importance of subtle electrostatic interactions even in crowded biomolecular environments where other short-ranged forces are often thought to dominate.

TOC Graphic



Keywords: Protein-protein interactions, electrostatics, non-centrosymmetric interactions, micro sphere formation, whey proteins, coarse graining

In a set of experimental studies it has been shown that 1:1 mixtures of lysozyme (lys) and α -lactalbumin (α -lac) self-assemble into well defined micrometer sized spheres.¹⁻³ This process is highly dependent on the salt concentration and microspheres are formed only for the calcium depleted *apo* form of α -lac at salt concentrations below 100 mM, while for the calcium loaded *holo* form of α -lac the self assembly vanishes. Contrary to phase separation of similarly charged proteins,^{4,5} lys and apo α -lac phase separate due to attractive electrostatic interactions between the two oppositely charged proteins. Interestingly, this electrostatic interaction is far from isotropic and it has been shown that the anisotropic charge distribution of α -lac, combined with the high positive net charge of lys gives rise to strong directional protein-protein interactions.^{6,7} The remaining

question is how such long-ranged non-centrosymmetric interactions from charged residues influence protein assembly and phase transition? We here present a mesoscopic simulation study of aqueous protein mixtures where we investigate the contribution of anisotropic protein-protein interactions on phase transition of proteins as well as on micro-structuring in dense protein environments.

Previous studies on anisotropic protein-protein interactions⁸⁻¹⁴ typically describe proteins as patchy spherical particles with attractive sites interacting via angularly dependent square well potentials. These models rely mainly on ambiguously placed attractive patches and not on anisotropy owing to non-uniform surface charge distributions of real proteins. One exception is the embedded discreet charge model^{9,11,15} where charged residues are projected onto a hard sphere. Although approximately accounting for anisotropic electrostatics, this model treats the excluded volume as spherical symmetric. To remedy this we here develop a protein model that captures anisotropic interactions due to both excluded volume and electrostatic interactions while at the same time being fast enough for simulation studies of phase transitions involving *many* protein molecules.

Simulating phase behavior at an atomistic level is unfeasible due to the vast number of solvent and solute molecules in the system. We therefore gradually coarse grained the proteins – see illustration in Figure 1 – while maintaining (i) the salt dependent potential of mean force between lys/ α -lac and (ii) the distance dependent average alignment of α -lac’s dipole moment with respect to lys. The latter, unit less property ranges between ± 1 where +1 corresponds to a fully aligned dipole. The final model, (c), maintains all charged residues as in the crystal structure, while the remaining part is represented by a central particle. As shown in the supporting information this model closely reproduces the well tested reference model (a).

Metropolis Monte Carlo simulations in the canonical ensemble, NVT , were used for two-protein interactions when coarse graining model (a) while simulations in the isobaric-isothermal ensemble, NpT were used to study many-protein interactions and phase behavior with model (c). The system energy for a given configuration was $U = \sum_{i<j} 4\epsilon_L [(\frac{\sigma_{ij}}{r_{ij}})^{12} - (\frac{\sigma_{ij}}{r_{ij}})^6] + \frac{e^2 z_i z_j}{4\pi\epsilon_0\epsilon_r r_{ij}} \zeta$ where ϵ_L is the depth of the Lennard-Jones potential, σ_i the diameter, and $\sigma_{ij} = (\sigma_i + \sigma_j)/2$. r_{ij} is the




Figure 1: Coarse graining starting from an implicit solvent, explicit salt model where amino acids are represented by spheres (a).^{6,16} From (a)→(b) Debye-Hückel electrostatics; (b)→(c) neutral particles are collected into a central sphere. See supporting information for details.

separation between particle i and j , z_i the charge valency, e the electron unit charge, and $\epsilon_0\epsilon_r$ is the dielectric permittivity of water. For explicit salt, $\zeta = 1$; for implicit salt, $\zeta = e^{-\kappa r_{ij}}$ where κ is the inverse Debye screening length. For NpT simulations, the histogram method was used to sample volume probability distributions, $P(V)$, related to the constrained Gibbs free energy, $G = -kT \ln P(V)$.¹⁷ Minima in G correspond to (meta-)stable phases and to obtain coexistence concentrations, G was re-weighted, $G' = G \pm \Delta pV$ such that two minima had equal free energies.¹⁸ The Helmholtz free energy is $A = G' - pV$. We used parallel tempering in both p and κ to efficiently sample configurational space at low salt concentrations where strongly attached clusters form. For simulation details, see supporting information.

Table 1: Model details for lys and α -lac. For all particles $\epsilon_{LJ} = 0.05 kT$ in model (a) and (b), $\epsilon_{LJ} = 0.075 kT$ in model (c). $\sigma_i = 30 \text{ \AA}$ for the central sphere (see reference⁶ for σ of other particles).

Protein	α -lac				lys
	apo	holo	mutated ^a	smeared ^b	
PDB entry	1F6R	1F6S	1F6R	1F6R	4LZT
N_{residues}	124	125	124	124	130
Net charge, pH 7.5	-5.9	-4.5	-5.9	-5.9	7.0
Dipole moment ^c (D)	78	68	34	3	20

^a Residues D79 and K99 are swapped.

^b Net charge is smeared out over all titratable sites.

^c Calculated with respect to the protein mass center.

Using model (c) we performed NpT simulations with 40 proteins to investigate the Helmholtz free energy of bulk 1:1 protein mixtures containing lys and four different forms of α -lac: apo, holo, smeared, and mutated apo – see Table 1 and Figure 2. Phase separation – indicated by a common

tangent of two points in the Helmholtz free energy curve – was observed only for mixtures containing the apo forms and at low salt concentrations. This is in perfect agreement with experimental observations.² Increasing the salt concentration leads to reduced electrostatic interactions between the oppositely charged lys and α -lac, and eventually suppresses phase separation. The same is the case for holo α -lac, where the bound calcium ion reduces the net charge of α -lac.




Figure 2: Helmholtz free energy for 1:1 mixtures of lys with: apo, holo, mutated, smeared α -lac at 30 mM (solid lines) and 100 mM (dashed lines) salt concentrations.

To identify the contribution of anisotropic electrostatics to phase separation we artificially reduced the α -lac dipole moment from 78 to 34 Debye by swapping merely two charged amino acids (Table 1). As seen in Figure 3 this mutation – which does not alter the net charge – drastically decreased the critical salt concentration at coexistence and widens the immiscibility gap of lys/apo α -lac. We also approached isotropic electrostatics by smearing out the total charge of apo α -lac on *all* titratable residues and, as seen in Figure 2, lack of directional interactions completely obstructs phase separation in the investigated salt concentration range.

It is instructive to follow the alignment of α -lac's dipole moment with respect to the mass center of lys – see Figure 3, right: At infinite protein dilution the apo form of α -lac is aligned by more than 70% when in contact with lys and then levels off to zero when the proteins separate. As expected, by diminishing the dipole moment with a point mutation, this alignment is reduced. A similar behavior is seen in the dense protein phases but to a lesser degree due to the surrounding proteins. Still, the apo form of α -lac is aligned by $\sim 40\%$ implying that anisotropic electrostatics are involved in the dense phase structuring.

Experimentally, micro-sphere formation occurs at low salt and at total protein concentrations between 0.075–1.05 mM² implying that the equilibrium concentration of the dilute phase is in the sub milli-molar range. Our model predicts a dilute phase concentration 5 to 70 times higher, and thus underestimates the width of the coexistence curve. Similar issues have been reported for lysozyme.^{11,19} One reason could be lack of charge regulation in our model, although this is probably of minor importance.^{6,20} It has also been proposed that apo α -lac undergoes conformational changes into a semi-folded molten globule state upon binding to lysozyme, leading to exposure of hydrophobic residues.³ Such a mechanism – which will be temperature dependent^{8,19} – likely leads to a short-range attraction unaccounted for by our rigid protein model. Although quantitative agreement is not the chief goal of this work, we attempted to remedy the above discrepancy by pragmatically incorporating a square well potential – depth one kT , width 3.1 Å, corresponding to a water layer – between the neutral spheres in model (c). This spherically symmetric potential is justified by lack of angular correlations in the hydrophobic interactions for the present system – see supporting information. Figure 3 shows that this simple addition significantly improves the agreement with the experimental dilute phase density. Further, the simulated two-body potential of mean force, $w(R)$, is related to the thermodynamic dissociation constant, $K_d^{-1} \approx 4\pi \int_{\text{contact}}^{\infty} (e^{-w(R)/kT} - 1) R^2 dR$.²¹ Without short ranged attraction, $K_d = 737 \mu\text{M}$ at 39 mM salt and pH 7.5, merely 32% off the experimental value of $490 \pm 70 \mu\text{M}$.⁷ Including the above square well potential, we obtained $K_d = 446 \mu\text{M}$. Thus, the short ranged attraction improves not only agreement with the experimental phase behavior, but also consistency at the two-protein level. Note that our reference model (a) was originally parameterized to reproduce lys-lys osmotic second virial coefficients¹⁶ and that we have made no efforts to fit it to experimental data for the current lys/ α -lac system.

We have shown that anisotropic electrostatic interactions resulting from a non-uniform distribution of amino acid charges play an important role in protein self-assembly. Due to the high charge anisotropy of α -lac, this protein aligns in the electric field of lys – both in dilute and concentrated protein environments, showing that directional electrostatics are influential even at crowded condi-




Figure 3: *Left*: Fluid-fluid coexistence curve of lys and α -lac in different forms and with extra short ranged attraction (SR). The shaded area shows the experimentally observed concentration range of the dilute phase; error bars reflect the standard deviation of five independent MC runs. *Right*: α -lac dipole moment alignment with respect to lys (25 mM salt) as a function of protein separation in the dense phases (solid) and at infinite dilution (dotted); The vertical line indicates the position of the $w(r)$ minima.

tions. Phase separation is sensitive to subtle charge changes and a single amino acid point mutation that minimizes the protein dipole moment of α -lac – while retaining the net charge – is enough to significantly shift the immiscibility gap to lower salt concentrations. This dipole moment reduction changes the close contact alignment and, consequently, the structure of the dense phase. Artificially smearing out the charge over the protein surface – and thus approaching isotropic electrostatics – completely inhibits phase separation. These results suggest that phase behavior of aqueous protein mixtures is sensitive to anisotropic electrostatics – both qualitatively and quantitatively – and that inclusion of directionality in theoretical models is essential for accurate prediction of coexistence curves.

This work was supported by the Linnaeus center Organizing Molecular Matter and the eSSSENCE program at Lund University. Computational resources were provided by LUNARC.

Supporting Information Available

Simulation details and model development. This material is available free of charge via the Internet at <http://pubs.acs.org/>.

References

- (1) Nigen, M.; Croguennec, T.; Renard, D.; Bouhallab, S. Temperature Affects the Supramolecular Structures Resulting from α -Lactalbumin-Lysozyme Interaction. *Biochem.* **2007**, *46*, 1248–1255.
- (2) Nigen, M.; Croguennec, T.; Bouhallab, S. Formation and Stability of α -Lactalbumin-Lysozyme Spherical Particles: Involvement of Electrostatic Forces. *Food Hydrocolloids* **2009**, *23*, 510–518.
- (3) Nigen, M.; Le Tilly, V.; Croguennec, T.; Drouin-Kucma, D.; Bouhallab, S. Molecular Interaction between Apo or Holo α -lactalbumin and Lysozyme: Formation of Heterodimers as Assessed by Fluorescence Measurements. *Biochim. Biophys. Acta* **2009**, *1794*, 709–715.
- (4) Taratuta, V. G.; Holschbach, A.; Thurston, G. M.; Blankschtein, D.; Benedek, G. B. Liquid-Liquid Phase Separation of Aqueous Lysozyme Solutions: Effects of pH and Salt Identity. *J. Phys. Chem.* **1990**, *94*, 2140–2144.
- (5) Shukla, A.; Mylonas, E.; Di Cola, E.; Finet, S.; Timmins, P.; Narayanan, T.; Svergun, D. I. Absence of Equilibrium Cluster Phase in Concentrated Lysozyme Solutions. *PNAS* **2008**, *105*, 5075–80.
- (6) Persson, B. A.; Lund, M. Association and Electrostatic Steering of α -Lactalbumin and Lysozyme Heterodimers. *Phys. Chem. Chem. Phys.* **2009**, *11*, 8879–8885.
- (7) Salvatore, D. B.; Duraffourg, N.; Favier, A.; Persson, B. A.; Lund, M.; Delage, M.-M.; Silvers, R.; Schwalbe, H.; Croguennec, T.; Bouhallab, S.; Forge, V. Investigation at Residue Level of the Early Steps during the Assembly of Two Proteins into Supramolecular Objects. *Biomacromol.* **2011**, *12*, 2200–2210.
- (8) Lomakin, A.; Asherie, N.; Benedek, G. B. Alelotopic Interactions of Globular Proteins. *Proc. Natl. Acad. Sci.* **1999**, *96*, 9465–9468.

- (9) Carlsson, F.; Malmsten, M.; Linse, P. Monte Carlo Simulations of Lysozyme Self-Association in Aqueous Solution. *J. Phys. Chem. B* **2001**, *105*, 12189–12195.
- (10) Kern, N.; Frenkel, D. Fluid-Fluid Coexistence in Colloidal Systems with Short-Ranged Strongly Directional Attraction. *J. Chem. Phys.* **2003**, *118*, 9882–9889.
- (11) Rosch, T. W.; Errington, J. R. Investigation of the Phase Behavior of an Embedded Charge Protein Model through Molecular Simulation. *J. Phys. Chem. B* **2007**, *111*, 12591–12598.
- (12) Liu, H.; Kumar, S. K.; Sciortino, F. Vapor-Liquid Coexistence of Patchy Models: Relevance to Protein Phase Behavior. *J. Chem. Phys.* **2007**, *127*, 084902–1–084902–5.
- (13) Wentzel, N.; Gunton, J. D. Effect of Solvent on the Phase Diagram of a Simple Anisotropic Model of Globular Proteins. *J. Phys. Chem* **2008**, *112*, 7803–7809.
- (14) Gögelein, C.; Nägele, G.; Tuinier, R.; Gibaud, T.; Stradner, A.; Schurtenberger, P. A simple patchy colloid model for the phase behavior of lysozyme dispersions. *J. Chem. Phys.* **2008**, *129*, 085102–1–085102–12.
- (15) Abramo, M. C.; Caccamo, C.; Costa, D.; Pellicane, G.; Ruberto, R. Molecular Dynamics of an Embedded-Charge Model of Lysozyme Aqueous Solutions. *J. Phys. Chem. B* **2010**, *114*, 9109–9118.
- (16) Lund, M.; Jönsson, B. A Mesoscopic Model for Protein-Protein Interactions in Solution. *Biophys. J.* **2003**, *85*, 2940–2947.
- (17) Hunter, J. E.; Reinhardt, W. P. Finite-Size Scaling Behavior of The Free Energy Barrier between Coexisting Phases: Determination of the Critical Temperature and Interfacial Tension of the Lennard-Jones fluid. *J. Chem. Phys.* **1995**, *103*, 8627–8637.
- (18) Brukhno, A. V.; Åkesson, T.; Jönsson, B. Phase Behavior in Suspensions of Highly Charged Colloids. *J. Chem. Phys.* **2009**, *113*, 6766–6774.

- (19) Biesheuvel, P. M.; Lindhoud, S.; de Vries, R.; Cohen Stuart, M. A. Phase Behavior of Mixtures of Oppositely Charged Nanoparticles: Heterogeneous Poisson-Boltzmann Cell Model Applied to Lysozyme and Succinylated Lysozyme. *Langmuir* **2006**, 22, 1291–300.
- (20) Maarten Biesheuvel, P.; Lindhoud, S.; Cohen Stuart, M.; de Vries, R. Phase Behavior of Mixtures of Oppositely Charged Protein Nanoparticles at Asymmetric Charge Ratios. *Phys. Rev. E* **2006**, 73, 041408.
- (21) Wennerström, H. *H. Statistical Mechanical Description of Surfactant Self-assembly. In Organized Solutions: Surfactants in Science and Technology*; Marcel Dekker: New York, 1992.