


**Fig. 3** Application of Snatzke MO formalism to  $\Delta^4$ -uronates: (a) chiral  $\alpha$ -alkoxy acrylate chromophore (absolute conformation of the dehydropyran ring is chosen as  $^1H_2$ ); (b) relevant MOs of the isoelectronic 2,3-dimethylbutadiene dianion; (c) formal multiplication of  $\pi_b$  and  $\pi^*$  giving the electron density; (d) charge translation and  $\mu$ ; (e) concomitant charge rotation and  $m$



**Fig. 2** Superposition of CD curves with (a) negative (**1–5**) and (b) positive Cotton effects (**6** and **7**). These compounds show a single absorption peak at 235–240 nm.

within the dihydropyran ring and this is done for the absolute conformation  $^1H_2$  chosen on a left-handed helical path. Such rotation of charge along the line connecting C4–O<sub>ring</sub> generates  $m$  which, by the right-hand rule, is oriented antiparallel to  $\mu$ . The resulting Cotton effect is, therefore, negative and the absolute

conformation of the dihydropyran ring in compounds **1–5** is  $^1H_2$ .

For compounds **6** and **7** (a positive Cotton effect measured) the charge is transferred on a right-handed helical trajectory, thus predicting a  $^2H_1$  absolute conformation for their ring systems.

The  $^1H$  NMR data for the compounds are in full agreement with the prediction of their absolute conformation concluded from this new semi-empirical rule. CD spectroscopy provides the absolute conformation, information that is simply not obtainable using NMR analysis.

We thank Professors Dan Quinn and Lei Geng for their helpful suggestions and the National Institutes of Health (GM38060) for supporting this research.

## Notes and References

† E-mail: robert-linhardt@uiowa.edu

- H. G. Bazin, I. Capila and R. J. Linhardt, *Carbohydr. Res.*, 1998, in the press.
- B. M. Gimarc, *Acc. Chem. Res.*, 1974, **7**, 384.
- H. Bock, *Angew. Chem.*, 1977, **89**, 631.
- R. B. Woodward and R. Hoffmann, *The Conservation of Orbital Symmetry*, 1970, Verlag Chemie, Academic Press, Weinheim.
- F. Ciardelli and P. Salvadori, *Fundamental Aspects and Recent Developments in ORD and CD*, Heyden, London, 1973; M. Legrand and M. Rougier, in *Stereochemistry, Fundamentals and Methods*, ed. H. Kagan, Thieme, Stuttgart, 1977, vol. 2, p. 33.
- G. Snatzke, *Angew. Chem., Int. Ed. Engl.*, 1979, **18**, 363.
- G. Snatzke and H. Kagan, *Pure Appl. Chem.*, 1979, **51**, 769.
- A. Moscovitz, *Tetrahedron*, 1961, **13**, 48; A. Moscovitz, K. Mislow, A. Glass and C. Djerassi, *J. Am. Chem. Soc.*, 1961, **84**, 1945.
- A. Konoval, G. Snatzke and P. Thies, *Tetrahedron*, 1978, **34**, 253.

Received in Cambridge, UK, 27th April 1998; 8/03108H