See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/250811823

Symmetrical hydrogen bonds in the crystal structure of calcium bis(dihydrogen arsenate): a neutron-diffraction study

ARTICLE in JOURNAL OF THE CHEMICAL SOCIETY CHEMICAL COMMUNICATIONS · JANUARY 1971 DOI: 10.1039/c29710001566

CITATIONS READS 7

3 AUTHORS:



Giovanni Ferraris

Università degli Studi di Torino

343 PUBLICATIONS 3,868 CITATIONS

SEE PROFILE



Derry W. Jones

University of Bradford

185 PUBLICATIONS 1,590 CITATIONS

SEE PROFILE



J. Yerkess

University of Bradford

29 PUBLICATIONS 311 CITATIONS

SEE PROFILE

Symmetrical Hydrogen Bonds in the Crystal Structure of Calcium Bis(dihydrogen arsenate): a Neutron-diffraction Study

By G. Ferraris

(Istituto di Mineralogia e Geochimica dell' Università, 10123 Torino, Italy)

and D. W. Jones* and J. Yerkess

(School of Chemistry, University of Bradford, Bradford BD7 1DP)

Summary Neutron-diffraction analysis (final R-value 0.025 over 930 independent reflexions) of the crystal structure of calcium bis(dihydrogen arsenate), $\text{Ca}(\text{H}_2\text{AsO}_4)_2$, strongly indicates that inter-ion oxygen-oxygen separations of 2.436(3) and 2.444(3) Å contain symmetrical hydrogen bonds.

Confirmation by neutron-diffraction crystal-structure analysis of the occurrence of symmetrical hydrogen bonds between oxygen atoms of separate approximately tetrahedral oxyanions is uncommon. For the crystal structure of anhydrous calcium bis(dihydrogen arsenate), $Ca(H_2-AsO_4)_2$, which crystallises with two formula units in the triclinic space group $P\bar{1}$, single-crystal X-ray analysis¹ revealed two "very short"² inter-ion oxygen-oxygen contacts: $O(5) \cdots O(5') = 2 \cdot 454(8)$ and $O(8) \cdots O(8') = 2 \cdot 455(8)$ Å, both across symmetry centres. In order chiefly to investigate the symmetry or otherwise of the

corresponding hydrogen bonds, an accurate neutron-diffraction analysis of this structure has been carried out. All hydrogen atoms were located unequivocally by Fourier difference syntheses; the positions of all atoms were refined anisotropically by least squares to a final disagreement index, R, of 0.025 over 930 independent reflexions. Assignment of the space group to $P\bar{1}$ rather than P1 is supported by the low R-value, the occurrence of a single Fourier-difference peak of acceptable shape, the absence of thermal-parameter anomalies in the least-squares refinement, and the absence of a measured piezoelectric effect. Oxygen-oxygen separations of $O(5) \cdots O(5') = 2.436(3)$ and $O(8) \cdots O(8') = 2.444(3)$ Å are very close to those derived from the X-ray analysis; O(5)-H(4) = 1.218(3) and O(8)-H(5) = 1.222(3) Å.

In the two formally symmetrical hydrogen bonds, two kinds of consideration point to centring of the hydrogens, rather than statistical occupation of a pair of sites³ separated by 0.2 Å or so. First, the thermal parameters of H(4) and H(5) are close to those of O(5) and O(8), rather than being higher than those of nearby atoms. Secondly, for inter-anion hydrogen bonds in a series of acid hydrogen arsenates in which hydrogen positions have been determined directly by neutron diffraction (CaHAsO₄,2H₂O;⁴ Na₂HAsO₄,7H₂O;⁵ and CaHAsO₄,H₂O⁶), we have examined the variation of several quantities with O···H distance. For the most linear of these graphs, that of the $O-H \cdots O$ angle, the linearity is accurately preserved on the assumption that the $O(5) \cdots O(5')$ and $O(8) \cdots O(8')$ hydrogen bonds in Ca- $(H_2AsO_4)_2$ are genuinely symmetrical.

(Received, October 19th, 1971; Com. 1827.)

G. Chiari and G. Ferraris, Atti Acc. Sci. (Torino), 1971, 105, 725.
J. C. Speakman, Chem. Comm., 1967, 32.
W. C. Hamilton and J. A. Ibers, 'Hydrogen Bonding in Solids', Benjamin, New York, 1968.
G. Ferraris, D. W. Jones, and J. Yerkess, Acta Cryst., 1971, B27, 349.
G. Ferraris, D. W. Jones, and J. Yerkess, Acta Cryst., 1971, B27, 354.
G. Ferraris, D. W. Jones, and J. Yerkess, Acta Cryst., 1972, B28, in the press.