

# Mechanism Investigations of the Endo Cycloisomerization of Alkynols through Isolation and Characterization of Ruthenium Complexes from the Reactions of Alkynes with a Ruthenium Complex


Pei Nian Liu,<sup>\*,†</sup> Ting Bin Wen,<sup>‡</sup> Kun Dong Ju,<sup>†</sup> Herman H.-Y. Sung,<sup>‡</sup> Ian D. Williams,<sup>‡</sup> and Guochen Jia<sup>\*,†</sup>

<sup>†</sup>Key Laboratory for Advanced Materials and Institute of Fine Chemicals, East China University of Science and Technology, 130 Meilong Road, Shanghai, 200237, People's Republic of China

<sup>‡</sup>Department of Chemistry, Hong Kong University of Science & Technology, Clear Water Bay, Kowloon, Hong Kong, People's Republic of China

## Supporting Information

**ABSTRACT:** In our efforts to probe the reaction mechanism of the endo cycloisomerization of alkynols catalyzed by the complex  $[\text{Ru}(\text{N}_3\text{P})(\text{OAc})](\text{BPh}_4)$  (**1**;  $\text{N}_3\text{P} = N,N$ -bis[(pyridin-2-yl)methyl][2-(diphenylphosphino)phenyl]methanamine), the reactions of **1** with alkynes were investigated. Several complexes related to intermediates in the catalytic reactions were isolated and characterized. In the presence of DIPEA ( $N,N$ -diisopropylethanamine), complex **1** reacted with 3-butyne-1-ol to afford the Ru–oxocyclocarbene complex  $\{\text{Ru}(\text{N}_3\text{P})[\text{C}(\text{CH}_2)_3\text{O}](\text{OAc})\}(\text{BPh}_4)$  (**4**). Under anhydrous conditions, the reaction of complex **1** with *tert*-butylacetylene afforded  $[\text{Ru}(\text{N}_3\text{P})(\eta^3\text{-}^t\text{BuC}\equiv\text{CC}=\text{CHBu}^t)](\text{BPh}_4)$  (**8**). In the presence of water, the reaction of complex **1** with *tert*-butylacetylene afforded  $[\text{Ru}(\text{N}_3\text{P})(\text{CO})(\eta^1\text{-CH}_2\text{Bu}^t)](\text{BPh}_4)$  (**13**). All the reactions are likely to proceed through ruthenium vinylidene intermediates. These results support that ruthenium vinylidene complexes are involved as the key intermediates in the cycloisomerization of alkynols catalyzed by complex **1**.



## INTRODUCTION

Endocyclic enol ethers are very useful synthetic starting materials for the construction of diverse oxygen-containing heterocycles which are important structural units in a number of naturally occurring and biologically active molecules.<sup>1–6</sup> Catalytic endo cycloisomerization of alkynols represents the most straightforward approach to obtain endocyclic enol ethers,<sup>7</sup> with the advantages of high efficiency and 100% atom economy.<sup>8</sup> In the pioneering work in this area, McDonald and his co-workers discovered that molybdenum<sup>4,5,9</sup> and tungsten<sup>2,3,8–10</sup> carbonyl complexes can effectively catalyze the endo cycloisomerization of a range of alkynols. Such catalytic endo-cycloisomerization of alkynols has been successfully applied to the synthesis of various natural products,<sup>2</sup> glycals,<sup>3</sup> nucleosides,<sup>4</sup> and clinically valuable drugs.<sup>5</sup> Endo cycloisomerization of alkynols catalyzed by Mo and W complexes was generally believed to involve initial formation of a vinylidene intermediate, and the following intramolecular nucleophilic addition of the OH group to the vinylidene ligand would generate an endocyclic enol ether linked to the transition metal. Then protonation of the metal–carbon bond would afford the product endocyclic enol ether.<sup>11</sup> The mechanism is supported by DFT studies,<sup>12</sup> although experimental evidence for the mechanism is still scarce.<sup>13</sup>

In the past several years, other systems such as  $\text{CpRuL}_n^+ / 20\text{--}40\text{ mol } \%$  phosphine ligands,<sup>14</sup>  $[\text{Rh}(\text{PR}_3)_3\text{Cl}] / 30\text{--}55\text{ mol } \%$  phosphine ligands,<sup>15</sup>  $\text{AgNO}_3$  (10 mol %),  $\text{Pd}(\text{II})/\text{Cu}(\text{I})$ ,  $\text{Au}(\text{I})$  (17–30 mol %),<sup>16</sup> and  $\text{Pd}(\text{MeCN})_2\text{Cl}_2$ <sup>17</sup> were also found to be

catalytically active for the endo cycloisomerization of alkynols. Endo cycloisomerization of alkynols catalyzed by the Rh and Ru systems was proposed to proceed through vinylidene intermediates, while those catalyzed by  $\text{Ag}(\text{I})$ ,  $\text{Au}(\text{I})$ ,  $\text{Cu}(\text{I})$ , and  $\text{Pd}(\text{II})$  complexes were proposed to proceed through  $\eta^2$ -alkyne complex intermediates.

Ruthenium complexes containing a tripodal tetradentate ligand are appealing in catalysis, since the coordination of ruthenium with a tetradentate ligand offers rigid frameworks and two mutually cis coordination sites for substrates in an octahedral mononuclear complex.<sup>18</sup> Moreover, high structural stability can be achieved due to the chelate effect,<sup>19</sup> which offers a good opportunity to investigate catalytic reaction mechanisms through isolating relatively stable intermediates.<sup>20</sup>


Recently, we have reported the preparation of the new ruthenium complex  $[\text{Ru}(\text{N}_3\text{P})(\text{OAc})](\text{BPh}_4)$  (**1**;  $\text{N}_3\text{P} = N,N$ -bis[(pyridin-2-yl)methyl][2-(diphenylphosphino)phenyl]methanamine) and its catalytic properties for the cycloisomerization of a variety of alkynols with  $\text{C}\equiv\text{C}$  attached to aryl or alkyl groups (Scheme 1).<sup>21</sup> The reactions catalyzed by complex **1** exhibit high regioselectivity and reactivity to give exclusively endo products in good to excellent yields without any additive. Herein, we report our results in the investigation of the mechanism of the catalytic reactions through isolation and characterization of several ruthenium complexes related to the

Received: February 22, 2011

Published: April 05, 2011



### Scheme 1. Endo Cycloisomerization of Alkynols Catalyzed by Complex 1



reaction intermediates in the endo cycloisomerization of alkynols catalyzed by complex 1.

## RESULTS AND DISCUSSION

**Reaction of Complex 1 with 3-Butyn-1-ol.** Because ruthenium(II) complexes can react with terminal alkynes to generate vinylidene complexes<sup>22</sup> which can be attacked by weak nucleophiles such as alcohols,<sup>11,23</sup> water,<sup>24</sup> and carboxylic acids;<sup>25</sup> it seems reasonable to assume that the endo cycloisomerization of alkynols catalyzed by complex 1 proceeds through a vinylidene intermediate. A plausible mechanism of the catalytic reactions mediated by complex 1 is depicted in path I of Scheme 2 using 3-butyne-1-ol (2) as an example of the substrate. The alkynol 2 can react with complex 1 to give initially the  $\eta^2$ -alkyne complex A, which could then rearrange to the vinylidene intermediate B. Intramolecular attack by the hydroxyl group on the  $\alpha$ -carbon of the vinylidene ligand in B followed by deprotonation would afford the vinyl complex C. Protonation of the metal–carbon bond in C would then give the endocyclic enol ether 3 and regenerate complex 1. As reactions involving direct nucleophilic attack on  $\eta^2$ -alkyne rather than vinylidene ligands have also been suggested for some of the related transformations, for example, in addition reactions of amides to alkynes<sup>26</sup> and in addition reactions of carboxylic acids to terminal alkynes,<sup>27</sup> it is also possible that C can be formed from alkyne complex A without the involvement of vinylidene intermediate B.

In our initial attempts to probe the mechanism of the endo cycloisomerization of alkynols catalyzed by complex 1, the reactions of complex 1 with stoichiometric amounts of various alkynols were carried out. Unfortunately, we have failed to isolate or identify experimentally the active intermediates involved in the catalytic reactions. In an effort to gain indirect evidence for the involvement of the vinyl intermediate C in the catalytic reactions, the reaction of complex 1 with 3-butyne-1-ol (2) was carried out in the presence of *N,N*-diisopropylethanamine (DIPEA), which may neutralize the proton generated in the formation of C and thus quench the reaction of C to 1 and 3.

The reaction of complex 1 with alkynol 2 in the presence of DIPEA in refluxing THF for 1 h afforded the new Ru–oxacyclocarbene complex  $[\text{Ru}(\text{N}_3\text{P})(=\text{C}_4\text{H}_6\text{O})(\text{OAc})](\text{BPh}_4)$  (4) (Scheme 3). Several reactions of ruthenium complexes with alkynols to give oxacyclocarbene complexes have been reported.<sup>28</sup>

The Ru–oxacyclocarbene complex 4 has been characterized by  $^1\text{H}$ ,  $^{13}\text{C}\{^1\text{H}\}$ , and  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy, 2D-NMR experiments, including H,H-COSY, HSQC, and HMQC, and mass spectrometry. Consistent with the structure shown in Scheme 3, the  $^1\text{H}$  NMR spectrum of complex 4 (in  $\text{CD}_2\text{Cl}_2$ ) displays the methyl signal of the acetyl group at 1.97 ppm. With the help of 2D-NMR experiments, the signals of the six protons of the five-membered tetrahydrofuranlydene are located in the  $^1\text{H}$  NMR spectrum as a quartet at 4.25 ppm and a multiplet at 3.95–4.01 ppm for  $\text{C}^1\text{H}_2$ , two multiplets at 1.81–1.90 and 1.66–1.75 ppm for  $\text{C}^2\text{H}_2$ , and two multiplets at 1.92–2.04 and 3.65–3.72 ppm for  $\text{C}^3\text{H}_2$ . In the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum of complex 4, the carbon signals of the acetyl group appears at 25.1 and 177.2 ppm. The four carbon atoms of the five-membered oxaring show three singlets at 81.7 ( $\text{C}^1$ ), 22.5 ( $\text{C}^2$ ), and 53.2 ( $\text{C}^3$ ) ppm and a doublet at 312.7 ppm ( $\text{C}^4$ ,  $J_{\text{CP}} = 14$  Hz), which indicates the presence of  $\text{Ru}=\text{C}$ .<sup>28</sup> Consistent with this structure, the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum displays a singlet at 53.4 ppm and the HRMS spectrum shows the parent peak of the complex cation of 4 at  $m/z$  704.1570 (calcd 704.1611).


The structure of 4 has been confirmed by an X-ray diffraction study. Yellow single crystals of complex 4 suitable for X-ray diffraction analysis were obtained via slow diffusion of  $\text{Et}_2\text{O}$  into a  $\text{CH}_2\text{Cl}_2$  solution of complex 4. The crystal data and refinement details are given in Table 1, and selected bond lengths and angles are given in Table 2. An ORTEP view of the cation of complex 4 is shown in Figure 1. The coordination geometry around ruthenium in 4 can be viewed as a distorted octahedron with a tetradentate  $\text{N}_3\text{P}$  ligand, an acetyl ligand, and an oxocarbene ligand. The tertiary amine N atom is trans to the acetyl ligand. The two pyridyl units in 4 are cis to each other. The P atom is trans to a pyridyl N atom, and the carbene ligand is trans to the other pyridyl N atom. The  $\text{Ru}=\text{C}$  bond length ( $\text{Ru}(1)–\text{C}(3) = 1.906(4)$  Å) is similar to those of other Ru–alkoxycarbene complexes.<sup>28</sup> The solid-state structure is in agreement with the solution NMR spectroscopic data.

A plausible mechanism for the formation of complex 4 from complex 1 and alkynol 2 is shown in path II of Scheme 2. Complex 1 could coordinate alkyne 2, followed by rearrangement to give the vinylidene intermediate B. Subsequent intramolecular addition of the hydroxyl group to the  $\alpha$ -carbon of the vinylidene ligand in B could generate intermediate C, accompanied by the loss of a proton which could be neutralized by DIPEA. The reaction of C to form 1 and 3 could be quenched because protonation of the metal–carbon bond in C could not proceed due to the low acidity of  $\text{DIPEA-H}^+$ . On the other hand, complex C could be protonated to give complex 4.


In the cycloisomerization of alkynol 2 catalyzed by complex 1 without DIPEA, complex 4 could be detected by  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy. Moreover, complex 4 is also catalytically active in the cycloisomerization of alkynol. For example, when a THF solution of 4-pentyn-1-ol (5) was refluxed in the presence of 1 mol % complex 4 for 5 h, 100% conversion of 5 to product 6 was achieved (Scheme 4). This result demonstrates that complex 4 could be converted to intermediate C under the conditions for catalytic cycloisomerization of alkynol.

**Reaction of Complex 1 with *tert*-Butylacetylene (7) in the Absence of Water.** In the proposed mechanism for endo cycloisomerization of alkynols catalyzed by complex 1 (Scheme 2, path I) and for the formation of complex 4 from the reaction of 1 with 3-butyne-1-ol (2) (Scheme 2, path II), vinylidene complexes have been suggested as important intermediates. To gain more

**Scheme 2.** Proposed Mechanisms of the Complex 1 Catalyzed Endo Cycloisomerization of Alkynols and the Formation of Complex 4



**Scheme 3.** Reaction of Complex 1 with Alkynol 2 in the Presence of DIPEA To Afford Complex 4



evidence for the involvement of vinylidene intermediates, the reaction of complex 1 with *tert*-butylacetylene (7) was performed. When a THF solution containing complex 1 and 7 was refluxed for 1 h, complex 1 disappeared completely and the new butenyne ruthenium complex  $[\text{Ru}(\text{N}_3\text{P})(\eta^3\text{-}^t\text{BuC}_3\text{CHBu}^t)]\text{-}(\text{BPh}_4)$  (**8**) was produced, which can be isolated in 77% yield (Scheme 5). Related reactions of ruthenium complexes with alkynes to afford butenyne ruthenium complexes have been reported.<sup>29</sup> Butenyne ruthenium complexes have also been suggested as intermediates in the catalytic dimerization of terminal acetylenes.<sup>18e,30</sup>

The structure of complex 8 has been unambiguously confirmed by single-crystal X-ray analysis as well as its spectroscopic data. The  $^1\text{H}$  NMR spectrum of complex 7 showed  $^t\text{Bu}$  resonances at 0.76 ( $\text{H}^1$ ) and 1.29 ppm ( $\text{H}^8$ ) and the resonance of proton  $\text{H}^3$  at 4.80 ppm. The  $^{13}\text{C}\{^1\text{H}\}$  spectrum exhibits the resonances of tertiary carbons of  $^t\text{Bu}$  at 35.3 ( $\text{C}^2$ ) and 33.7 ( $\text{C}^7$ ) ppm and the resonances of methyl groups at 29.7 ( $\text{C}^1$ ) and 32.5 ( $\text{C}^8$ ) ppm. The resonance of the double-bonded carbon tethered with a proton was observed at 133.6 ppm. The resonances of  $\text{C}^4$ ,  $\text{C}^5$ , and  $\text{C}^6$  were found at 144.7, 48.9, and 123.7 ppm, respectively. The mass spectrum of complex 8 showed a strong signal of complex cation of 8 with  $m/z$  738.4191 (calcd 738.2546).

Yellow single crystals of complex 8 suitable for single-crystal X-ray diffraction analysis were obtained at 4 °C by slow diffusion of  $\text{Et}_2\text{O}$  into a  $\text{CH}_2\text{Cl}_2$  solution containing complex 8. The crystal data and refinement details are given in Table 1, and selected bond lengths and angles are given in Table 3. Figure 2 shows the X-ray structure of complex 8. The coordination geometry

around ruthenium in 8 is approximately octahedral, and the compound contains a tetradentate  $\text{N}_3\text{P}$  ligand and an  $\eta^3\text{-Bu}^t\text{C}_3\text{CHBu}^t$  ligand. The  $\eta^3\text{-Bu}^t\text{C}_3\text{CHBu}^t$  ligand is trans to the tertiary amine N atom and a pyridine N atom and the P atom is trans to another pyridine N atom. The bond lengths of  $\text{Ru}(1)\text{-C}(35)$  (2.212(5) Å),  $\text{Ru}(1)\text{-C}(36)$  (2.139(5) Å), and  $\text{Ru}(1)\text{-C}(37)$  (2.074(6) Å) and the bond angles of  $\text{C}(36)\text{-C}(35)\text{-C}(31)$  (142.7(6)°),  $\text{C}(35)\text{-C}(36)\text{-C}(37)$  (143.8(5)°),  $\text{C}(36)\text{-C}(37)\text{-C}(38)$  (139.3(6)°), and  $\text{C}(37)\text{-C}(38)\text{-C}(39)$  (127.4(6)°) are similar to those of other  $\text{Ru}-\eta^3\text{-RC}_3\text{CHR}'$  complexes.<sup>29</sup> The solid-state structure is in agreement with the solution NMR spectroscopic data.

A plausible mechanism for the reaction of complex 1 with *tert*-butylacetylene to afford complex 8 is depicted in Scheme 6.<sup>31</sup> First, complex 1 could react with acetylene 7 to generate the  $\eta^2$ -alkyne complex 9, which would rearrange to the vinylidene intermediate 10. One more molecule of acetylene 7 might react with 10 to form intermediate 11, which could undergo an intramolecular C–C coupling reaction to afford the 16-electron ruthenium complex 12, which could evolve to complex 8 after coordination of ruthenium with the  $\text{C}\equiv\text{C}$  moiety.

**Reaction of 1 with *tert*-Butylacetylene in the Presence of  $\text{H}_2\text{O}$ .** In our previous effort to gain indirect evidence for the involvement of the transformation in the catalytic reactions, we have carried out the reaction of complex 1 with phenylacetylene in the presence of  $\text{H}_2\text{O}$ . The reaction was found to give a benzyl carbonyl complex, which is most likely formed by nucleophilic attack of water at a vinylidene intermediate followed by an deinsertion reaction. Several related metal-assisted  $\text{C}\equiv\text{C}$  bond

Table 1. Crystal Data and Structure Refinement Details for Complexes 4, 8, 13, and 19


	4 · 1.5CH <sub>2</sub> Cl <sub>2</sub>	8 · CH <sub>2</sub> Cl <sub>2</sub>	13 · CH <sub>2</sub> Cl <sub>2</sub> · 0.25(hexane)	17 · 4THF
empirical formula	C <sub>62.5</sub> H <sub>60</sub> BCl <sub>3</sub> N <sub>3</sub> O <sub>3</sub> PRu	C <sub>68</sub> H <sub>69</sub> BCl <sub>2</sub> N <sub>3</sub> PRu	C <sub>63.5</sub> H <sub>64.5</sub> BCl <sub>2</sub> N <sub>3</sub> OPRu	C <sub>126</sub> H <sub>130</sub> B <sub>2</sub> N <sub>6</sub> O <sub>6</sub> P <sub>2</sub> Ru <sub>2</sub>
formula wt	1150.33	1142.01	1099.43	2110.06
temp (K)	173(2)	173(2)	173(2)	173(2)
wavelength (Å)	0.710 73	1.541 78	1.541 78	1.541 78
cryst syst	monoclinic	monoclinic	monoclinic	triclinic
space group	C2/c	P2 <sub>1</sub> /c	C2/c	P1
unit cell dims				
<i>a</i> (Å)	33.168(3)	22.9187(16)	33.8739(9)	14.3083(7)
<i>b</i> (Å)	9.0552(8)	13.7663(3)	12.3263(4)	15.0767(7)
<i>c</i> (Å)	36.835(3)	21.4402(10)	27.7974(6)	15.4095(7)
α (deg)	90	90	90	112.824(5)
β (deg)	92.1830(10)	117.178(7)	93.945(2)	104.771(4)
γ (deg)	90	90	90	107.571(4)
<i>V</i> (Å <sup>3</sup> )	11055.1(16)	6017.6(5)	11579.0(5)	2646.2(2)
<i>Z</i>	8	4	8	1
calcd density (Mg/m <sup>3</sup> )	1.382	1.261	1.261	1.324
abs coeff (mm <sup>−1</sup> )	0.507	3.495	3.626	3.065
<i>F</i> (000)	4760	2384	4580	1104
θ range for data collec (deg)	1.68–27.50	4.68–66.99	4.92–66.99	5.92–71.55
no. of rflns collected	32 951	20 042	16 943	13 956
no. of indep rflns	12 229 ( <i>R</i> (int) = 0.0569)	9764 ( <i>R</i> (int) = 0.0984)	9833 ( <i>R</i> (int) = 0.0546)	9512 ( <i>R</i> (int) = 0.0237)
completeness to θ = 25.00° (%)	97.2	91.1	95.2	94.9
max, min transmissn	1.00, 0.83	1.00, 0.58	1.00, 0.73	1.00, 0.81
no. of data/restraints/params	12 229/1/670	9764/0/690	9833/4/664	9512/0/649
goodness of fit on <i>F</i> <sup>2</sup>	1.040	1.036	1.012	1.055
final <i>R</i> indices ( <i>I</i> > 2σ( <i>I</i> ))	<i>R</i> 1 = 0.0650, <i>wR</i> 2 = 0.1366	<i>R</i> 1 = 0.0694, <i>wR</i> 2 = 0.1701	<i>R</i> 1 = 0.0548, <i>wR</i> 2 = 0.1233	<i>R</i> 1 = 0.0317, <i>wR</i> 2 = 0.0778
largest diff peak, hole (e Å <sup>−3</sup> )	0.978, −1.107	1.553, −1.121	1.458, −1.556	0.893, −1.024

Table 2. Selected Bond Lengths (Å) and Angles (deg) for Complex 4


Bond Distances			
Ru(1)–C(3)	1.906(4)	Ru(1)–O(1)	2.070(3)
Ru(1)–N(1)	2.114(3)	Ru(1)–N(10)	2.116(3)
Ru(1)–N(20)	2.175(3)	Ru(1)–P(1)	2.2843(10)
C(3)–C(4)	1.510(5)	C(4)–C(5)	1.520(6)
C(5)–C(6)	1.502(7)	C(6)–O(3)	1.478(5)
C(3)–O(3)	1.328(4)		
Bond Angles			
C(3)–Ru(1)–O(1)	98.20(13)	C(3)–Ru(1)–N(1)	94.41(13)
O(1)–Ru(1)–N(1)	161.56(11)	C(3)–Ru(1)–N(10)	88.66(14)
O(1)–Ru(1)–N(10)	87.72(12)	N(1)–Ru(1)–N(10)	79.16(12)
C(3)–Ru(1)–N(20)	170.18(13)	O(1)–Ru(1)–N(20)	84.65(11)
N(1)–Ru(1)–N(20)	80.78(12)	N(10)–Ru(1)–N(20)	82.05(12)
C(3)–Ru(1)–P(1)	93.42(11)	O(1)–Ru(1)–P(1)	98.27(8)
N(1)–Ru(1)–P(1)	94.32(8)	N(10)–Ru(1)–P(1)	173.30(9)
N(20)–Ru(1)–P(1)	95.47(9)	O(3)–C(3)–Ru(1)	125.8(3)
C(4)–C(3)–Ru(1)	126.5(3)	C(3)–C(4)–C(5)	104.3(3)
C(6)–C(5)–C(4)	102.9(4)	O(3)–C(6)–C(5)	103.9(3)
C(3)–O(3)–C(6)	113.0(3)	O(3)–C(3)–C(4)	107.4(3)

cleavage reactions of 1-alkynes with water have been reported to afford complexes containing a carbonyl and an  $\eta^1$ -alkyl with one fewer carbon atom, especially for metal complexes of ruthenium,<sup>32</sup> osmium,<sup>33</sup> and iridium.<sup>34</sup> The related reactions of metal acetylides or vinylidenes with water have been also reported.<sup>35</sup>

As complex 1 can also catalyze the endo cycloisomerization of alkynols with C≡C attached to aliphatic chains, it would be of interest to see if a similar reaction will also occur for the reaction of 1 with aliphatic alkynes in the presence of water. For this purpose, the reaction of complex 1 with *tert*-butylacetylene in the presence of H<sub>2</sub>O was carried out. After refluxing in THF for 8 h,

Figure 1. ORTEP diagram of the complex cation 4<sup>+</sup>.

#### Scheme 4. Complex 4 Catalyzed Endo Cycloisomerization of Alkynol 5 To Afford 6



the new complex 13 was produced and was isolated in 87% yield (Scheme 7).






Scheme 5. Reaction of Complex 1 and *tert*-Butylacetylene (7) To Afford Complex 8

Table 3. Selected Bond Lengths (Å) and Angles (deg) for Complex 8

Bond Distances			
Ru(1)–C(37)	2.074(6)	Ru(1)–C(36)	2.139(5)
Ru(1)–C(35)	2.212(5)	Ru(1)–N(1)	2.156(5)
Ru(1)–N(2)	2.139(5)	Ru(1)–N(3)	2.172(4)
Ru(1)–P(1)	2.2658(15)	C(31)–C(35)	1.493(8)
C(35)–C(36)	1.263(8)	C(36)–C(37)	1.392(8)
C(37)–C(38)	1.335(8)	C(38)–C(39)	1.510(8)
Bond Angles			
C(37)–Ru(1)–C(36)	38.6(2)	C(37)–Ru(1)–N(2)	82.3(2)
C(36)–Ru(1)–N(2)	85.5(2)	C(37)–Ru(1)–N(1)	171.8(2)
C(36)–Ru(1)–N(1)	138.00(19)	N(2)–Ru(1)–N(1)	90.16(18)
C(37)–Ru(1)–N(3)	102.9(2)	C(36)–Ru(1)–N(3)	140.0(2)
N(2)–Ru(1)–N(3)	76.62(17)	N(1)–Ru(1)–N(3)	78.28(17)
C(37)–Ru(1)–C(35)	72.1(2)	C(36)–Ru(1)–C(35)	33.7(2)
N(2)–Ru(1)–C(35)	94.1(2)	N(1)–Ru(1)–C(35)	105.5(2)
N(3)–Ru(1)–C(35)	170.07(19)	C(37)–Ru(1)–P(1)	88.26(17)
C(36)–Ru(1)–P(1)	95.05(16)	N(2)–Ru(1)–P(1)	164.03(12)
N(1)–Ru(1)–P(1)	99.84(13)	N(3)–Ru(1)–P(1)	93.15(13)
C(35)–Ru(1)–P(1)	95.21(16)	C(36)–C(35)–Ru(1)	70.0(4)
C(31)–C(35)–Ru(1)	147.3(5)	C(35)–C(36)–Ru(1)	76.3(4)
C(37)–C(36)–Ru(1)	68.2(3)	C(38)–C(37)–Ru(1)	147.2(5)
C(36)–C(37)–Ru(1)	73.2(3)	C(36)–C(35)–C(31)	142.7(6)
C(35)–C(36)–C(37)	143.8(5)	C(36)–C(37)–C(38)	139.3(6)
C(37)–C(38)–C(39)	127.4(6)		

Complex 13 has been characterized by  $^1\text{H}$ ,  $^{13}\text{C}\{^1\text{H}\}$ , and  $^{31}\text{P}\{^1\text{H}\}$  NMR and mass spectroscopy. The  $^1\text{H}$  NMR spectrum of complex 13 in  $\text{CD}_2\text{Cl}_2$  shows the resonance of  $^t\text{Bu}$  at 0.54 ppm and the resonance of the methylene tethered with  $^t\text{Bu}$  at 0.90 ppm as a doublet ( $J_{\text{HH}} = 10.8$  Hz) and at 1.59 ppm as a triplet ( $J_{\text{HH}} = J_{\text{HP}} = 10.8$  Hz). In the  $^{13}\text{C}\{^1\text{H}\}$  NMR spectrum, the signal of the CO ligand was observed at 207.0 ppm as a doublet ( $J_{\text{CP}} = 18$  Hz) due to coupling with the phosphorus atom in the tetradentate ligand. Furthermore, the mass spectrum showed the ion peak of the complex cation of 13 at  $m/z$  674.3293 (calcd 674.1869).

The structure of 13 has also been confirmed by X-ray diffraction. A colorless single crystal suitable for single-crystal X-ray diffraction analysis was obtained by slow diffusion of hexane into a  $\text{CH}_2\text{Cl}_2$  solution of complex 13. A view of the molecular structure of complex cation of 13 is shown in Figure 3. Crystal data and refinement details are given in Table 1, and selected bond lengths and bond angles of complex 13 are given in Table 4. The geometry around ruthenium in 13 can be viewed as a distorted octahedron. The tertiary amine N atom is trans to the CO ligand. The two pyridyl units in 13 are cis to each other, and the  $\text{CH}_2\text{CMe}_3$  group is trans to one of the two pyridyl nitrogen atoms. For the two protons of the methylene in  $\text{Ru}-\text{CH}_2\text{Bu}^t$ , the



Figure 2. ORTEP diagram of the complex cation  $8^+$ .Scheme 6. Proposed Mechanism for the Reaction of Complex 1 and *tert*-Butylacetylene To Afford Complex 8

two  $\text{H}-\text{C}-\text{Ru}-\text{P}$  dihedral angles are 3 and  $109^\circ$ , respectively. The structural feature may account for why one proton of  $\text{Ru}-\text{CH}_2\text{Bu}^t$  couples with the phosphorus atom ( $J_{\text{HP}} = 10.8$  Hz) while the other does not in the  $^1\text{H}$  NMR spectrum.

Scheme 8 shows a plausible mechanism for the formation of complex 13. Complex 1 could react with *tert*-butylacetylene to give vinylidene complex 10 via intermediate 9. 10 could be attacked by  $\text{H}_2\text{O}$  to give intermediate 14. After tautomerization and the loss of the acetyl ligand, the resulting intermediate 15 could undergo a deinsertion reaction to afford complex 13.



Scheme 9. Reactions of Complex 8

Figure 4. ORTEP diagram of complex  $17^{2+}$ .

poor solubility in common solvents. As expected, isolated complex 8 reacted with *tert*-butylacetylene and  $\text{H}_2\text{O}$  to produce complex 13. The transformation is accelerated by adding HOAc. Apparently, HOAc can easily protonate the  $\eta^3$ -butynyl ligand in complex 8 to regenerate complex 1.

## CONCLUSION

Vinylidene complexes have been suggested as the key intermediates in the endo cycloisomerization of alkynols catalyzed by complex 1. In this work, we have successfully isolated and characterized complexes 4, 8, and 13 from the reactions of 1

with alkynes. Production of these complexes can also be interpreted via vinylidene intermediates. The experimental results provide indirect evidence that the catalytic cycloisomerization of alkynols catalyzed by complex 1 involves vinylidene complexes as the key intermediates.

## EXPERIMENTAL SECTION

**General Considerations.** All manipulations were carried out under a nitrogen atmosphere by using standard Schlenk techniques, unless otherwise stated. Solvents were distilled under nitrogen from sodium benzophenone (hexane, diethyl ether, THF, benzene) or

**Table 5. Selected Bond Lengths (Å) and Angles (deg) in Complex 17**

Bond Distances			
Ru(1)–N(10)	2.0650(16)	Ru(1)–N(1)	2.0958(16)
Ru(1)–N(20)	2.1019(17)	Ru(1)–O(1)	2.1468(13)
Ru(1)–O(1A)	2.1681(13)	Ru(1)–P(1)	2.2502(5)
Bond Angles			
N(10)–Ru(1)–N(1)	80.94(6)	N(10)–Ru(1)–N(20)	162.96(7)
N(1)–Ru(1)–N(20)	82.58(7)	N(10)–Ru(1)–O(1)	96.01(6)
N(1)–Ru(1)–O(1)	168.69(6)	N(20)–Ru(1)–O(1)	99.33(6)
N(10)–Ru(1)–O(1A)	84.77(6)	N(1)–Ru(1)–O(1A)	90.28(6)
N(20)–Ru(1)–O(1A)	91.02(6)	O(1)–Ru(1)–O(1A)	78.58(6)
N(10)–Ru(1)–P(1)	96.28(5)	N(1)–Ru(1)–P(1)	92.68(5)
N(20)–Ru(1)–P(1)	88.77(5)	O(1)–Ru(1)–P(1)	98.49(4)
O(1A)–Ru(1)–P(1)	176.98(4)	Ru(1)–O(1)–Ru(1A)	101.42(6)

calcium hydride (dichloromethane).  $[\text{Ru}(\text{N}_3\text{P})(\text{OAc})](\text{BPh}_4)$  (**1**) was prepared according to the literature method.<sup>21</sup> Other chemicals were used as received from Aldrich. Elemental analyses were performed by M-H-W Laboratories (Phoenix, AZ). Mass spectra were collected on a MALDI Micro MX mass spectrometer (MALDI) and an API QSTAR XL System (ESI).  $^1\text{H}$ ,  $^{13}\text{C}\{^1\text{H}\}$ , and  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra were collected on a Bruker AV 400 MHz NMR spectrometer.  $^2\text{D}$  NMR spectra were collected on a JEOL EX 400 MHz NMR spectrometer.  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  NMR chemical shifts are relative to TMS or the residue of deuterium solvents, and  $^{31}\text{P}\{^1\text{H}\}$  NMR chemical shifts are relative to 85%  $\text{H}_3\text{PO}_4$ .

**Complex 4.** A mixture of  $[\text{Ru}(\text{N}_3\text{P})(\text{OAc})](\text{BPh}_4)$  (**1**; 0.15 g, 0.16 mmol), 3-buten-1-ol (**2**; 60  $\mu\text{L}$ , 0.76 mmol), and DIPEA (130  $\mu\text{L}$ , 0.76 mmol) in THF (5 mL) was stirred under  $\text{N}_2$  at 80 °C for 1 h. The mixture was cooled to room temperature and concentrated to approximately 3 mL. The precipitated solid was collected by filtration, washed with THF,  $\text{Et}_2\text{O}$ , and hexane, and dried under vacuum to afford complex **4** as a yellow solid (68%, 0.11 g).  $^{31}\text{P}\{^1\text{H}\}$  NMR (162.0 MHz,  $\text{CD}_2\text{Cl}_2$ , 25 °C):  $\delta$  53.4 ppm (s).  $^1\text{H}$  NMR (400.1 MHz,  $\text{CD}_2\text{Cl}_2$ , 25 °C):  $\delta$  8.72–8.70 (m, 1H), 8.38 (d,  $J$  = 4.8 Hz, 1H), 7.70–7.66 (m, 2H), 7.63–7.57 (m, 2H), 7.44–7.34 (m, 14H), 7.29–7.17 (m, 3H), 7.06–6.99 (m, 12H), 6.87 (t,  $J$  = 7.2 Hz, 4H), 6.78–6.74 (m, 2H), 6.57 (d,  $J$  = 7.8 Hz, 1H), 4.64 (d,  $J$  = 13.0 Hz, 1H;  $\text{ArCH}_2$ ), 4.25 (q,  $J_1$  = 16.4,  $J_2$  = 8.7 Hz;  $\text{Ru}=\text{C}(\text{O}-)\text{CH}_2\text{CH}_2\text{CH}_2-$ ), 4.06 (d,  $J$  = 17.4 Hz, 1H;  $\text{PyCH}_2$ ), 4.01–3.95 (m, 1H;  $\text{Ru}=\text{C}(\text{O}-)\text{CH}_2\text{CH}_2\text{CH}_2-$ ), 3.76 (d,  $J$  = 13.2 Hz, 1H;  $\text{ArCH}_2$ ), 3.72–3.65 (m, 1H;  $\text{Ru}=\text{C}(\text{O}-)\text{CH}_2$ ), 3.40 (d,  $J$  = 15.4 Hz, 1H;  $\text{PyCH}_2$ ), 3.22 (d,  $J$  = 15.4 Hz, 1H;  $\text{PyCH}_2$ ), 3.06 (d,  $J$  = 17.4 Hz, 1H;  $\text{PyCH}_2$ ), 2.04–1.92 (m, 1H;  $\text{Ru}=\text{C}(\text{O}-)\text{CH}_2$ ), 1.97 (s, 3H;  $\text{COCH}_3$ ), 1.90–1.81 (m, 1H;  $\text{Ru}=\text{C}(\text{O}-)\text{CH}_2\text{CH}_2$ ), 1.75–1.66 (m, 1H;  $\text{Ru}=\text{C}(\text{O}-)\text{CH}_2\text{CH}_2$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $\text{CD}_2\text{Cl}_2$ , 100.6 MHz, 25 °C):  $\delta$  312.7 (d,  $J$  = 14 Hz,  $\text{Ru}=\text{C}$ ), 177.2 ( $\text{CH}_3\text{CO}$ ), 165.4, 157.4, 156.6, 149.0, 138.4, 137.8, 136.4, 135.8, 134.5, 134.4, 132.6, 132.5, 132.1, 129.9, 129.4, 128.2, 128.1, 127.9, 127.7, 126.1, 122.3, 120.1, 81.7 ( $\text{Ru}=\text{C}(\text{O}-)\text{CH}_2\text{CH}_2\text{CH}_2-$ ), 74.2 ( $\text{PyCH}_2$ ), 67.7 (d,  $J$  = 4 Hz,  $\text{ArCH}_2$ ), 65.9 ( $\text{PyCH}_2$ ), 53.2 ( $\text{Ru}=\text{C}(\text{O}-)\text{CH}_2\text{CH}_2\text{CH}_2-$ ), 25.1 ( $\text{CH}_3\text{CO}$ ), 22.5 ( $\text{Ru}=\text{C}(\text{O}-)\text{CH}_2\text{CH}_2\text{CH}_2-$ ) ppm. Anal. Calcd for  $\text{C}_{61}\text{H}_{57}\text{BN}_3\text{O}_3\text{PRu} \cdot 0.5(\text{hexane})$ : C, 72.10; H, 6.05; N, 3.94. Found: C, 72.64; H, 5.94; N, 3.97. HRMS (ESI):  $m/z$  calcd for  $\text{C}_{37}\text{H}_{37}\text{N}_3\text{O}_3\text{PRu}^+$  704.1611, found 704.1570 [ $\text{M} - \text{BPh}_4$ ] $^+$ .

**Complex 8.** A mixture of complex **1** (0.3 g, 0.3 mmol) and *tert*-butylacetylene (**7**; 0.38 mL, 3.0 mmol) in THF (12 mL) was stirred under  $\text{N}_2$  in a sealed tube at 90 °C for 20 min. The reaction mixture was cooled to room temperature, and the solution was concentrated to 1 mL; then  $\text{Et}_2\text{O}$  (10 mL) was added to give a precipitate. The solvent was removed by filtration, and the solid was dissolved in  $\text{CH}_2\text{Cl}_2$  (1 mL); then  $\text{Et}_2\text{O}$  (10 mL) was added to the solution to afford a precipitate. The solid was collected by

filtration, washed with  $\text{Et}_2\text{O}$ , and dried under vacuum to afford complex **8** as a yellow solid (71%, 0.21 g).  $^{31}\text{P}\{^1\text{H}\}$  NMR (162.0 MHz,  $\text{CD}_2\text{Cl}_2$ , 25 °C):  $\delta$  50.2 ppm (s).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 400.1 MHz, 25 °C):  $\delta$  9.75 (d,  $J$  = 5.2 Hz, 1H), 7.74 (d,  $J$  = 5.4 Hz, 1H), 7.49–7.39 (m, 10H), 7.36–7.18 (m, 7H), 7.10–6.99 (m, 11H), 6.89–6.80 (m, 8H), 6.72–6.65 (m, 3H), 6.41 (d,  $J$  = 7.8 Hz, 1H), 4.80 (s, 1H;  $\text{C}=\text{CHC}(\text{CH}_3)_3$ ), 4.70 (d,  $J$  = 13.1 Hz, 1H;  $\text{ArCH}_2$ ), 4.54 (d,  $J$  = 15.0 Hz, 1H;  $\text{PyCH}_2$ ), 4.07 (d,  $J$  = 13.2 Hz, 1H;  $\text{ArCH}_2$ ), 3.68 (d,  $J$  = 16.7 Hz, 1H;  $\text{PyCH}_2$ ), 3.64 (d,  $J$  = 14.7 Hz, 1H;  $\text{PyCH}_2$ ), 3.10 (d,  $J$  = 14.7 Hz, 1H;  $\text{PyCH}_2$ ), 1.29 (s, 9H;  $\text{C}=\text{CC}(\text{CH}_3)_3$ ), 0.76 (s, 9H;  $\text{C}=\text{CHC}(\text{CH}_3)_3$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100.6 MHz,  $\text{CDCl}_3$ , 25 °C):  $\delta$  164.9, 164.4, 163.9, 163.4, 155.7, 148.6, 144.7 ( $\text{C}=\text{CC}=\text{CH}$ ), 141.2, 136.8, 136.4, 134.7, 133.6 ( $\text{C}=\text{CHC}(\text{CH}_3)_3$ ), 132.7, 132.6, 130.9, 129.5, 129.3, 129.0, 128.7, 128.3, 128.1, 127.9, 127.6, 127.2, 125.7, 123.7 ( $\text{C}=\text{CC}=\text{CH}$ ), 123.4, 122.0, 121.4, 70.5 ( $\text{PyCH}_2$ ), 67.8 ( $\text{ArCH}_2$ ), 64.8 ( $\text{PyCH}_2$ ), 48.9 ( $\text{C}=\text{CC}=\text{CH}$ ), 35.3 ( $\text{C}=\text{CHC}(\text{CH}_3)_3$ ), 33.7 ( $\text{C}=\text{CC}(\text{CH}_3)_3$ ), 32.5 ( $\text{C}=\text{CC}(\text{CH}_3)_3$ ), 29.7 ( $\text{C}=\text{CHC}(\text{CH}_3)_3$ ). Anal. Calcd for  $\text{C}_{67}\text{H}_{69}\text{BN}_3\text{OPRu}$ : C, 74.85; H, 6.47; N, 3.91. Found: C, 74.99; H, 6.19; N, 4.08. HRMS (MALDI, matrix CHCA):  $m/z$  calcd for  $\text{C}_{43}\text{H}_{47}\text{N}_3\text{PRu}^+$  738.2546, found 738.4191 [ $\text{M} - \text{BPh}_4$ ] $^+$ .

**Complex 13.** A mixture of complex **1** (0.10 g, 0.10 mmol), *tert*-butylacetylene (**7**; 0.25 mL, 2.0 mmol), and  $\text{H}_2\text{O}$  (0.2 mL) in THF (8 mL) was stirred under  $\text{N}_2$  in a sealed tube at 90 °C for 6 h. The reaction mixture was cooled to room temperature, and  $\text{Na}_2\text{SO}_4$  was added. After 2 h, the solid was removed by filtration, the filtrate was concentrated to approximately 1 mL, and  $\text{Et}_2\text{O}$  (7 mL) was added to give a white precipitate. The solid was collected by filtration, washed with a mixed solvent of  $\text{CH}_2\text{Cl}_2$  and  $\text{Et}_2\text{O}$  (1:5), and dried under vacuum to afford complex **13** as a white solid (76%, 76 mg).  $^{31}\text{P}\{^1\text{H}\}$  NMR (162.0 MHz,  $\text{CD}_2\text{Cl}_2$ , 25 °C):  $\delta$  56.7 ppm (s).  $^1\text{H}$  NMR (400.1 MHz,  $(\text{CD}_3)_2\text{CO}$ , 25 °C):  $\delta$  8.81 (d,  $J$  = 5.5 Hz, 1H), 8.25 (d,  $J$  = 5.3 Hz, 1H), 7.76–7.68 (m, 2H), 7.64–7.42 (m, 9H), 7.39–7.35 (m, 8H), 7.29–7.21 (m, 3H), 7.19–7.14 (m, 4H), 7.02–6.98 (m, 2H), 6.93 (t,  $J$  = 7.4 Hz, 8H), 6.78 (t,  $J$  = 7.2 Hz, 4H), 5.07 (d,  $J$  = 14.1 Hz, 1H;  $\text{ArCH}_2$ ), 5.05 (d,  $J$  = 16.1 Hz;  $\text{PyCH}_2$ ), 4.73 (d,  $J$  = 17.2 Hz;  $\text{PyCH}_2$ ), 4.45 (d,  $J$  = 16.1 Hz;  $\text{PyCH}_2$ ), 4.32 (dd,  $J_1$  = 14.1 Hz,  $J_2$  = 1.0 Hz, 1H;  $\text{ArCH}_2$ ), 4.23 (d,  $J$  = 17.2 Hz;  $\text{PyCH}_2$ ), 1.59 (t,  $J$  = 10.8 Hz, 1H;  $\text{RuCH}_2$ ), 0.90 (d,  $J$  = 10.8 Hz, 1H;  $\text{RuCH}_2$ ), 0.54 (s, 9H;  $\text{CH}_2\text{C}(\text{CH}_3)_3$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR (100.6 MHz,  $\text{CD}_2\text{Cl}_2$ , 25 °C):  $\delta$  207.0 (d,  $J$  = 18 Hz), 165.1, 164.7, 164.2, 163.7, 156.6, 156.0, 152.0, 150.2, 141.7, 141.5, 138.3, 137.8, 137.5, 136.4, 135.7, 135.2, 133.9, 133.7, 133.6, 133.0, 132.3, 131.9, 131.8, 130.6, 129.9, 129.6, 128.8, 128.7, 128.6, 128.5, 126.6, 126.2, 126.1, 124.8, 124.7, 122.9, 122.4, 121.7, 70.6 ( $\text{PyCH}_2$ ), 66.8 ( $\text{PyCH}_2$ ), 62.6 (d,  $J$  = 4 Hz,  $\text{ArCH}_2$ ), 37.3 ( $\text{C}(\text{CH}_3)_3$ ), 33.6 ( $\text{C}(\text{CH}_3)_3$  and  $\text{RuCH}_2$ ). Anal. Calcd for  $\text{C}_{61}\text{H}_{59}\text{BN}_3\text{OPRu}$ : C, 73.78; H, 5.99; N, 4.23. Found: C, 73.45; H, 6.02; N, 4.09. HRMS (MALDI, matrix CHCA):  $m/z$  calcd for  $\text{C}_{37}\text{H}_{39}\text{N}_3\text{OPRu}^+$  674.1869, found 674.3293 [ $\text{M} - \text{BPh}_4$ ] $^+$ .

**[D<sub>2</sub>] Complex 16.** A mixture of complex **1** (0.10 g, 0.10 mmol), *tert*-butylacetylene (**7**; 0.25 mL, 2.0 mmol) and  $\text{D}_2\text{O}$  (0.2 mL) in THF (8 mL) was stirred under  $\text{N}_2$  in a sealed tube at 90 °C for 6 h. The reaction mixture was cooled to room temperature, and  $\text{Na}_2\text{SO}_4$  was added. After 2 h, the solid was removed by filtration, the filtrate was concentrated to approximately 1 mL, and  $\text{Et}_2\text{O}$  (7 mL) was added to give a white precipitate. The solid was collected by filtration, washed with a mixed solvent of  $\text{CH}_2\text{Cl}_2$  and  $\text{Et}_2\text{O}$  (1:5), and dried under vacuum to afford complex **16** as a white solid (68%, 68 mg).  $^{31}\text{P}\{^1\text{H}\}$  NMR (162.0 MHz,  $\text{CD}_2\text{Cl}_2$ , 25 °C):  $\delta$  56.8 ppm (s).  $^1\text{H}$  NMR (400.1 MHz,  $(\text{CD}_3)_2\text{CO}$ , 25 °C):  $\delta$  8.82 (d,  $J$  = 5.6 Hz, 1H), 8.26 (d,  $J$  = 5.2 Hz, 1H), 7.78–7.68 (m, 2H), 7.66–7.46 (m, 9H), 7.38–7.34 (m, 8H), 7.29–7.23 (m, 3H), 7.19–7.14 (m, 4H), 7.06–6.70 (m, 2H), 6.93 (t,  $J$  = 7.4 Hz, 8H), 6.78 (t,  $J$  = 7.2 Hz, 4H), 5.08 (d,  $J$  = 14.5 Hz, 2H;  $\text{ArCH}_2$  and  $\text{PyCH}_2$ ), 4.76 (d,  $J$  = 17.0 Hz, 1H;  $\text{PyCH}_2$ ), 4.50 (d,  $J$  = 16.1 Hz;  $\text{PyCH}_2$ ), 4.35 (dd,  $J_1$  = 14.2 Hz,  $J_2$  = 1.2 Hz, 1H;  $\text{ArCH}_2$ ), 4.27 (d,  $J$  = 17.2 Hz;  $\text{PyCH}_2$ ), 0.53 (s, 9H;  $\text{CH}_2\text{C}(\text{CH}_3)_3$ ).  $^2\text{D}$  NMR (61.3 MHz,  $(\text{CH}_3)_2\text{CO}$  (set as  $\delta$  2.20 ppm), 25 °C):  $\delta$  1.66 (1D;  $\text{RuCD}_2$ ); 0.99 ppm (1D;  $\text{RuCD}_2$ ). Anal. Calcd for  $\text{C}_{61}\text{H}_{57}\text{D}_2\text{BN}_3\text{OPRu}$ : C, 73.63; H+D, 6.18; N, 4.22. Found: C, 73.75; H, 5.84; N, 4.26. HRMS (MALDI, matrix



CHCA):  $m/z$  calcd (%) for  $C_{37}H_{39}N_3OPRu^+$  676.1994, found 676.3060  $[M - BPh_4]^+$ .

**Complex 17.** A mixture of complex 8 (0.050 g, 0.047 mmol) and  $H_2O$  (90  $\mu L$ , 5.0 mmol) in THF (5 mL) was heated under  $N_2$  at 80 °C for 20 h. The mixture was cooled to room temperature. The precipitated solid was collected by filtration, washed with a solvent mixture of THF (0.5 mL) and  $Et_2O$  (5 mL), and dried under vacuum to afford complex 17 as a red solid (57%, 0.024 g). Anal. Calcd for  $C_{110}H_{98}B_2N_6O_2P_2Ru_2 \cdot 6H_2O$ : C, 68.46; H, 5.75; N, 4.35. Found: C, 68.35; H, 5.82; N, 4.30.

**X-ray Crystallography Studies of Compounds 4, 8, 13, and 17.** The crystals were mounted on glass fibers, and the diffraction intensity data of 4 were collected with a Bruker CCD diffractometer with monochromated Mo K $\alpha$  radiation ( $\lambda = 0.71073 \text{ \AA}$ ) at 173 K. Lattice determination and data collection were carried out using SMART v.5.625 software. Data reduction and absorption correction were performed using SAINT v 6.26 and SADABS v 2.03. However, the diffraction intensity data of 8, 13, and 17 were collected with an Oxford Diffraction GeminiS Ultra with CCD area detector with monochromated Cu K $\alpha$  radiation ( $\lambda = 1.54178 \text{ \AA}$ ) at 173 K. Lattice determination, data collection, and reduction were carried out using CrysAlisPro 171.32.5. Absorption correction was performed using SADABS built into the CrysAlisPro program suite. Structure solution and refinement for all three compounds were performed by using the SHELXTL v.6.10 software package. They were solved by direct methods and refined by full-matrix least squares on  $F^2$ . All non-hydrogen atoms were refined anisotropically. The hydrogen atoms were introduced at their geometric positions and refined as riding atoms.

## ■ ASSOCIATED CONTENT

**S Supporting Information.** CIF files giving X-ray structural information for complexes 4, 8, 13, and 17. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*P.N.L.: fax, (+)86-21-64252758; e-mail, [liupn@ecust.edu.cn](mailto:liupn@ecust.edu.cn). G.J.: fax, (+)852-23581594; e-mail, [chjiag@ust.hk](mailto:chjiag@ust.hk).

## ■ ACKNOWLEDGMENT

This work was supported by the Hong Kong Research Grant Council (Project No. HKUST 601709), the National Natural Science Foundation of China (Project No. 20902020), the Shanghai Pujiang Talent Program (Project No. 09J1403500), and the Fundamental Research Funds for the Central Universities.

## ■ REFERENCES

- (1) (a) Alonso, F. I.; Beletsaya, P.; Yus, M. *Chem. Rev.* **2004**, *104*, 3079. (b) Tius, M. A. In *Modern Allene Chemistry*; Krause, N., Hashmi, A. S. K., Eds.; Wiley-VCH: Weinheim, Germany, Vol. 2, 2004; p 834. (c) Tani, K.; Kataoka, Y. In *Catalytic Heterofunctionalization*; Togni, A., Grützmacher, H., Eds.; Wiley-VCH: Weinheim, Germany, 2001; p 171.
- (2) (a) Koo, B.; McDonald, F. E. *Org. Lett.* **2007**, *9*, 1737; (b) Koo, B.; McDonald, F. E. *Org. Lett.* **2005**, *7*, 3621; (c) McDonald, F. E.; Reddy, K. S. *Angew. Chem.* **2001**, *113*, 3765; *Angew. Chem., Int. Ed.* **2001**, *40*, 3653.
- (3) (a) McDonald, F. E.; Reddy, K. S.; Díaz, Y. *J. Am. Chem. Soc.* **2000**, *122*, 4304. (b) Wipf, P.; Graham, T. H. *J. Org. Chem.* **2003**, *68*, 8798. (c) Moilanena, S. B.; Tan, D. S. *Org. Biomol. Chem.* **2005**, *3*, 798. (d) Davidson, M. H.; McDonald, F. E. *Org. Lett.* **2004**, *6*, 1601.

- (4) McDonald, F. E.; Gleason, M. M. *J. Am. Chem. Soc.* **1996**, *118*, 6648.
- (5) McDonald, F. E.; Gleason, M. M. *Angew. Chem.* **1995**, *107*, 356; *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 350.
- (6) Trost, B. M.; Rhee, Y. H. *Org. Lett.* **2004**, *6*, 4311.
- (7) McDonald, F. E. *Chem. Eur. J.* **1999**, *5*, 3103.
- (8) (a) Trost, B. M. *Acc. Chem. Res.* **2002**, *35*, 695. (b) Anastas, P. N.; Kirchhoff, M. M. *Acc. Chem. Res.* **2002**, *35*, 686. (c) Trost, B. M.; Toste, D. F.; Pinkerton, A. B. *Chem. Rev.* **2001**, *101*, 2067. (d) Trost, B. M. *Angew. Chem.* **1995**, *107*, 285; *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 259. (e) Trost, B. M. *Science* **1991**, *254*, 1471.
- (9) (a) McDonald, F. E.; Connolly, C. B.; Gleason, M. M.; Towne, T. B.; Treiber, K. D. *J. Org. Chem.* **1993**, *58*, 6952. (b) Ramana, C. V.; Mallik, R.; Gonnade, R. G. *Tetrahedron* **2008**, *64*, 219.
- (10) (a) McDonald, F. E.; Reddy, K. S. *J. Organomet. Chem.* **2001**, *617–618*, 444. (b) McDonald, F. E.; Bowman, J. L. *Tetrahedron Lett.* **1996**, *37*, 4675. (c) Alcázar, E.; Pletcher, J. M.; McDonald, F. E. *Org. Lett.* **2004**, *6*, 3877.
- (11) (a) McDonald, F. E.; Schultz, C. C. *J. Am. Chem. Soc.* **1994**, *116*, 9363. (b) McDonald, F. E.; Schultz, C. C.; Chatterjee, A. K. *Organometallics* **1995**, *14*, 3628.
- (12) (a) Nowroozi-Isfahani, T.; Musaev, D. G.; McDonald, F. E.; Morokuma, K. *Organometallics* **2005**, *24*, 2921. (b) Sheng, Y.; Musaev, D. G.; Reddy, K. S.; McDonald, F. E.; Morokuma, K. *J. Am. Chem. Soc.* **2002**, *124*, 4149.
- (13) Weyershausen, B.; Dötz, K. H. *Eur. J. Inorg. Chem.* **1999**, 1057.
- (14) (a) Trost, B. M.; Rhee, Y. H. *J. Am. Chem. Soc.* **2002**, *124*, 2528. (b) Trost, B. M.; Rudd, M. T.; Costa, M. G.; Lee, P. I.; Pomerantz, A. E. *Org. Lett.* **2004**, *6*, 4235. (c) Varela-Fernandez, A.; Gonzalez-Rodriguez, C.; Varela, J. A.; Castedo, L. C. *Saa Org. Lett.* **2009**, *11*, 5350. (d) Zacuto, M. J.; Tomita, D.; Pirada, Z.; Xu, F. *Org. Lett.* **2010**, *12*, 684.
- (15) Trost, B. M.; Rhee, Y. H. *J. Am. Chem. Soc.* **2003**, *125*, 7482.
- (16) (a) Arimitsu, S.; Hammond, G. B. *J. Org. Chem.* **2007**, *72*, 8559. (b) Jury, J. C.; Swamy, N. K.; Yazici, A.; Willis, A. C.; Pyne, S. G. *J. Org. Chem.* **2009**, *74*, 5523.
- (17) (a) Qing, F.-L.; Gao, W.-Z.; Ying, J. *J. Org. Chem.* **2000**, *65*, 2003. (b) Qing, F.-L.; Gao, W.-Z. *Tetrahedron Lett.* **2000**, *41*, 7727.
- (18) (a) Bianchini, C.; Peruzzini, M.; Zanobini, F.; Frediani, P.; Albinati, A. *J. Am. Chem. Soc.* **1991**, *113*, 5453. (b) Bianchini, C.; Frediani, P.; Masi, D.; Peruzzini, M.; Zanobini, F. *Organometallics* **1994**, *13*, 4616. (c) Bianchini, C.; Bohanna, C.; Esteruelas, M. A.; Frediani, P.; Meli, A.; Oro, L. A.; Peruzzini, M. *Organometallics* **1992**, *11*, 3837. (d) Goodwin, J. M.; Olmstead, M. M.; Patten, T. E. *J. Am. Chem. Soc.* **2004**, *126*, 14352. (e) Chen, X.; Xue, P.; Sung, H. H. Y.; Williams, I. D.; Peruzzini, M.; Bianchini, C.; Jia, G. *Organometallics* **2005**, *24*, 4330.
- (19) (a) Simmons, E. L. *J. Chem. Educ.* **1979**, *56*, 578. (b) Major, R. C.; Zhu, X.-Y. *J. Am. Chem. Soc.* **2003**, *125*, 8454. (c) Crabtree, R. H. In *The Organometallic Chemistry Of The Transition Metals*, 4th ed.; Wiley: Hoboken, NJ, 2005; p 3.
- (20) (a) McBee, J. L.; Bell, A. T.; Tilley, T. D. *J. Am. Chem. Soc.* **2008**, *130*, 16562. (b) Liu, P. N.; Zhou, Z. Y.; Lau, C. P. *Chem. Eur. J.* **2007**, *13*, 8610.
- (21) Liu, P. N.; Su, F. H.; Wen, T. B.; H. Sung, H.-Y.; Williams, I. D.; Jia, G. *Chem. Eur. J.* **2010**, *16*, 7889.
- (22) (a) Bruce, M. I. *Chem. Rev.* **1991**, *91*, 197. (b) Puerta, M. C.; Valerga, P. *Coord. Chem. Rev.* **1999**, *193–195*, 977. (c) Rigaut, S.; Touchard, D.; Dixneuf, P. H. *Coord. Chem. Rev.* **2004**, *248*, 1585. (d) Katayama, H.; Ozawa, F. *Coord. Chem. Rev.* **2004**, *248*, 1703. (e) Che, C. M.; Ho, C. M.; Huang, J. S. *Coord. Chem. Rev.* **2007**, *251*, 2145. For reviews on catalysis involving vinylidene intermediates, see, for example: (f) Bruneau, C.; Dixneuf, P. H. *Acc. Chem. Res.* **1999**, *32*, 311; (g) Bruneau, C.; Dixneuf, P. In *Metal Vinylidenes and Allenylidenes in Catalysis*; Wiley-VCH: Weinheim, Germany, 2008. (h) Bruneau, C.; Dixneuf, P. H. *Angew. Chem.* **2006**, *118*, 2232; *Angew. Chem., Int. Ed.* **2006**, *45*, 2176.
- (23) For stoichiometric reactions, see, for example: (a) Ouzzine K.; Le Bozec, H.; Dixneuf, P. H. *J. Organomet. Chem.* **1986**, *317*, C25.

- (b) Le Bozec, H.; Ouzzine, K.; Dixneuf, P. H. *Organometallics* **1991**, *10*, 2768. (c) Pilette, D.; Ouzzine, K.; Le Bozec, H.; Dixneuf, P. H.; Rickard, C. E. F.; Roper, W. R. *Organometallics* **1992**, *11*, 809. For catalytic reactions, see, for example: (d) Trost, B. M.; Dyker, G.; Kulawiec, R. J. *J. Am. Chem. Soc.* **1990**, *112*, 7809. (e) Trost, B. M.; Flygare, J. A. *J. Am. Chem. Soc.* **1992**, *114*, 5476. (f) Trost, B. M.; Kulawiec, R. J. *J. Am. Chem. Soc.* **1992**, *114*, 5579. (g) Nishibayashi, Y.; Takei, I.; Hidai, M. *Organometallics* **1997**, *16*, 3091. (h) Trost, B. M.; Vidal, B.; Thommen, M. *Chem. Eur. J.* **1999**, *5*, 1055. (i) Gemel, C.; Trimmel, G.; Slugovc, C.; Kremel, S.; Mereiter, K.; Schmid, R.; Kirchner, K. *Organometallics* **1996**, *15*, 3998. (j) Derien, S.; Dixneuf, P. H. *J. Chem. Soc., Chem. Commun.* **1994**, 2551. (k) Derien, S.; Jan, D.; Dixneuf, P. H. *Tetrahedron* **1996**, *52*, 5511. (l) Derien, S.; Gomez Vicente, B.; Dixneuf, P. H. *Chem. Commun.* **1997**, 1405. (m) Dérien, S.; Dixneuf, P. H. *J. Organomet. Chem.* **2004**, *689*, 1382.
- (24) Addition of water to vinylidene intermediates has been suggested in Ru-catalyzed hydration of terminal alkynes; see, for example: (a) Tokunaga, M.; Wakatsuki, Y. *Angew. Chem.* **1998**, *110*, 3024; *Angew. Chem., Int. Ed.* **1998**, *37*, 2867. (b) Grotjahn, D. B.; Incarvito, C. D.; Rheingold, A. L. *Angew. Chem.* **2001**, *113*, 4002; *Angew. Chem., Int. Ed.* **2001**, *40*, 3884. (c) Grotjahn, D. B.; Lev, D. A. *J. Am. Chem. Soc.* **2004**, *126*, 12232. (d) Alvarez, P.; Bassetti, M.; Gimeno, J.; Mancini, G. *Tetrahedron Lett.* **2001**, *42*, 8467. (e) Suzuki, T.; Tokunaga, M.; Wakatsuki, Y. *Tetrahedron Lett.* **2002**, *43*, 7531. (f) Chen, Y.; Ho, D. M.; Lee, C. J. *J. Am. Chem. Soc.* **2005**, *127*, 12184. (g) Tokunaga, M.; Suzuki, T.; Koga, N.; Fukushima, T.; Horiuchi, A.; Wakatsuki, Y. *J. Am. Chem. Soc.* **2001**, *123*, 11917.
- (25) For catalytic reactions involving addition of carboxylic acids to vinylidene intermediates, see, for example: (a) Mahé, R.; Dixneuf, P. H.; Rcolier, S. L. *Tetrahedron Lett.* **1986**, *27*, 6333. (b) Doucet, H.; Martin-Vaca, B.; Bruneau, C.; Dixneuf, P. H. *J. Org. Chem.* **1995**, *60*, 7247. (c) Picquet, M.; Bruneau, C.; Dixneuf, P. H. *Chem. Commun.* **1997**, 1201. (d) Picquet, M.; Fernandez, A.; Bruneau, C.; Dixneuf, P. H. *Eur. J. Org. Chem.* **2000**, 2361. (e) Jiménez-Tenorio, M.; Puerta, M. C.; Valerga, P.; Moreno-Dorado, F. J.; Guerra, F. M.; Massanet, G. M. *Chem. Commun.* **2001**, 2324. (f) Melis, K.; Samulkiewicz, P.; Rynkowski, J.; Verpoort, F. *Tetrahedron Lett.* **2002**, *43*, 2713.
- (26) See for example: (a) Goossen, L. J.; Salih, K. S. M.; Blanchot, M. *Angew. Chem.* **2008**, *120*, 8620; *Angew. Chem., Int. Ed.* **2008**, *47*, 8492. (b) Goossen, L. J.; Blanchot, M.; Salih, K. S. M.; Blanchot, M. *Synthesis* **2009**, 2283 and references therein.
- (27) (a) Goossen, L. J.; Rodríguez, N.; Goossen, K. *Angew. Chem.* **2008**, *120*, 3144; *Angew. Chem., Int. Ed.* **2008**, *47*, 3100. (b) Goossen, L. J.; Paetzold, J.; Koley, D. *Chem. Commun.* **2003**, 706. (c) Nicks, F.; Aznar, R.; Sainz, D.; Muller, G.; Demonceau, A. *Eur. J. Org. Chem.* **2009**, 5020.
- (28) (a) Le Bozec, H.; Ouzzine, K.; Dixneuf, P. H. *Organometallics* **1991**, *10*, 2768. (b) Beddoes, R. L.; Grime, R. W.; Hussain, Z. I.; Whiteley, M. W. *J. Organomet. Chem.* **1996**, *526*, 371. (c) Barthel-Rosa, L. P.; Maitra, K.; Fischer, J.; Nelson, J. H. *Organometallics* **1997**, *16*, 1714. (d) Leung, W.-H.; Chan, E. Y. Y.; Williams, I. D.; Wong, W.-T. *Organometallics* **1997**, *16*, 3234. (e) Hansen, H. D.; Nelson, J. H. *Organometallics* **2000**, *19*, 4740. (f) Gamasa, M. P.; Gimeno, J.; Martín-Vaca, B. M.; Isea, R.; Vegas, A. *J. Organomet. Chem.* **2002**, *651*, 22. (g) Keller, A.; Jasionka, B.; Głowiak, T.; Ershov, A.; Matusiak, R. *Inorg. Chim. Acta* **2003**, *344*, 49. (h) Pavlik, S.; Mereiter, K.; Puchberger, M.; Kirchner, K. *J. Organomet. Chem.* **2005**, *690*, 5497. (i) Kopf, H.; Pietraszuk, C.; Hübner, E.; Burzlaff, N. *Organometallics* **2006**, *25*, 2533.
- (29) (a) Jia, G.; Rheingold, A. L.; Meek, D. W. *Organometallics* **1989**, *8*, 1378. (b) Jia, G.; Gallucci, J. C.; Rheingold, A. L.; Haggerty, B. S.; Meek, D. W. *Organometallics* **1991**, *10*, 3459. (c) Field, L. D.; Magill, A. M.; Jensen, P. *Organometallics* **2008**, *27*, 6539. (d) Lynam, J. M.; Nixon, T. D.; Whitwood, A. C. *J. Organomet. Chem.* **2008**, *693*, 3103. (e) Bassetti, M.; Marini, S. *Organometallics* **2002**, *21*, 4815. (f) Tenorio, M. A. J.; Tenorio, M. J.; Puerta, M. C.; Valerga, P. *Organometallics* **2000**, *19*, 1333. (g) Yi, C. S.; Liu, N. *Organometallics* **1997**, *16*, 3910. (h) Yang, S.-M.; Chan, M. C.-W.; Cheung, K.-K.; Che, C.-M.; Peng, S.-M. *Organometallics* **1997**, *16*, 2819. (i) Santos, A.; López, J.; Matas, L.; Ros, J.; Galán, A.; Echavarren, A. M. *Organometallics* **1993**, *12*, 4215. (j) Pavlik, S.; Gemel, C.; Slugovc, C.; Mereiter, K.; Schmid, R.; Kirchner, K. *J. Organomet. Chem.* **2001**, *617–618*, 301. (k) Liles, D. C.; Verhoeven, P. F. M. *J. Organomet. Chem.* **1996**, *522*, 33.
- (30) (a) Chen, C.-K.; Tong, H.-C.; Hsu, C.-Y. C.; Lee, C.-Y.; Fong, Y. H.; Chuang, Y.-S.; Lo, Y.-H.; Lin, Y.-C.; Wang, Y. *Organometallics* **2009**, *28*, 3358. (b) Slugovc, C.; Mereiter, K.; Zobetz, E.; Schmid, R.; Kirchner, K. *Organometallics* **1996**, *15*, 5275. (c) Bianchini, C.; Peruzzini, M.; Zanolini, F.; Frediani, P.; Albinati, A. *J. Am. Chem. Soc.* **1991**, *113*, 5453. (d) Bianchini, C.; Frediani, P.; Masi, D.; Peruzzini, M.; Zanolini, F. *Organometallics* **1994**, *13*, 4616. (e) Barbaro, C.; Bianchini, C.; Peruzzini, M.; Polo, A.; Zanolini, F.; Frediani, P. *Inorg. Chim. Acta* **1994**, *220*, 5.
- (31) (a) Cheng, C.-J.; Tong, H.-C.; Fong, Y.-H.; Wang, P.-Y.; Kuo, Y.-L.; Lo, Y.-H.; Lind, C.-H. *Dalton Trans.* **2009**, 4435. (b) Jia, G.; Meek, D. W. *Organometallics* **1991**, *10*, 1444.
- (32) (a) Mountassir, C.; Hedda, T. B.; Bozec, H. Le. *J. Organomet. Chem.* **1990**, 388, c13. (b) Bianchini, C.; Casares, J. A.; Peruzzini, M.; Romero, A.; Zanolini, F. *J. Am. Chem. Soc.* **1996**, *118*, 4585. (c) Lynam, J. M.; Welby, C. E.; Whitwood, A. C. *Organometallics* **2009**, *28*, 1320.
- (33) Sullivan, B. P.; Smythe, R. S.; Kober, E. M.; T. Meyer, J. J. *Am. Chem. Soc.* **1982**, *104*, 4701.
- (34) (a) O'Connor, J. M.; Pu, L. *J. Am. Chem. Soc.* **1990**, *112*, 9013. (b) Jiménez, M. V.; Sola, E.; Martínez, A. P.; Lahoz, F. J.; Oro, L. A. *Organometallics* **1999**, *18*, 1125. (c) Chin, C. S.; Chong, D.; Maeng, B.; Ryu, J.; Kim, H.; Kim, M.; Lee, H. *Organometallics* **2002**, *21*, 1739. (d) Albertin, G.; Antoniutti, S.; Bacchi, A.; Pelizzi, G.; Piasente, F. *Dalton Trans.* **2003**, 2881. (e) Sridevi, V. S.; Fan, W. Y.; Leong, W. K. *Organometallics* **2007**, *26*, 1173.
- (35) (a) Abu Salah, O. M.; Bruce, M. I. *J. Chem. Soc., Dalton Trans.* **1974**, 2302. (b) Bruce, M. I.; Swincer, A. G. *Aust. J. Chem.* **1980**, *33*, 1471. (c) Knaup, W.; Werner, H. *J. Organomet. Chem.* **1991**, *411*, 471. (d) Buil, M. L.; Esteruelas, A.; Lopez, A. M.; Onate, E. *Organometallics* **1997**, *16*, 3169. (e) Grotjahn, D. B.; Kragulj, E. J.; Zeinalipour-Yazdi, C. D.; Miranda-Soto, V.; Lev, D. A.; Cooksy, A. L. *J. Am. Chem. Soc.* **2008**, *130*, 10860.