

Co-ordination chemistry of the mixed phosphorus(III)–phosphorus(V) phosphazene ligand $\text{Ph}_2\text{PNP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NPPh}_2$: crystal structures of *cis*- $[\text{Pt}(\text{CH}_3)_2\text{X}\{\text{Ph}_2\text{PNP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NPPh}_2-P,P'\}]$ ($\text{X} = \text{CH}_3$ or I)[†]

Alexandra M. Z. Slawin, Martin B. Smith and J. Derek Woollins*

Department of Chemistry, Loughborough University, Loughborough, Leics., UK LE11 3TU

The ligand $\text{Ph}_2\text{PNP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NPPh}_2$ **I** was smoothly oxidised at both terminal PPh_2 moieties with either elemental sulfur or grey selenium to give $\text{Ph}_2\text{P}(\text{E})\text{NP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NP}(\text{E})\text{Ph}_2$ ($\text{E} = \text{S}$ **II** or Se **III**) in good yields. Oxidation of **I** with either aqueous H_2O_2 (30% w/w) or Bu^tOOH (in decane) gave the known dioxide compound $[\text{Ph}_2\text{P}(\text{O})]_2\text{NH}$ presumably *via* a sequence of oxidation, protonation and P–P bond-cleavage reactions. Complexation of **I** to several palladium(II), platinum(II) and gold(I) starting materials, at ambient temperature, afforded new metal complexes in which **I** adopts either *P,P'*-chelating or *P,P'*-bridging co-ordination modes. Reaction of **I** (1 equivalent) with the zerovalent compounds $[\text{M}(\text{PPh}_3)_4]$ ($\text{M} = \text{Pt}$ or Pd) in toluene proceeded cleanly, with P–P bond cleavage, and formation of the known four-membered metallacycles $[\text{M}(\text{Ph}_2\text{PNPPh}_2-P,P')_2]$. All compounds were characterised by a combination of IR and NMR (^1H , ^{31}P - $\{^1\text{H}\}$, ^{195}Pt - $\{^1\text{H}\}$) spectroscopy and elemental analyses. Furthermore, the molecular structures of *cis*- $[\text{Pt}(\text{CH}_3)_2\text{X}\{\text{Ph}_2\text{PNP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NPPh}_2-P,P'\}]$ ($\text{X} = \text{CH}_3$ or I) have been determined by single-crystal X-ray diffraction and represent the first examples of MP_4N_2 metallacycles incorporating a late transition metal. The PtP_4N_2 rings in both compounds adopt 'tub-like' geometries.


Bis(diphenylphosphino)amine, $\text{Ph}_2\text{PNHPPh}_2$ (hereafter abbreviated dppa), is an excellent ligand for complexation to a range of metals, reminiscent in many respects to the more familiar diphosphine ligand $\text{Ph}_2\text{PCH}_2\text{PPh}_2$ (dppm).^{1–13} Moreover dppa is also a valuable phosphorus precursor to some attractive P–N containing compounds **A–C**. These may either be non-ligated or stabilised by co-ordination to a metal centre.^{8–10,12–17} Recently there has been interest in the unusual P–P coupled diphosphazene $\text{Ph}_2\text{PNP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NPPh}_2$ **I**, loosely related to the carbon backbone ligand $\text{Ph}_2\text{P}(\text{CH}_2)_4\text{PPh}_2$ (dppb).^{18–20} Indeed several synthetic routes (based on oxidative coupling reactions) to **I** utilising the lithiated anion $[\text{Ph}_2\text{PNPPh}_2]^-$ **A** have been developed, yet surprisingly the co-ordination chemistry of **I** has so far remained poorly understood.²¹ In contrast, dppa and **A–C** can be complexed to a range of transition metals and, furthermore, several different ligating modes have been observed.^{1–17,21–24}

The present work describes the preparation and characterisation of $\text{Ph}_2\text{P}(\text{E})\text{NP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NP}(\text{E})\text{Ph}_2$ ($\text{E} = \text{S}$ or Se) in which the P–P bond remains intact and also new complexes of **I** in which both *P,P'*-chelating and *P,P'*-bridging bonding modes are observed. X-Ray crystallography has been used to determine unambiguously the structures of the platinum(II) complexes *cis*- $[\text{Pt}(\text{CH}_3)_2\text{X}\{\text{Ph}_2\text{PNP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NPPh}_2-P,P'\}]$ ($\text{X} = \text{CH}_3$ or I) which represent the first examples of seven-membered MP_4N_2 (M = late transition-metal ion) ring compounds. Part of this work has previously been communicated.²¹

Experimental

General

Unless otherwise stated, manipulations were performed under an oxygen-free nitrogen atmosphere using predried solvents and standard Schlenk techniques. The ligand $\text{Ph}_2\text{PNP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NPPh}_2$ **I** was prepared according to the procedure



described by Braunstein *et al.*¹⁹ The metal complexes $[\text{M}(\text{X})\text{-Y}(\text{cod})]$ ($\text{M} = \text{Pt}$ or Pd ; $\text{X}, \text{Y} = \text{Cl}, \text{Br}, \text{I}$ or CH_3 ; cod = cycloocta-1,5-diene),^{25–28} $[\text{PtCl}_2(\text{SEt}_2)_2]$,²⁹ $[\{\text{Pd}(\mu\text{-Cl})(\text{L-L})\}_2]$ [$\text{HL-L} = N,N$ -dimethylbenzylamine ($\text{C}_9\text{H}_{13}\text{N}$) or N,N -dimethyl-1-naphthylamine ($\text{C}_{12}\text{H}_{13}\text{N}$)],³⁰ $[\text{AuCl}(\text{tht})]$ (tht = tetrahydrothiophene)³¹ and $[\text{M}(\text{PPh}_3)_4]$ ³² ($\text{M} = \text{Pt}$ or Pd) were prepared according to literature methods. Aqueous H_2O_2 (30% w/w, Fluka), anhydrous Bu^tOOH in decane (Aldrich Chemical Co.) and MeI (Fisons) were used without further purification. Potassium tetrachloroplatinate(II), sodium tetrachloropalladate(II) and tetrachloroauric(III) acid were provided on loan by Johnson Matthey plc.

Infrared spectra were recorded as KBr pellets in the range 4000–220 cm^{-1} on a Perkin-Elmer System 2000 Fourier-transform spectrometer, ^1H NMR spectra (250 MHz) on a Bruker AC250 Fourier-transform spectrometer with chemical shifts (δ) in ppm (± 0.01) to high frequency of $\text{Si}(\text{CH}_3)_4$ and coupling constants (J) in Hz (± 0.1 Hz), ^{31}P - $\{^1\text{H}\}$ NMR spectra (36.2 or 101.3 MHz) either on a JEOL FX90Q or Bruker AC250 Fourier-transform spectrometer with chemical shifts (δ) in ppm (± 0.1) to high frequency of 85% H_3PO_4 and coupling constants (J) in Hz (± 3) and ^{195}Pt - $\{^1\text{H}\}$ NMR spectra (53.7 MHz) on a Bruker AC250 Fourier-transform spectrometer with chemical shifts (δ) referenced to external H_2PtCl_6 (in $\text{D}_2\text{O-HCl}$). All NMR spectra were measured in CDCl_3 unless otherwise stated. Elemental analyses (Perkin-Elmer 2400 CHN

[†] This paper is dedicated to Professor Sir Geoffrey Wilkinson FRS who provided inspiration and friendship through many years. He is sadly missed.

elemental analyser) were performed by the Loughborough University Service within the Department of Chemistry.

Preparation of the compounds

Ph₂P(S)NP(Ph)₂P(Ph)₂NP(S)Ph₂ II. To a solution of Ph₂PNP(Ph)₂P(Ph)₂NPPH₂ (0.352 g, 0.458 mmol) in thf (30 cm³) was added elemental sulfur (0.029 g, 0.905 mmol) as a solid in one portion. After stirring for ca. 5 min a white solid **II** deposited and the resulting mixture was stirred for 2.5 h. The solid was collected by suction filtration. Yield: 0.303 g, 80%. Selected spectroscopic data: δ (¹H) 7.83–7.49 and 7.35–7.18 (aromatic H); IR (KBr) 1207, 1174 (ν_{PN}), 603 cm⁻¹ (ν_{PS}).

Ph₂P(Se)NP(Ph)₂P(Ph)₂NP(Se)Ph₂ III. To the solids Ph₂PNP(Ph)₂P(Ph)₂NPPH₂ (0.440 g, 0.572 mmol) and grey Se (0.096 g, 1.216 mmol) was added tetrahydrofuran (thf) (45 cm³). The mixture was heated under reflux for 2 h, cooled and evaporated to dryness under reduced pressure. Extraction into CH₂Cl₂ (ca. 50 cm³), filtration through a small Celite pad and evaporation to dryness gave a pale yellow solid. Recrystallisation from CH₂Cl₂–Et₂O gave the product **III** as a white solid. Yield: 0.402 g, 76%. Selected spectroscopic data: δ (¹H) 7.83–7.51 and 7.35–7.18 (aromatic H); IR (KBr) 1206, 1174, 1160 (ν_{PN}), 550 cm⁻¹ (ν_{PSe}).

Reaction of Ph₂PNP(Ph)₂P(Ph)₂NPPH₂ with aqueous H₂O₂ (30% w/w). Under aerobic conditions, a thf (1 cm³) solution of Ph₂PNP(Ph)₂P(Ph)₂NPPH₂ (0.050 g, 0.065 mmol) was treated with aqueous H₂O₂ (30% w/w, 1 drop, ca. 0.038 g). The solution was stirred for ca. 21 h during which time [Ph₂P(O)]₂NH was deposited, collected by suction filtration and dried *in vacuo*. Yield: 0.042 g, 77%. Alternatively the same product was obtained using anhydrous Bu^oOOH in decane. An authentic sample of [Ph₂P(O)]₂NH was prepared by oxidation of dppa with aqueous H₂O₂ according to ref. 13.

cis-[PtCl₂(Ph₂PNP(Ph)₂P(Ph)₂NPPH₂-P,P')] 1. Under aerobic conditions, to a solution of [PtCl₂(cod)] (0.074 g, 0.198 mmol) in CH₂Cl₂ (20 cm³) was added Ph₂PNP(Ph)₂P(Ph)₂NPPH₂ (0.156 g, 0.203 mmol). The solution was stirred for 10 min, filtered and the volume concentrated by evaporation under reduced pressure to ca. 1–2 cm³. Addition of Et₂O (25 cm³) gave the product **1** which was collected by suction filtration and dried *in vacuo*. Yield: 0.185 g, 90%. Alternatively **1** was synthesized from [PtCl₂(SEt₂)] in 84% yield. Selected spectroscopic data: δ (¹H) 7.62–7.04 (aromatic H); IR (KBr) 1206, 1181, 1161 (ν_{PN}), 306, 279 cm⁻¹ (ν_{PtCl}).

The following compounds were prepared in a similar manner (yields in parentheses): *cis*-[PtBr₂(Ph₂PNP(Ph)₂P(Ph)₂NPPH₂-P,P')] **2** (92%), *cis*-[PtI₂(Ph₂PNP(Ph)₂P(Ph)₂NPPH₂-P,P')] **3** (93%), *cis*-[PdCl₂(Ph₂PNP(Ph)₂P(Ph)₂NPPH₂-P,P')] **4** (76%), *cis*-[PdBr₂(Ph₂PNP(Ph)₂P(Ph)₂NPPH₂-P,P')] **5** (90%), *cis*-[Pt(CH₃)₂(Ph₂PNP(Ph)₂P(Ph)₂NPPH₂-P,P')] **6** (91%) and *cis*-[Pt(CH₃)Cl(Ph₂PNP(Ph)₂P(Ph)₂NPPH₂-P,P')] **7** (63%). Selected spectroscopic data: **2**, δ (¹H) 7.62–7.04 (aromatic H); IR (KBr) 1197, 1180, 1161 cm⁻¹ (ν_{PN}); **3**, δ (¹H) 7.61–7.03 (aromatic H); IR (KBr) 1181, 1161 cm⁻¹ (ν_{PN}); **4**, IR (KBr) 1182, 1161 (ν_{PN}); 304, 276 cm⁻¹ (ν_{PdCl}); **5**, IR (KBr) 1182, 1162 cm⁻¹ (ν_{PN}); **6**, δ (¹H) 7.43–6.97 (aromatic H) and 0.09 [Pt–CH₃, *J*(PtH) 65.7, *J*(PH) 12.7 Hz]; **7**, δ (¹H) 7.61–7.08 (aromatic H) and 0.27 (Pt–CH₃); IR (KBr) 279 cm⁻¹ (ν_{PtCl}).

Slow evaporation of a CH₂Cl₂–light petroleum (b.p. 60–80 °C) filtrate gave crystals of complex **6** suitable for X-ray crystallography.

cis-[Pt(CH₃)I(Ph₂PNP(Ph)₂P(Ph)₂NPPH₂-P,P')] 8. A solution of *cis*-[Pt(CH₃)₂(Ph₂PNP(Ph)₂P(Ph)₂NPPH₂-P,P')] (0.029 g, 0.0292 mmol) in CH₂Cl₂ (ca. 0.6 cm³) was treated with iodomethane (ca. 0.18 g, 43-fold molar excess). After stirring for ca. 30 min, light petroleum (b.p. 60–80 °C) (3 cm³) was

added and the solid product **8** collected by suction filtration. Yield: 0.028 g, 88%. Selected spectroscopic data: δ (¹H) 7.58–7.05 (aromatic H) and 0.63 (Pt–CH₃). Slow diffusion of light petroleum into a CDCl₃ solution of complex **8** over the course of ca. 8 d gave crystals suitable for X-ray crystallography.

[PdCl(C₉H₁₂N)]₂{ μ -Ph₂PNP(Ph)₂P(Ph)₂NPPH₂} 9. To a yellow solution of [Pd(μ -Cl)(C₉H₁₂N)]₂ (0.040 g, 0.0724 mmol) in CH₂Cl₂ (10 cm³) was added solid Ph₂PNP(Ph)₂P(Ph)₂NPPH₂ (0.056 g, 0.0728 mmol). The solution was stirred for 30 min, filtered and the volume concentrated under reduced pressure to ca. 1–2 cm³. Addition of light petroleum (15 cm³) gave the solid product **9** which was collected by suction filtration and dried *in vacuo*. Yield: 0.080 g, 84%. Selected spectroscopic data: δ (¹H) 7.97–6.25 (aromatic H), 3.79 (N–CH₂) and 2.50 [*J*(PH) 2.3 Hz, N(CH₃)₂].

In a similar manner the complex [PdCl(C₁₂H₁₂N)]₂{ μ -Ph₂PNP(Ph)₂P(Ph)₂NPPH₂} **10** was prepared in 90% yield. Selected spectroscopic data: δ (¹H) 7.84–5.30 (aromatic H) and 3.09 [*J*(PH) 2.3 Hz, N(CH₃)₂].

[AuCl]₂{ μ -Ph₂PNP(Ph)₂P(Ph)₂NPPH₂} 11. To a solution of [AuCl(tht)] (0.104 g, 0.324 mmol) in CH₂Cl₂ (5 cm³) was added solid Ph₂PNP(Ph)₂P(Ph)₂NPPH₂ (0.126 g, 0.164 mmol). After ca. 5 min solid **11** deposited and the mixture was stirred for ca. 70 min. The product was collected by suction filtration and dried *in vacuo*. Yield: 0.153 g, 77%. Selected spectroscopic data: δ (¹H) 7.66–7.29 (aromatic H); IR (KBr) 1191, 1176, 1152 (ν_{PN}), 324 cm⁻¹ (ν_{AuCl}).

Reaction of Ph₂PNP(Ph)₂P(Ph)₂NPPH₂ with [Pt(PPh₃)₄]. To the solids Ph₂PNP(Ph)₂P(Ph)₂NPPH₂ (0.129 g, 0.168 mmol) and [Pt(PPh₃)₄] (0.208 g, 0.167 mmol) was added toluene (20 cm³). After stirring for ca. 24 h the white product [Pt(Ph₂PNP(Ph)₂P(Ph)₂NPPH₂-P,P')] **12** was collected by suction filtration and dried *in vacuo*. Yield: 0.145 g, 90%. Examination of the filtrate residue by ³¹P-{¹H} NMR spectroscopy revealed only PPh₃ [δ (P) –4.6] was present. In a similar manner the reaction of Ph₂PNP(Ph)₂P(Ph)₂NPPH₂ (0.070 g, 0.091 mmol) with [Pd(PPh₃)₄] (0.103 g, 0.089 mmol) was studied and found to give [Pd(Ph₂PNP(Ph)₂P(Ph)₂NPPH₂-P,P')] **13**. Yield: 0.061 g, 78%.

No reaction between [M(PPh₃)₄] (M = Pt or Pd) and Ph₂P(E)NP(Ph)₂P(Ph)₂NP(E)Ph₂ (E = S or Se) in toluene was observed even after prolonged (>10 d) stirring at ambient temperature.

X-Ray crystallography

The crystal structures of compounds **6** and **8** were obtained using a Rigaku AFC7S diffractometer with graphite-monochromated Cu-K α radiation (λ = 1.541 78 Å) and ω scans at room temperature. Details of the data collections and refinements are given in Table 3. Empirical absorption corrections (DIFABS)³³ were applied. The structures were solved by the heavy-atom method.³⁴ In **8** the CH₃ and I groups are disordered. The molecule was refined with both C and I atoms in 50% occupancy at each site. The disordered I atom positions were refined anisotropically and the disordered C atom positions isotropically, with no H atoms on the 50% occupancy methyl carbons being included in the refinement. All of the other non-hydrogen atoms were refined anisotropically. The C–H atoms were idealised and fixed (C–H 0.95 Å). No additional constraints or restraints were applied. Refinement was by full-matrix least-squares methods based on *F*. Calculations were performed using TEXSAN.³⁵

Atomic coordinates, thermal parameters, and bond lengths and angles have been deposited at the Cambridge Crystallographic Data Centre (CCDC). See Instructions for Authors, *J. Chem. Soc., Dalton Trans.*, 1997, Issue 1. Any request to the CCDC for this material should quote the full literature citation and the reference number 186/570.

Table 1 Microanalytical data for the new compounds (calculated values in parentheses)

Compound	Analysis (%)		
	C	H	N
II	69.2 (69.2)	4.55 (4.85)	3.25 (3.35)
III	62.2 (62.2)	4.15 (4.35)	2.95 (3.0)
1	55.1 (55.7)	4.0 (3.9)	2.7 (2.7)
2	51.2 (51.3)	3.6 (3.6)	2.6 (2.5)
3	47.4 (47.35)	3.2 (3.3)	1.5 (2.3)
4	60.25 (60.95)	4.4 (4.25)	3.0 (2.95)
5	55.05 (55.7)	3.9 (3.9)	2.45 (2.7)
6	60.0 (60.4)	4.25 (4.65)	2.6 (2.8)
7	56.6 (58.0)	4.15 (4.3)	2.8 (2.75)
8	53.25 (53.20)	4.15 (3.95)	2.7 (2.55)
9	61.6 (62.05)	4.95 (4.65)	3.9 (4.0)
10	59.55 (60.0)	4.9 (4.9)	4.3 (4.25)
11	46.45 (46.75)	2.9 (3.25)	2.4 (2.25)

Results and Discussion

Oxidation reactions of $\text{Ph}_2\text{PNP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NPPh}_2$

Previous work by ourselves^{36,37} has shown that dppa can be selectively monooxidised to afford the unsymmetrical ligands $\text{Ph}_2\text{PNHP}(\text{E})\text{Ph}_2$ (E = O, S or Se). Further oxidation leads to the formation of the doubly oxidised compounds $[\text{Ph}_2\text{P}(\text{E})]_2\text{NH}$.^{13,37–39} We have found that compound **I** can also be smoothly oxidised, under conditions similar to those employed for dppa, with elemental sulfur or grey selenium in thf to yield $\text{Ph}_2\text{P}(\text{E})\text{NP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NP}(\text{E})\text{Ph}_2$ (E = S **II**, E = Se **III**), both isolated as white air-stable solids in 80 and 76% yields respectively. Moreover the P–P bond in these ligands is not cleaved (X-ray evidence).²¹ This contrasts with the analogous reaction of **I** in thf, with either aqueous H_2O_2 (30% w/w) or anhydrous Bu^tOOH (in decane) which did not lead to the expected formation of $\text{Ph}_2\text{P}(\text{O})\text{NP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NP}(\text{O})\text{Ph}_2$ **IV** but instead gave the known compound $[\text{Ph}_2\text{P}(\text{O})]_2\text{NH}$ in 77% yield. However, careful monitoring (by $^{31}\text{P}\{-^1\text{H}\}$ NMR spectroscopy) of the reaction of **I** (in thf) with H_2O_2 revealed the formation of both $[\text{Ph}_2\text{P}(\text{O})]_2\text{NH}$ [$\delta(\text{P})$ 18.8] and a second species tentatively assigned to **IV** [$\delta \text{P}^{\text{III}}$, P^{V} 14.1, 9.6, $J(\text{PP})$ not resolved]. After leaving samples to stand for several hours only $[\text{Ph}_2\text{P}(\text{O})]_2\text{NH}$ precipitated from solution. No attempts to prepare **IV** using alternative oxidants or varying the experimental conditions have been tried. We, and others,¹⁸ have observed that solutions of **I** in various solvents are unstable and decompose. Hence, in thf we have identified a range of phosphorus-containing compounds including dppa, $\text{Ph}_2\text{PNHP}(\text{O})\text{Ph}_2$ and $[\text{Ph}_2\text{P}(\text{O})]_2\text{NH}$ in addition to several other uncharacterised species.

The mixed phosphorus(v) compounds **II** and **III** have the expected analytical (Table 1) and spectroscopic properties (Table 2 and Experimental section). Hence in the IR spectra of **II** and **III** $\nu_{(\text{PS})}$ and $\nu_{(\text{PSe})}$ are found at 603 and 550 cm^{-1} respectively, cf. $\text{Ph}_2\text{PNHP}(\text{S})\text{Ph}_2$ [$\nu_{(\text{PS})}$ 634 cm^{-1}], $[\text{Ph}_2\text{P}(\text{S})]_2\text{NH}$ [$\nu_{(\text{PS})}$ 645 cm^{-1}], $\text{Ph}_2\text{PNHP}(\text{Se})\text{Ph}_2$ [$\nu_{(\text{PSe})}$ 551 cm^{-1}] and $[\text{Ph}_2\text{P}(\text{Se})]_2\text{NH}$ [$\nu_{(\text{PSe})}$ 546 cm^{-1}].^{37,38} In the $^{31}\text{P}\{-^1\text{H}\}$ NMR spectrum of **III** the $^1J(\text{PSe})$ coupling constant of 714 Hz is smaller in magnitude than that reported for $\text{Ph}_2\text{PNHP}(\text{Se})\text{Ph}_2$ (757 Hz) and $[\text{Ph}_2\text{P}(\text{Se})]_2\text{NH}$ (793 Hz).^{37,38} The crystal structure of **III** has previously been reported by us and is in accord with the formulation shown.²¹


Chelate complexes of $\text{Ph}_2\text{PNP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NPPh}_2$

Treatment of $[\text{M}(\text{X})\text{Y}(\text{cod})]$ (M = Pt or Pd; X, Y = Cl, Br, I or CH_3 ; cod = cycloocta-1,5-diene) with compound **I** (1 : 1 molar ratio) in CH_2Cl_2 at room temperature gave the complexes *cis*- $[\text{M}(\text{X})\text{Y}\{\text{Ph}_2\text{PNP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NPPh}_2-P,P'\}]$ **1–7** in reasonable to good yields (63–93%). Alternatively complex **1** can be readily

Table 2 Selected spectroscopic data for compounds **II**, **III** and **1–13**

Compound	δ			Additional data
	$\text{P}^{\text{III}a}$	P^{V}	Pt	
II	47.2	13.5		$J(\text{PP})$ 7.2
III	38.4	14.5		$J(\text{PSe})$ 714 $J(\text{PP})$ 7.2
1	22.9 (4049)	1.1	–4157	
2	23.0 (4001)	0.3	–4408	
3	22.7 (3845)	–0.9	–4902	
4	48.8 ^b	–1.4		
5	47.8 ^b	–2.1		
6	51.7 (2111)	–0.1 ^c	–4354	$J(\text{PP})$ 13.3
7	54.4 (2006) ^d	4.1, 0.8 ^e	–4281	
	29.1 (4694) ^f			
8	48.5 (2072) ^d	2.3, –1.7 ^e	–4602	
	31.3 (4556) ^{fg}			
9	60.6	15.8		
10	62.4	15.3		
11	50.8	16.6		$J(\text{PP})$ 8.3
12	–15.9 (1667) ^h			
	–17.4 (1660) ⁱ			
13	–6.5 ^j			

^a Values in parentheses denote $^1J(\text{PtP})/\text{Hz}$. ^b Decomposes in solution. ^c $^3J(\text{PtP})/\text{Hz}$ not observed. ^d P *trans* to CH_3 . ^e Several $J(\text{PP})$ values between 5 and 35 Hz. ^f P *trans* to X (Cl or I). ^g Measured *in situ*. ^h Measured *in situ* in CH_2Cl_2 – CD_2Cl_2 (ω : 35 Hz). ⁱ Taken from ref. 8. ^j Measured *in situ* in CH_2Cl_2 – CD_2Cl_2 .



- 1** M = Pt, X = Y = Cl
- 2** M = Pt, X = Y = Br
- 3** M = Pt, X = Y = I
- 4** M = Pd, X = Y = Cl
- 5** M = Pd, X = Y = Br
- 6** M = Pt, X = Y = CH_3
- 7** M = Pt, X = CH_3 , Y = Cl
- 8** M = Pt, X = CH_3 , Y = I

prepared from $[\text{PtCl}_2(\text{SEt}_2)_2]$. The *cis* geometries of **1–7** were established in solution by multinuclear NMR spectroscopy (Table 2). The chemical shifts ($\delta \text{P}^{\text{III}}$) for **1–3** are typically observed at ca. 23.0 ppm (cf. **I** $\delta \text{P}^{\text{III}}$ 43), whilst on going from Cl > Br > I in the platinum(II) series **1–3** there is a small reduction in the magnitude of the $^1J(\text{PtX})$ coupling constant [$J(\text{PtP})$ 4049 for **1**, 4001 for **2**, 3845 Hz for **3**]. For all the dihalogeno complexes $J(\text{PP})$ was not resolved (at 36.2 MHz) whereas for the dimethylplatinum(II) complex **6** $J(\text{PP})$ is ca. 13 Hz. In the $^{195}\text{Pt}\{-^1\text{H}\}$ NMR spectra of **1–3** and **6** the expected 1 : 2 : 1 triplet was observed whilst the resonances for **7** and **8** were doublet of doublets (Table 2). When the complexes **1** and **2** are left to stand in CDCl_3 for prolonged periods (ca. 5 d) only trace amounts of a decomposition product (δP 21.0) are observed and tentatively assigned as $[\text{Ph}_2\text{P}(\text{O})]_2\text{NH}$. In contrast $^{31}\text{P}\{-^1\text{H}\}$ NMR spectra of freshly prepared CDCl_3 solutions of *cis*- $[\text{PdX}_2\{\text{Ph}_2\text{PNP}(\text{Ph})_2\text{P}(\text{Ph})_2\text{NPPh}_2-P,P'\}]$ (X = Cl **4** or Br **5**) indicate only single phosphorus-containing species. However after ca. 2 h conversion into new species ($\delta \text{P}^{\text{III}}$, P^{V} 52.9, 20.3 in the case of **4**; 51.7, 19.2 in the case of **5**) is then observed whereas after several hours insoluble products are formed and identified by IR (KBr pellets) as *cis*- $[\text{PdX}_2(\text{Ph}_2\text{PNHPPh}_2-P,P')]$ (X = Cl or Br).[‡] Clearly the P–P bond is susceptible to cleavage under these conditions.

Monitoring the reaction of complex **6** in toluene with an

[‡] Authentic samples of *cis*- $[\text{PdX}_2(\text{Ph}_2\text{PNHPPh}_2-P,P')]$ were prepared from $[\text{PdX}_2(\text{cod})]$ (X = Cl or Br) and 1 equivalent of $\text{Ph}_2\text{PNHPPh}_2$ in CH_2Cl_2 .


Table 3 Details of the X-ray data collections and refinements for compounds **6** and **8**

	6	8
Empirical formula	C ₅₀ H ₄₆ N ₂ P ₄ Pt	C ₄₈ H ₄₃ IN ₂ P ₄ Pt
<i>M</i>	993.91	1105.78
Crystal colour, habit	Clear, plate	Yellow, prism
Crystal dimensions/mm	0.20 × 0.03 × 0.29	0.12 × 0.21 × 0.21
Crystal system	Orthorhombic	Orthorhombic
Space group	<i>Pccn</i>	<i>Pbca</i>
<i>a</i> /Å	12.255(6)	20.807(8)
<i>b</i> /Å	34.724(9)	34.700(4)
<i>c</i> /Å	20.847(8)	12.453(5)
<i>U</i> /Å ³	8871	8990
<i>Z</i>	8	8
<i>D_c</i> /g cm ⁻³	1.49	1.63
<i>μ</i> /mm ⁻¹	7.64	12.63
2θ _{max} /°	120.0	119.9
<i>F</i> (000)	3984	4336
Measured reflections	7294	7366
Observed reflections [<i>I</i> > 2.0σ(<i>I</i>)]	2968	2748
Reflection: parameter ratio	5.8:1	5.3:1
Minimum, maximum transmission	0.73, 1.00	0.59, 1.00
Weighting scheme <i>p</i>	0.004	0.002
No. variables	515	523
Final <i>R</i> , <i>R'</i>	0.037, 0.032	0.045, 0.041
Maximum Δ/σ	0.64	0.64
Largest difference peak, hole/e Å ⁻³	0.62, -0.74	0.66, -0.95

$$R = \sum |F_o| - |F_c| / \sum |F_o|, R' = [\sum w(|F_o| - |F_c|)^2 / \sum w F_o^2]^{1/2}; \text{weighting scheme defined in ref. 35.}$$

Table 4 Selected bond distances (Å) and angles (°) for complexes **6** and **8**

	6	8
P(1)–N(1)	1.648(8)	1.628(10)
N(1)–P(2)	1.575(8)	1.589(9)
P(2)–P(3)	2.290(4)	2.242(5)
P(3)–N(2)	1.561(8)	1.552(10)
N(2)–P(4)	1.647(8)	1.652(10)
Pt–P(1)	2.283(3)	2.284(4)
Pt–P(4)	2.285(3)	2.279(4)
Pt–C(49)	2.21(1)	
Pt–C(50)	2.209(9)	
Pt–P(1)–N(1)	120.3(3)	118.9(4)
P(1)–N(1)–P(2)	126.9(5)	130.4(6)
N(1)–P(2)–P(3)	105.6(4)	104.0(4)
P(2)–P(3)–N(2)	110.1(4)	108.5(4)
P(3)–N(2)–P(4)	138.5(6)	139.6(7)
N(2)–P(4)–Pt	121.2(3)	120.4(4)
P(4)–Pt–P(1)	94.0(1)	94.0(1)
P(1)–Pt–C(49)	93.6(3)	
P(1)–Pt–C(50)	173.3(2)	
P(4)–Pt–C(49)	172.4(3)	
P(4)–Pt–C(50)	92.2(2)	
C(49)–Pt–C(50)	80.2(3)	

**Fig. 1** Crystal structure of *cis*-[Pt(CH₃)₂{Ph₂PNP(Ph)₂P(Ph)₂NPPh₂-*P,P'*}] **6**. Phenyl groups and CH protons omitted for clarity. The structure of complex **8** is very similar and is not illustrated. The numbering scheme is the same for the PtP₄N₂ rings in **6** and **8**


excess of CH₃I by ³¹P-{¹H} NMR spectroscopy indicated the exclusive formation of *cis*-[Pt(CH₃)I{Ph₂PNP(Ph)₂P(Ph)₂NPPh₂-*P,P'*}] **8** (Tables 1 and 2 for characterising data). We did not observe the formation of any intermediate platinum(IV), species *e.g.* [PtMe₃I{Ph₂PNP(Ph)₂P(Ph)₂NPPh₂-*P,P'*}].

The crystal structure of compound **6** (Table 4, Fig. 1) reveals square-planar co-ordination of the platinum(II) centre with the PtP₄N₂ ring adopting a 'tub-like' geometry. This can be envisaged as being made up of a Pt–P(1)–N(1)–P(2) plane of atoms [maximum deviation from mean plane 0.08 Å for N(1)] which is folded with respect to the Pt–P(2)–P(3)–P(4) mean plane (fold angle 109°). The N(2) atom lies out (0.37 Å) of the Pt–P(2)–P(3)–P(4) plane and is on the same side of the molecule as the Pt–P(1)–N(1)–P(2) plane. The bond lengths and angles in **6** are broadly as anticipated. The P(2)–P(3) distance is 2.290(4) Å whilst the P–N bond lengths divide into two groups: P(1)–N(1) and P(4)–N(2) are 1.648(8) and 1.647(8) Å respectively whilst P(2)–N(1) and P(3)–N(2) are 1.575(8) and 1.561(8) Å respectively reflecting some degree of bond alternation as one would

expect from simple bonding descriptions. Interestingly the P–P distance in **6** is elongated with respect to the same bond in free **1** [2.232(2) Å],¹⁹ **III** [2.218(4) Å]²¹ and the gold(I) compound [(AuCl)₂{μ-Ph₂PNP(Ph)₂P(Ph)₂NPPh₂}] [2.215(7) Å]²¹ whilst the P–N bond lengths are not so dramatically influenced. The lengthening of the P–P bond does not appear to be a consequence of ring formation since in the cyclic arsenic salt [AsPPh₂NP(Ph)₂P(Ph)₂NPPh₂]-4C₄H₈O the P–P distance is 2.241(3) Å.¹⁸ The crystal structure of **8** (Table 4 for selected bond lengths and angles) is very similar to that of **6**.

Bridging complexes of Ph₂PNP(Ph)₂P(Ph)₂NPPh₂

The bridge-cleavage reaction of [Pd(μ-Cl)(L–L)]₂ [HL–L = *N,N*-dimethylbenzylamine (C₉H₁₃N) or *N,N*-dimethyl-1-naphthylamine (C₁₂H₁₃N)] with 1 equivalent of compound **1** in CH₂Cl₂ yields the dimeric complexes [Pd(L–L)Cl]₂{μ-Ph₂PNP(Ph)₂P(Ph)₂NPPh₂}] (L–L = C₉H₁₂N **9** or C₁₂H₁₂N **10**). Related cyclopalladated dimers with bridging Ph₂X–(CH₂)₂XPh₂ (X = P or As) ligands are known.⁴⁰ In the ³¹P-{¹H} NMR spectra of **9** and **10** the phosphorus(v) resonance is found at higher frequency (typically by *ca.* 15 ppm) than that observed for the chelate complexes **1–8** (typically *ca.* 0 ppm, Table 2). We have also prepared the dimeric gold(I) compound [(AuCl)₂{μ-Ph₂PNP(Ph)₂P(Ph)₂NPPh₂}] **11** as a white solid in 77% yield.



Oxidative-addition reactions with ligands I–III

A preliminary study on the reactivity of **I–III** with the zerovalent compounds [M(PPh₃)₄] (M = Pt or Pd) was undertaken in order to probe the susceptibility of the P–P bond to cleavage. We find that reaction of **I** with [M(PPh₃)₄] (1 : 1 ratio) in toluene yields the homoleptic metal(II) complexes [M(Ph₂PNPPh₂-P,P')] (M = Pt **12** or Pd **13**). ³¹P-{¹H} NMR spectroscopy revealed the predominant species upon evaporation of the toluene filtrate to be PPh₃. The ³¹P spectral data (Table 2) for **12** are in good agreement with those previously reported by Browning and Farrar.⁸ Monitoring these reactions by ³¹P-{¹H} NMR spectroscopy showed that oxidative addition was both clean and facile with only [M(Ph₂PNPPh₂-P,P')] and displaced PPh₃ identified as the sole products. We did not detect the formation of any intermediates (e.g. the d¹⁰ species **D**) in these reactions. Mixed metal(0) complexes of the type [M(PR₃)₂(P–P)] (P–P = chelating diphosphine) have previously been documented.⁴¹

Neutral compounds of the type [M{Ph₂PNP(E)Ph₂-P,E}]₂ **E** (*cis*) and **F** (*trans*) are known^{36,37} and can be readily synthesized from the monoxidised ligand Ph₂PNHP(E)Ph₂ and [MCl₂(cod)] (M = Pt or Pd) in the presence or absence of base. We were interested to see whether compounds **II** and **III** would be suitable precursors and undergo P–P cleavage to yield analogous complexes. Treatment of [M(PPh₃)₄] (M = Pt or Pd) with **II** (or **III**) in toluene for several days at ambient temperature gave no reaction (**II** and **III** recovered unchanged) which may in part reflect either the low solubility of **II** (or **III**) in toluene or the poor donor ability of these ligands. Related metal complexes containing the neutral chelating ligand [R₂P(E)]₂NH (R = alkyl or aryl, E = S or Se) are, to the best of our knowledge, restricted only to the cationic palladium(II) compound [Pd{(PrⁱPS)₂-NH}{(PrⁱPS)₂N}]Cl in which one of the ligands is protonated.⁴²

Conclusion

The results of our study clearly illustrate that compound **I** has the propensity to form several late transition-metal complexes in which the ligand adopts either a *P,P'*-chelating or *P,P'*-bridging mode. We also observe that the P–P bond in **I** is readily cleaved by [M(PPh₃)₄], but under similar conditions no reaction was observed with **II** (or **III**).

Acknowledgements

We are grateful to the EPSRC for funding and to Johnson Matthey plc for loans of precious metals.

References

- D. I. Arnold, F. A. Cotton and F. E. Kühn, *Inorg. Chem.*, 1996, **35**, 4733, 5764.
- F. A. Cotton and F. E. Kühn, *Inorg. Chim. Acta*, 1996, **252**, 257.
- R. Graziani and U. Casellato, *Acta Crystallogr., Sect. C*, 1996, **52**, 850.

- M. Paul and H. Schmidbaur, *Chem. Ber.*, 1996, **129**, 77.
- D. Pohl, J. Ellermann, M. Moll, F. A. Knoch and W. Bauer, *Z. Anorg. Allg. Chem.*, 1996, **622**, 283.
- P. Bhattacharyya and J. D. Woollins, *Polyhedron*, 1995, **14**, 3367 and refs. therein.
- D. Pohl, J. Ellermann, F. A. Knoch and M. Moll, *J. Organomet. Chem.*, 1995, **495**, C6.
- C. S. Browning and D. H. Farrar, *J. Chem. Soc., Dalton Trans.*, 1995, 521.
- D. Pohl, J. Ellermann, F. A. Knoch, M. Moll and W. Bauer, *Chem. Ber.*, 1994, **127**, 2167.
- D. Pohl, J. Ellermann, F. A. Knoch, M. Moll and W. Bauer, *J. Organomet. Chem.*, 1994, **481**, 259.
- G. Liehr, G. Szucsányi and J. Ellermann, *J. Organomet. Chem.*, 1984, **265**, 95.
- H. Schmidbaur, S. Lauteschläger and B. Milewski-Mahrle, *J. Organomet. Chem.*, 1983, **254**, 59.
- F. T. Wang, J. Najdzionek, K. L. Leneker, H. Wasserman and D. M. Braitsch, *Synth. React. Inorg. Metal-Org. Chem.*, 1978, **8**, 119.
- M. Gómez, G. Muller, J. Sales and X. Solans, *J. Chem. Soc., Dalton Trans.*, 1993, 221.
- J. Ellermann and M. Lietz, *Z. Naturforsch., Teil B*, 1980, **35**, 64.
- J. Ellermann, J. Sutter, C. Schelle, F. A. Knoch and M. Moll, *Z. Anorg. Allg. Chem.*, 1993, **619**, 2006.
- J. Ellermann, J. Sutter, F. A. Knoch and M. Moll, *Angew. Chem., Int. Ed. Engl.*, 1993, **32**, 700.
- M. Dotzler, A. Schmidt, J. Ellermann, F. A. Knoch, M. Moll and W. Bauer, *Polyhedron*, 1996, **15**, 4425.
- P. Braunstein, R. Hasselbring, A. Tiripicchio and F. Uguzzoli, *J. Chem. Soc., Chem. Commun.*, 1995, 37.
- J. Ellermann, P. Gabold, F. A. Knoch, M. Moll, A. Schmidt and M. Schütz, *Z. Naturforsch., Teil B*, 1996, **51**, 201.
- A. M. Z. Slawin, M. B. Smith and J. D. Woollins, *Chem. Commun.*, 1996, 2095.
- K. V. Katti, H. W. Roesky and M. Rietzel, *Inorg. Chem.*, 1987, **26**, 4032.
- H. W. Roesky, K. V. Katti, U. Seseke, M. Witt, E. Egert, R. Herbst and G. M. Sheldrick, *Angew. Chem., Int. Ed. Engl.*, 1986, **25**, 477.
- H. W. Roesky, K. V. Katti, U. Seseke, H.-G. Schmidt, E. Egert, R. Herbst and G. M. Sheldrick, *J. Chem. Soc., Dalton Trans.*, 1987, 847.
- J. X. McDermott, J. F. White and G. M. Whitesides, *J. Am. Chem. Soc.*, 1976, **98**, 6521.
- D. Drew and J. R. Doyle, *Inorg. Synth.*, 1972, **13**, 47.
- E. Costa, P. G. Pringle and M. Ravetz, *Inorg. Synth.*, 1997, **31**, 284.
- H. C. Clark and L. E. Manzer, *J. Organomet. Chem.*, 1973, **59**, 411.
- G. B. Kauffman and D. O. Cowan, *Inorg. Synth.*, 1960, **6**, 211.
- A. C. Hope and E. C. Friedrich, *J. Am. Chem. Soc.*, 1968, **90**, 909.
- R. Uson, A. Laguna and J. Vicente, *J. Organomet. Chem.*, 1977, **131**, 471.
- R. Ugo, F. Cariati and G. La Monica, *Inorg. Synth.*, 1968, **11**, 105; D. R. Coulson, *Inorg. Synth.*, 1972, **13**, 121.
- DIFABS, N. G. Walker and D. Stuart, *Acta Crystallogr., Sect. A*, 1983, **39**, 158.
- DIRDIF 92, PATTY, P. T. Beurskens, G. Admiraal, G. Beurskens, W. P. Bosman, S. Garcia-Granda, R. O. Gould, J. M. M. Smits and C. Smykalla, The DIRDIF program system, Technical Report of the Crystallography Laboratory, University of Nijmegen, 1992.
- TEXSAN, Crystal Structure Analysis Package, Molecular Structure Corporation, Woodlands, TX, 1985, 1992.
- P. Bhattacharyya, A. M. Z. Slawin, M. B. Smith and J. D. Woollins, *Inorg. Chem.*, 1996, **35**, 3675.
- P. Bhattacharyya, A. M. Z. Slawin, D. J. Williams and J. D. Woollins, *J. Chem. Soc., Dalton Trans.*, 1995, 3189.
- P. Bhattacharyya, J. Novosad, J. Phillips, A. M. Z. Slawin, D. J. Williams and J. D. Woollins, *J. Chem. Soc., Dalton Trans.*, 1995, 1607.
- I. Haiduc, in *Inorganic Experiments*, ed. J. D. Woollins, VCH, Weinheim, 1994.
- J. Dehand, M. Pfeffer and M. Zinsius, *Inorg. Chim. Acta*, 1975, **13**, 229.
- B. R. James and M. B. Smith, unpublished work; P. A. Chaloner and G. T. L. Broadwood-Strong, *J. Chem. Soc., Dalton Trans.*, 1996, 1039.
- D. Cupertino, R. Keyte, A. M. Z. Slawin, J. D. Woollins and D. J. Williams, *Polyhedron*, 1996, **15**, 4441.

Received 11th April 1997; Paper 7/02500I