

# Electronic Substituent Effects of Furoxan and Furazan Systems

Roberta Fruttero,<sup>a</sup> Donatella Boschi,<sup>a</sup> Elisa Fornatto,<sup>a</sup>  
Anna Serafino,<sup>a</sup> Alberto Gasco<sup>\*a</sup> and Otto Exner<sup>b</sup>

<sup>a</sup>Department of 'Scienza e Tecnologia del Farmaco', Facoltà di Farmacia di Torino, Via Giuria 9, 10125 Torino, Italy

<sup>b</sup>Institute of Organic Chemistry and Biochemistry, Academy of Sciences of the Czech Republic, 166 10 Praha 6, Czech Republic

*J. Chem. Research (S)*,  
1998, 495

*J. Chem. Research (M)*,  
1998, 2545–2562

Heterocyclic systems 3-furoxanyl, 4-furoxanyl and 3-furazanyl have been characterised as substituents by their substituent constants  $\sigma_m$ ,  $\sigma_p$ ,  $\sigma_I$  and  $\sigma_R$ , determined from pK and  $^{19}\text{F}$  NMR shifts: they are strongly electron attracting but weakly conjugated.

Furoxans are important in biochemistry as NO donors under physiologic conditions.<sup>4</sup> Parallel with their pharmacology, we have also investigated their physical properties: lipophilicity,<sup>6</sup> dipole moments and electron distribution.<sup>7</sup> They may be described as electron overcrowded molecules. Here we report the Hammett substituent constants  $\sigma_m$  and  $\sigma_p$  of 4-methyl-3-furoxanyl (**1a**), 3-methyl-4-furoxanyl (**1b**) and also of the parent heterocycle without oxygen, 4-methyl-3-furazanyl (**1c**). We used the standard approach<sup>11</sup> based on pK values of substituted benzoic acids **2a–c** and **3a–c** in 50% ethanol. The methyl group on the heterocycle was necessary for the stability of compounds and scarcely influences the  $\sigma$  values.

All the heterocyclic systems are rather strongly electron attracting by the inductive mechanism, as strong as *e.g.* halogens, but seem to be only very slightly conjugated as weak donors. We also determined the inductive constants  $\sigma_I$  and resonance constants  $\sigma_R$  by a method<sup>20</sup> exploiting the  $^{19}\text{F}$  NMR shifts of substituted fluorobenzenes. This approach confirmed the inductive electron attracting character of these groups but suggested their conjugation as weak acceptors. Evidently, they can be weakly conjugated in either sense in different molecules.

O.E. thanks the Grant Agency of the Czech Republic for the financial support given with grant No. 203/96/1658. Financial support from MURST, Rome, is gratefully acknowledged by the Italian authors.

Techniques used: Potentiometry, NMR ( $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^{19}\text{F}$ ), microanalysis

Table 1: pK values of substituted benzoic acids (50% ethanol) and the derived  $\sigma$  values

Table 2:  $^{19}\text{F}$  NMR data of fluoro derivatives and the derived  $\sigma$  values


Fig. 1: Mesomeric formulas for the substituents **1a**, **1b**, **1c**

References: 27

Received, 13th May 1998; Accepted, 15th June 1998  
Paper E/8/03585G

## References cited in this synopsis

- G. Sorba, C. Medana, R. Fruttero, C. Cena, A. Di Stilo, U. Galli and A. Gasco, *J. Med. Chem.*, 1997, **40**, 463 and references therein.
- R. Calvino, A. Gasco and A. Leo, *J. Chem. Soc., Perkin Trans. 2*, 1992, 1643.
- V. Vřeteċka, R. Fruttero, A. Gasco and O. Exner, *J. Mol. Struct.*, 1994, **324**, 277.
- J. Shorter, *Pure Appl. Chem.*, 1997, **69**, 2947.
- R. W. Taft, E. Price, I. R. Fox, I. C. Lewis, K. K. Andersen and G. I. Davis, *J. Am. Chem. Soc.*, 1963, **85**, 709, 3146.



\*To receive any correspondence (e-mail: gasco@pharm.unito.it).