

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/280631743>

One-Pot Synthesis of GABA Amides via the Nucleophilic Addition of Amines to 3,3-Disubstituted Cyclopropenes

ARTICLE *in* ORGANIC & BIOMOLECULAR CHEMISTRY · JULY 2015

Impact Factor: 3.56 · DOI: 10.1039/C5OB01462J

READS

18

3 AUTHORS, INCLUDING:



Vladimir Maslivets

University of Kansas

1 PUBLICATION 0 CITATIONS

SEE PROFILE



Michael Rubin

University of Kansas

85 PUBLICATIONS 2,068 CITATIONS

SEE PROFILE

COMMUNICATION



Cite this: *Org. Biomol. Chem.*, 2015, **13**, 8993

Received 16th July 2015,

Accepted 29th July 2015

DOI: 10.1039/c5ob01462j

www.rsc.org/obc

One-pot synthesis of GABA amides *via* the nucleophilic addition of amines to 3,3-disubstituted cyclopropenes†

Vladimir A. Maslivets,^a Marina Rubina^a and Michael Rubin^{*a,b}

A one-pot synthesis of various GABA amides has been demonstrated, employing the nucleophilic addition of primary and secondary amines across the double bond of cyclopropene-3-carboxamides, followed by ring-opening of the resulting donor–acceptor cyclopropanes and subsequent *in situ* reduction of enamine (imine) intermediates.

γ -Aminobutyric acid (GABA) is the chief inhibitory neurotransmitter in the mammalian central nervous system playing a principal role in reducing neuronal excitability,¹ and is a species of immense importance for modern bioorganic and medicinal chemistry. This motif is omnipresent in natural products, including Bistramide A² and cyclic oligopeptides Microsclerodermins³ and Unguisins.⁴ GABA derivatives are also widely used in numerous over-the-counter and prescription medicinal agents, such as Noofen (Phenibut),⁵ Picamilon,⁶ Lioresal (Baclophen), Lyrica (Pregabalin)⁷ or anti-arthritis drug Trocade (Cipemastat)⁸ (Fig. 1).

In our studies⁹ of donor–acceptor cyclopropanes (DAC),¹⁰ we investigated the possibility to access substituted GABA derivatives *5* *via* the ring opening of DAC **1** (Scheme 1). The propensity of DACs toward ring cleavage is proportional to polarization of the C–C bond between electron-donating (EDG) and electron-withdrawing (EWG) groups. The requisite polarization is commonly achieved through installation of strong EWGs, typically two ester groups, additionally activated by a Lewis acid (“pull” strategy), which leads to products with an “extra” carboxylate moiety at the α -carbon. In our recent report⁹ we proposed the possibility to employ an alternative “push” strategy by taking advantage of our formal nucleophilic substitution methodology that allows for installation of





Fig. 1 Biologically active GABA derivatives.



^aDepartment of Chemistry, University of Kansas, 1251 Wescoe Hall Dr., Lawrence, KS 66045-7582, USA. E-mail: mrubin@ku.edu; Fax: +1(785) 864-5396; Tel: +1 (785) 864-5071

^bDepartment of Chemistry, North Caucasus State University, 1a Pushkin St., Stavropol 355009, Russian Federation

†Electronic supplementary information (ESI) available: Experimental procedures, physico-chemical and spectral data. See DOI: 10.1039/c5ob01462j

various *N*-moieties in the cyclopropane ring. Herein we demonstrate the proof of concept and employment of this strategy toward synthesis of GABA amide derivatives.

amination step (entry 2). In contrast, cyclic secondary amines, such as pyrrolidine (**8c**), morpholine (**8d**), *N*-ethyl- (**8e**), and *N*-benzylpiperazines (**8f**) afforded GABA amides **5ad–5af** in good yields (entries 3–6). *N,N*-Diisopropyl-1-methylcycloprop-2-ene-1-carboxamide (**7b**)¹⁵ and (1-methylcycloprop-2-en-1-yl)(pyrrolidin-1-yl)methanone (**7c**)¹⁵ proved to be similarly efficient as **7a** with a number of secondary amines (entries 7–12). Reaction with primary amines, such as phenethylamine (**8g**), benzylamine (**8h**), and *n*-butylamine (**8j**), also proceeded uneventfully, although somewhat more sluggishly (entries 13, 14, 16–19). It was also necessary to raise the temperature to 140 °C to drive the reaction with aniline (**8i**) to complete conversion (entry 15).

Finally, a possibility to induce a diastereoselective ring cleavage upon addition of chiral amines was probed by reacting cyclopropene **7a** with α -phenylethylamine (**8k**). Unfortunately, transfer of asymmetric information from a remote stereogenic center was not efficient, and the corresponding adduct **5ak** was produced as a 1 : 1 mixture of two diastereomers (entry 20).

Conclusions

In conclusion, we have successfully employed an alternative “push” strategy for ring opening of “push–pull” cyclopropanes generated *in situ* via the unassisted nucleophilic addition of electron-rich amines across the double bond of cyclopropene-3-carboxamides. This concept was utilized in efficient one-pot synthesis of GABA derivatives. Further investigations of this transformation are currently underway in our laboratories, which include (a) exploring the possibility of controlling the diastereoselectivity of small ring cleavage upon addition of chiral amines, assisted by Lewis acidic chelating agents; (b) investigating regio- and stereoselectivity of ring cleavage in 1,3,3-trisubstituted chiral cyclopropenes en route to chiral GABA derivatives possessing several contiguous stereogenic centers; (c) intercepting imine species **3** in diastereoselective reactions with C-nucleophiles. The results on these studies will be reported in due course.

This work was financed through the International Collaboration Program, supported by the Ministry of Education and Science of the Russian Federation and the Ministry of Education of Perm Krai. We are also grateful for the financial support by the Russian Foundation for Basic Research (grant #15-03-02661). Support for NMR instruments used in this project was provided by NIH Shared Instrumentation Grant #S10RR024664 and NSF Major Research Instrumentation Grant #0329648.

Notes and references

- (a) W. Sieghart, *Pharmacol. Rev.*, 1995, **47**, 181; (b) Y. Ben-Ari, *Nat. Rev. Neurosci.*, 2002, **3**, 728; (c) D. F. Owens and A. R. Kriegstein, *Nat. Rev. Neurosci.*, 2002, **3**, 715.
- (a) A. V. Statsuk, D. Liu and S. A. Kozmin, *J. Am. Chem. Soc.*, 2004, **126**, 9546; (b) M. T. Crimmins and A. C. DeBaillie, *J. Am. Chem. Soc.*, 2006, **128**, 4936; (c) J. T. Lowe, I. E. Wrona and J. S. Panek, *Org. Lett.*, 2007, **9**, 327.
- (a) C. A. Bewley, C. Debitus and D. J. Faulkner, *J. Am. Chem. Soc.*, 1994, **116**, 7631; (b) J. Zhu and D. Ma, *Angew. Chem., Int. Ed.*, 2003, **42**, 5348.
- L. Hunter and J. H. Chung, *J. Org. Chem.*, 2011, **76**, 5502.
- I. Lapin, *CNS Drug Rev.*, 2001, **7**, 471.
- O. V. Denisenko, O. A. Shandra, L. M. Karpov and L. I. Siomik, *Neurophysiology*, 2014, **46**, 284.
- (a) C. P. Taylor, T. Angelotti and E. Fauman, *Epilepsy Res.*, 2007, **73**, 137; (b) J. S. Bryans and D. J. Wustrow, *Med. Res. Rev.*, 1999, **19**, 149.
- (a) N. Borkakoti, *Biochem. Soc. Trans.*, 2004, **32**, 17; (b) F. J. Hemmings, M. Farhan, J. Rowland, L. Banken and R. Jain, *Rheumatology*, 2001, **40**, 537.
- J. E. Banning, J. Gentillon, P. G. Ryabchuk, A. R. Prosser, A. Rogers, A. Edwards, A. Holtzen, I. A. Babkov, M. Rubina and M. Rubin, *J. Org. Chem.*, 2013, **78**, 7601.
- For recent reviews, see: (a) T. F. Schneider, J. Kaschel and D. B. Werz, *Angew. Chem., Int. Ed.*, 2014, **53**, 5504; (b) F. de Nanteuil, F. De Simone, R. Frei, F. Benfatti, E. Serrano and J. Waser, *Chem. Commun.*, 2014, **50**, 10912; (c) S. Liao, X. Sun and Y. Tang, *Acc. Chem. Res.*, 2014, **47**, 2260; (d) H. K. Grover, M. R. Emmett and M. A. Kerr, *Org. Biomol. Chem.*, 2015, **13**, 655.
- (a) P. Ryabchuk, A. Edwards, N. Gerasimchuk, M. Rubina and M. Rubin, *Org. Lett.*, 2013, **15**, 6010; (b) P. Ryabchuk, M. Rubina, J. Xu and M. Rubin, *Org. Lett.*, 2012, **14**, 1752; (c) A. R. Prosser, J. E. Banning, M. Rubina and M. Rubin, *Org. Lett.*, 2010, **12**, 3968; (d) B. K. Alnasleh, W. M. Sherrill, M. Rubina, J. Banning and M. Rubin, *J. Am. Chem. Soc.*, 2009, **131**, 6906.
- For ring-retentive nucleophilic addition of amines to cyclopropenes, see: (a) E. I. Gritsenko, R. R. Khaliullin, V. V. Plemenkov and E. M. Faizullin, *Zh. Obshch. Khim.*, 1988, **58**, 2733; (b) M. Franck-Neumann, M. Miesch and H. Kempf, *Tetrahedron*, 1988, **44**, 2933; (c) R. M. Albert and G. B. Butler, *J. Org. Chem.*, 1977, **42**, 674.
- For other examples of nucleophilic displacement of halogen in cyclopropyl halides with nitrogen-based nucleophiles, see: (a) S. Y. Kang, S.-H. Lee, H. J. Seo, M. E. Jung, K. Ahn, J. Kim and J. Lee, *Bioorg. Med. Chem. Lett.*, 2008, **18**, 2385; (b) K. N. Shavrin, V. D. Gvozdev and O. M. Nefedov, *Russ. Chem. Bull.*, 2010, **59**, 396; (c) G. S. Basarab, P. Hill, C. J. Eyermann, M. Gowravaram, H. Kack and E. Osimoni, *Bioorg. Med. Chem. Lett.*, 2012, **22**, 5600; (d) T. H. Walls, S. C. Grindrod, D. Beraud, L. Zhang, A. R. Baheti, S. Dakshanamurthy, M. K. Patel, M. L. Brown and L. H. MacArthur, *Bioorg. Med. Chem.*, 2012, **20**, 5269; (e) Y. Zhu and Y. Gong, *J. Org. Chem.*, 2015, **80**, 490.
- J. E. Banning, A. R. Prosser, B. K. Alnasleh, J. Smarker, M. Rubina and M. Rubin, *J. Org. Chem.*, 2011, **76**, 3968.
- (a) W. M. Sherrill, R. Kim and M. Rubin, *Synthesis*, 2009, 1477; (b) R. Kim, W. M. Sherrill and M. Rubin, *Tetrahedron*, 2010, **66**, 4947.