

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/46287123>

# Rapid Reaction of Hydrogen Sulfide with the Neutrophil Oxidant Hypochlorous Acid to Generate Polysulfides

ARTICLE in CHEMICAL RESEARCH IN TOXICOLOGY · SEPTEMBER 2010

Impact Factor: 3.53 · DOI: 10.1021/tx100266a · Source: PubMed

---

CITATIONS

48

---

READS

38

## 2 AUTHORS:



Péter Nagy

National Institute of Oncology

56 PUBLICATIONS 991 CITATIONS

SEE PROFILE



Christine Winterbourn

University of Otago

362 PUBLICATIONS 19,704 CITATIONS

SEE PROFILE

# Rapid Reaction of Hydrogen Sulfide with the Neutrophil Oxidant Hypochlorous Acid to Generate Polysulfides

Péter Nagy\*<sup>†</sup> and Christine C. Winterbourn

Department of Pathology, University of Otago Christchurch, P.O. Box 4345, Christchurch, New Zealand


Received August 4, 2010

H<sub>2</sub>S has been recognized as a signaling molecule and mediator of inflammation. Here, we report the kinetics and mechanism of its reaction with the neutrophil oxidant hypochlorous acid. Stopped flow studies, carried out at high pH, showed this reaction to be extremely fast, with a second-order rate constant extrapolated to be  $2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  at pH 7.4. The reaction produces polysulfides rather than polythionates and may represent a novel pathway for protein Cys-sulphydration, a recently proposed mechanism for H<sub>2</sub>S signaling.

Hydrogen sulfide has been shown to be produced in almost all tissues of the human body via trans-sulfuration pathways during the metabolism of cysteine by two pyridoxal phosphate-dependent enzymes: *cystathione-γ-lyase* (CSE) and *cystathione-β-synthetase* (CBS) (1). H<sub>2</sub>S is currently under active investigation to elucidate its biological functions and explore its therapeutic potential (2). It acts as a vasorelaxant as demonstrated by hypertension in CSE-deleted mice (3), it is proposed to increase the lifespan and thermotolerance in *Caenorhabditis elegans* (4), and it is capable of inducing hibernation in mice by decreasing oxygen consumption and reducing the metabolic rate (5). Along with nitric oxide and carbon monoxide, it is now regarded as a physiological gasotransmitter. Although many investigations suggest that, unlike NO or CO, it acts primarily via opening K<sub>ATP</sub> channels (6), a recent report proposed that in analogy with NO-mediated nitrosothiol formation, H<sub>2</sub>S signals through the formation of protein persulfides (ProteinCys-SSH) (3).

The chemistry of the reactions of H<sub>2</sub>S with biomolecules is poorly understood. For example, its cardioprotective (7) and neuroprotective (8, 9) effects were associated with its ability to scavenge physiological oxidants, but the kinetics and products of these reactions were not studied, and the physiological concentrations of H<sub>2</sub>S are controversial. In this contribution, we have investigated the reaction of H<sub>2</sub>S with hypochlorous acid (HOCl). HOCl is a neutrophil-derived oxidant that is thought to be primarily responsible for clearing invading microbes and has a major role in inflammation (10). It is produced in the myeloperoxidase-catalyzed oxidation of chloride by H<sub>2</sub>O<sub>2</sub> (11). Of particular relevance to this study, the mediatory role of H<sub>2</sub>S in leukocyte action is increasingly recognized (12).

Mixing OCl<sup>−</sup> with HS<sup>−</sup> resulted in a marked change in the UV-spectrum; see Figure S1 (Supporting Information). The largest absorbance change was an increase at 290 nm, and therefore, the kinetics of the reaction was followed at this wavelength. At physiological pH, the reaction was too fast to follow, but it could be monitored by stopped-flow spectroscopy at high OH<sup>−</sup> and relatively low reactant concentrations. Pseudo first-order conditions were employed by keeping the concentration of HS<sup>−</sup> at a large excess over [OCl<sup>−</sup>]. The observed exponential kinetic traces (see Figure S2a, Supporting Informa-



**Figure 1.** Hydroxide ion concentration dependency of the rate of the reaction of HS<sup>−</sup> with OCl<sup>−</sup> under pseudo first-order conditions. Data points represent the effective rate constants (that were calculated by dividing the obtained pseudo first-order rate constants by [HS<sup>−</sup>]) at different hydroxide ion concentrations. The dashed line represents the linear least-squares fit of the data.

tion) indicate that the rate law exhibits first order dependency on the concentration of OCl<sup>−</sup>. This was corroborated by the fact that the observed pseudo first-order rate constants remained the same when changing the concentration of OCl<sup>−</sup> at constant pH, [HS<sup>−</sup>], and ionic strength (not shown). Increasing the concentration of HS<sup>−</sup> resulted in a linear increase in the observed rate constants (Figure S2b, Supporting Information) indicating that the reaction is also first order for [HS<sup>−</sup>].

The rate of the reaction was inversely dependent on [OH<sup>−</sup>] (Figure 1). This indicates that HOCl is more reactive than OCl<sup>−</sup> and is consistent with analogous reactions of the hypohalites and with the protonated form of the oxidant being a better electrophile (13–16). However, the fact that the line on Figure 1 has a positive intercept indicates that OCl<sup>−</sup> reacts with a measurable rate under these conditions. The following mechanism is consistent with these observations:



where  $K_w$  is the ionic product of water, and  $K_a^{\text{HOCl}}$  is the acid dissociation constant of HOCl ( $\text{p}K_a^{\text{HOCl}} = 7.4$ ). Using the pre-

\* To whom correspondence should be addressed. E-mail: peter.nagy@otago.ac.nz; peter.nagy@oncol.hu (after 2010).

<sup>†</sup> Current address: Department of Molecular Toxicology, National Institute of Oncology, 1122 Budapest, Rath Gyorgy 7-9, Hungary.



equilibrium approximation for reaction 1, the following rate equation can be derived from this model:

$$-\frac{d[\text{HS}^-]}{dt} = \left( k_3 + \frac{k_2 K_1}{[\text{OH}^-]} \right) [\text{OCl}^-][\text{HS}^-] \quad (4)$$


On the basis of this rate equation,  $k_{\text{eff}}$  in Figure 1 will be defined as  $k_{\text{eff}} = (k_2 K_1 / [\text{OH}^-] + k_3)$ . Because an excess of  $[\text{HS}^-]$  was used over  $[\text{OCl}^-]$ , it is not necessary to correct  $k_2$  or  $k_3$  with a proportionality constant (17). Therefore,  $k_3 = (2.7 \pm 0.2) \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$  can be obtained from the intercept and  $k_2 = (4.8 \pm 0.1) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  from the slope of the line of Figure 1 using  $\text{p}K_{\text{w}} = 13.7$  and  $\text{p}K_{\text{a}}^{\text{HOCl}} = 7.4$ . These rate constants are close to the analogous rate constants for  $\text{OCl}^-$  and  $\text{HOCl}$  reacting with the thiolate form of cysteine:  $1.9 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$  and  $1.2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ , respectively (14, 18). The sulfhydryl group of cysteine is mostly protonated at physiological pH (19). However, the two  $\text{p}K_{\text{a}}$  values of  $\text{H}_2\text{S}$  are 6.76 and 19.2, respectively (20), which means that at pH 7.4, 81% would be present as  $\text{HS}^-$  and 19% as  $\text{H}_2\text{S}$ . The negative charge on the sulfur in  $\text{HS}^-$  makes it a better nucleophile compared to  $\text{H}_2\text{S}$ , and therefore, it is the more reactive form toward electrophiles. The  $\text{S}^{2-}$  form, which would be an even better nucleophile, has very little abundance at  $\text{pH} < 14$ .

On the basis of the  $[\text{OH}^-]$  dependency at high pH, the reaction of  $\text{HOCl}$  with  $\text{HS}^-$  is 5 orders of magnitude faster than that of  $\text{OCl}^-$ . If we assume that  $\text{H}_2\text{S}$  has negligible reactivity with  $\text{HOCl}$  at pH 7.4 and that the reaction proceeds predominantly via  $\text{HS}^-$  and  $\text{HOCl}$ , then the lower limit of the apparent rate constant between the two reactants at pH 7.4 can be extrapolated to the value of  $2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ .

We also investigated the products of the reaction. The peaks at 290 and 380 nm in the UV-spectrum at pH 7 suggest the formation of polysulfides (21). We found a linear correlation between  $[\text{OCl}^-]_0$  and the observed absorbance change at both pH 7.4 and pH 13, with the linear fits passing through the origin (Figure S3, Supporting Information). This would be consistent with a mechanism analogous to that described for cysteine (14) in which the initial oxidation step is followed by the formation of a disulfide ( $\text{HS-S}^-$ ) as in reactions 5–9.



Alternatively, reactions 6 and 7 could proceed via  $\text{OH}^+$  transfer to directly give  $\text{HSOH}$ . However, the  $\text{Cl}^+$  transfer mechanism would be consistent with a large body of literature on the two electron oxidation of nucleophiles with hypohalous acids; see refs 14–19 and the references therein. The disulfide species can undergo further exchange reactions to give polysulfides (reaction 10), where the number of atoms of zerovalent sulfur per polysulfide can vary between 1 and 8 depending on



**Figure 2.** Qualitative and quantitative analysis of the product(s) of the reaction of  $\text{HS}^-$  with  $\text{OCl}^-$ .  $\text{OCl}^-$  (2.5 mM) was reacted with 150 mM  $\text{HS}^-$  at pH 7.4 (in 50 mM phosphate buffer). The reaction mixture was acidified by the addition of  $\text{H}_3\text{PO}_4$  to a final pH of  $\sim 2$ . After 2 h of incubation in the dark at room temperature, the yellow precipitate was extracted into  $\text{CHCl}_3$ , and the dashed line represents the spectrum of the organic phase. The solid line shows the UV-spectrum of 0.5 mM authentic cyclooctasulfur in chloroform. On the basis of the similarities of the two spectra and a previous qualitative analysis of a similar precipitate by tandem mass spectrometry (22), we conclude that acidification of the reaction mixture resulted in the precipitation of cyclooctasulfur. Because of a 5-fold dilution during extraction, the solid line represents the expected amount of  $\text{S}_8$  formation upon a 100% conversion of  $\text{OCl}^-$  to  $\text{S}_8$  in the above experiment. The similar absorbance of the two samples suggests that  $\text{OCl}^-$  generates predominantly polysulfides rather than polythionates when reacted with an excess of  $\text{HS}^-$  (for further details see Supporting Information).

the pH and the relative amount of the oxidized versus reduced forms (21).



Under the conditions of our kinetic analysis (high pH and an excess of  $[\text{HS}^-]$  over the oxidant), the dominant form (>99%) is the disulfide (21). However, at pH 7.4 in a 1 to 1 mixture of  $\text{HS}^-$  and  $\text{S}_0$ , the two dominant species would be  $\text{HS}_4\text{S}^-$  ( $\sim 60\%$ ) and  $\text{HS}_3\text{S}^-$  ( $\sim 40\%$ ) (21).

As reactions 8 and 9 are fast (compared to reactions 6 and 7) at  $\text{pH} > 13$  (see Stopped-Flow Studies in Supporting Information), we have no information on the nature or the rate of these reactions or the protonation state of the intermediates. However, these rates could be smaller than that of reaction 6 at physiological pH, where reaction 6 becomes very fast.

Polysulfides are relatively stable at higher or neutral pH, but under acidic conditions, they precipitate from aqueous solution as elemental sulfur.




Upon acidification of our reaction mixtures, we observed the formation of a yellow precipitate. The precipitate dissolved in chloroform, resulting in an UV-spectrum analogous to that of cyclooctasulfur (Figure 2). The amount of precipitated  $\text{S}_8$  at different  $[\text{OCl}^-]$  was quantified using a previously reported procedure (22) (see Supporting Information and Figure 2). At an excess of  $[\text{HS}^-]$  over  $[\text{OCl}^-]$ , the majority of the oxidizing equivalents were converted to elemental sulfur, suggesting a minor contribution (if at all) of pathways that would result in the formation of polythionates (oxysulfur species, such as  $\text{S}_2\text{O}_3^{2-}$ ,  $\text{S}_4\text{O}_6^{2-}$ ,  $\text{SO}_3^{2-}$ , or  $\text{SO}_4^{2-}$ ).


Although the calculated second order rate constant of the reaction of  $\text{HS}^-$  with  $\text{HOCl}$  at pH 7.4 means the reaction is highly favorable, whether  $\text{H}_2\text{S}$  acts as an antioxidant toward  $\text{HOCl}$  (9, 23) will depend on its local concentration. Different authors have reported widely varying values for physiological

### Scheme 1. Proposed Models for the Formation of Protein Persulfides

#### Model A



#### Model B



Protein persulfides were generated by (A) direct oxidation of the protein Cys thiol or (B) direct oxidation of  $\text{H}_2\text{S}$ .

concentrations of  $\text{H}_2\text{S}$ . However, the most recent studies favor a relatively low value (24), suggesting that  $\text{H}_2\text{S}$  has little importance as an antioxidant. However,  $\text{HOCl}$  could deplete  $\text{H}_2\text{S}$  at sites of inflammation, possibly producing bioactive products and influencing its signaling function. For example, it has recently been proposed that  $\text{H}_2\text{S}$  signals via protein sulfhydrylation (25) to give persulfide derivatives on Cys residues. To form a persulfide from a thiol and  $\text{H}_2\text{S}$ , one molar oxidizing equivalent is needed. The oxidant could either react with the Cys to give a sulfenic acid, which could react further with  $\text{HS}^-$  to give the persulfide (model A, Scheme 1). Alternatively, the oxidant could target  $\text{HS}^-$  directly to give polysulfides, as shown in this study, followed by a reaction with the Cys thiol to give the persulfide (model B, Scheme 1).

**Acknowledgment.** This work was supported by the Marsden Fund, and the equipment used was provided by the National Research Centre for Growth and Development. We are grateful to Dr. Guy Jameson who allowed us to use his stopped-flow apparatus and Prof. Anthony Kettle for helpful comments on the manuscript.

**Supporting Information Available:** Experimental details, concentration dependencies of the rate law, spectral titration, and quantitative analyses of the product(s). This material is available free of charge via the Internet at <http://pubs.acs.org>.

### References

- (1) Stipanuk, M. H., and Beck, P. W. (1982) Characterization of the enzymic capacity for cysteine desulfhydration in liver and kidney of the rat. *Biochem. J.* 206, 267–277.
- (2) Szabo, C. (2007) Hydrogen sulphide and its therapeutic potential. *Nat. Rev. Drug Discovery* 6, 917–935.
- (3) Yang, G. D., Wu, L. Y., Jiang, B., Yang, W., Qi, J. S., Cao, K., Meng, Q. H., Mustafa, A. K., Mu, W. T., Zhang, S. M., Snyder, S. H., and Wang, R. (2008)  $\text{H}_2\text{S}$  as a physiologic vasorelaxant: Hypertension in mice with deletion of cystathionine gamma-lyase. *Science* 322, 587–590.
- (4) Miller, D. L., and Roth, M. B. (2007) Hydrogen sulfide increases thermotolerance and lifespan in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. U.S.A.* 104, 20618–20622.
- (5) Blackstone, E., Morrison, M., and Roth, M. B. (2005)  $\text{H}_2\text{S}$  induces a suspended animation-like state in mice. *Science* 308, 518–518.
- (6) Zhao, W. M., Zhang, J., Lu, Y. J., and Wang, R. (2001) The vasorelaxant effect of  $\text{H}_2\text{S}$  as a novel endogenous gaseous K-ATP channel opener. *EMBO J.* 20, 6008–6016.
- (7) Elrod, J. W., Calvert, J. W., Morrison, J., Doeller, J. E., Kraus, D. W., Tao, L., Jiao, X. Y., Scalia, R., Kiss, L., Szabo, C., Kimura, H., Chow, C. W., and Lefer, D. J. (2007) Hydrogen sulfide attenuates myocardial ischemia-reperfusion injury by preservation of mitochondrial function. *Proc. Natl. Acad. Sci. U.S.A.* 104, 15560–15565.
- (8) Kimura, Y., and Kimura, H. (2004) Hydrogen sulfide protects neurons from oxidative stress. *FASAB J.* 18, 1165–1166.
- (9) Whiteman, M., Cheung, N. S., Zhu, Y. Z., Chu, S. H., Siau, J. L., Wong, B. S., Armstrong, J. S., and Moore, P. K. (2005) Hydrogen sulphide: a novel inhibitor of hypochlorous acid-mediated oxidative damage in the brain? *Biochem. Biophys. Res. Commun.* 326, 794–798.
- (10) Hampton, M. B., Kettle, A. J., and Winterbourn, C. C. (1998) Inside the neutrophil phagosome: oxidants, myeloperoxidase, and bacterial killing. *Blood* 92, 3007–3017.
- (11) Davies, M. J., Hawkins, C. L., Pattison, D. I., and Rees, M. D. (2008) Mammalian heme peroxidases: from molecular mechanisms to health implications. *Antioxid. Redox Signaling* 10, 1199–1234.
- (12) Zhang, H. L., and Bhatia, M. (2008) Hydrogen sulfide: A novel mediator of leukocyte activation. *Immunopharmacol. Immunotoxicol.* 30, 631–645.
- (13) Nagy, P., and Ashby, M. T. (2005) Reactive sulfur species: kinetics and mechanism of the oxidation of cystine by hypochlorous acid to give N,N'-dichlorocystine. *Chem. Res. Toxicol.* 18, 919–923.
- (14) Nagy, P., and Ashby, M. T. (2007) Reactive sulfur species: kinetics and mechanisms of the oxidation of cysteine by hypochlorous acid to give cysteine sulfenic acid. *J. Am. Chem. Soc.* 129, 14082–14091.
- (15) Nagy, P., and Ashby, M. T. (2007) Kinetics and mechanism of the oxidation of the glutathione dimer by hypochlorous Acid and catalytic reduction of the chloroamine product by glutathione reductase. *Chem. Res. Toxicol.* 20, 79–87.
- (16) Nagy, P., Beal, J. L., and Ashby, M. T. (2006) Thiocyanate is an efficient endogenous scavenger of the phagocytic killing agent hypobromous acid. *Chem. Res. Toxicol.* 19, 587–593.
- (17) Nagy, P., Jameson, G. N., and Winterbourn, C. C. (2009) Kinetics and mechanisms of the reaction of hypothiocyanous acid with 5-thio-2-nitrobenzoic acid and reduced glutathione. *Chem. Res. Toxicol.* 22, 1833–1840.
- (18) Armesto, X. L., Canle, M., Fernandez, M. I., Garcia, M. V., and Santabell, J. A. (2000) First steps in the oxidation of sulfur-containing amino acids by hypohalogenation: very fast generation of intermediate sulfonyl halides and halosulfonium cations. *Tetrahedron* 56, 1103–1109.
- (19) Nagy, P., and Winterbourn, C. C. (2010) Redox Chemistry of Biological Thiols, in *Advances in Molecular Toxicology* (Fishbein, J. C., Ed.) pp 183–222, Elsevier: New York.
- (20) Hughes, M. N., Centelles, M. N., and Moore, K. P. (2009) Making and working with hydrogen sulfide: The chemistry and generation of hydrogen sulfide in vitro and its measurement in vivo: a review. *Free Radical Biol. Med.* 47, 1346–1353.
- (21) Giggenbach, W. (1972) Optical-spectra and equilibrium distribution of polysulfide ions in aqueous-solution at 20 degrees. *Inorg. Chem.* 11, 1201–1207.
- (22) Nagy, P., Wang, X., Lemma, K., and Ashby, M. T. (2007) Reactive sulfur species: Hydrolysis of hypothiocyanite to give thiocarbamate-S-oxide. *J. Am. Chem. Soc.* 129, 15756–15757.
- (23) Laggner, H., Muellner, M. K., Schreier, S., Sturm, B., Hermann, M., Exner, M., Gmeiner, B. M. K., and Kapiotis, S. (2007) Hydrogen sulphide: A novel physiological inhibitor of LDL atherogenic modification by  $\text{HOCl}$ . *Free Radical Res.* 41, 741–747.
- (24) Whitfield, N. L., Kreimier, E. L., Verdial, F. C., Skovgaard, N., and Olson, K. R. (2008) Reappraisal of  $\text{H}_2\text{S}$  /sulfide concentration in vertebrate blood and its potential significance in ischemic preconditioning and vascular signaling. *Am. J. Physiol.-Regul. Integr. Comp. Physiol.* 294, R1930–R1937.
- (25) Mustafa, A. K., Gadalla, M. M., Sen, N., Kim, S., Mu, W., Gazi, S. K., Barrow, R. K., Yang, G., Wang, R., and Snyder, S. H. (2009)  $\text{H}_2\text{S}$  signals through protein S-sulfhydration. *Sci. Signal.* 2, ra72.

TX100266A