

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/239938538>

# Synthesis of hydrolytically stable porphyrin C- and S-glycoconjugates in high yields

ARTICLE in CHEMICAL COMMUNICATIONS · JANUARY 2001

Impact Factor: 6.83 · DOI: 10.1039/b008489l

CITATIONS

47

READS

34

4 AUTHORS, INCLUDING:



Charles Michael Drain

City University of New York - Hunter College

118 PUBLICATIONS 3,974 CITATIONS

SEE PROFILE



Xin Chen

CUNY Graduate Center

16 PUBLICATIONS 413 CITATIONS

SEE PROFILE



Richard W Franck

City University of New York - Hunter College

160 PUBLICATIONS 3,555 CITATIONS

SEE PROFILE

# Synthesis of hydrolytically stable porphyrin C- and S-glycoconjugates in high yields†

Paolo Pasetto, Xin Chen, Charles Michael Drain\* and Richard W. Franck\*

Department of Chemistry and Biochemistry, Hunter College and Graduate Center of The City University of New York, 695 Park Avenue, New York, NY 10021, USA. E-mail: cdrain@shiva.hunter.cuny.edu; rfranck@shiva.hunter.cuny.edu; Phone: (212)650-3791; Fax: (212)772-5332

Received (in Corvallis, OR, USA) 18th October 2000, Accepted 14th November 2000

First published as an Advance Article on the web 14th December 2000

**Two C-glucosyl porphyrins are prepared using Ramberg–Backlund and Lindsey methods for the key conversions, and thiosugars are shown to react with perfluorophenylporphyrins.**

Photofrin®, a mixture of hematoporphyrin oligomers, is currently used clinically for the photodynamic therapy (PDT) of cancers, but suffers from a variety of problems including solubility and dosing.<sup>1</sup> Because of their potential selective binding to various cell types, porphyrins appended with a variety of saccharides have been examined as possible agents for PDT.<sup>1,2</sup> Most recently, sugar-specific binding to rat hepatoma cells by porphyrin glycoconjugates has been described.<sup>2</sup> Their efficacy as antibiotics and anti-viral agents is also under intense investigation.<sup>1</sup> Since metalloporphyrins are well-established catalysts, the attachment of sugars to effect regio- and stereo-selective oxidations has also been examined, albeit with limited success.<sup>3</sup> Additionally, there are several requirements for the successful commercialization of any of these adducts: (a) effective synthesis in high yields, (b) stable products, in this case to hydrolysis, and (c) highly pure compounds.

It is interesting that with few exceptions,<sup>4–6</sup> the glycoconjugate has been oxygen linked *via* a glycosidic bond to a phenolic aryl porphyrin. When the syntheses of these O-linked materials are examined, one finds that disappointingly low yields are reported, considering that the Lindsey porphyrin synthesis yields are typically 45–65%.<sup>7</sup> We believe that one problem may be that the frequently used *O*-glycosyloxybenzaldehyde starting materials, when subjected to the Lindsey porphyrin synthesis using BF<sub>3</sub> catalysis, can undergo a competing and yield-reducing Suzuki *O* to *C*-glycoside rearrangement.<sup>8</sup> Additionally, a perceived problem with the *O*-glycosides is the inherent possibility for glycosyl cleavage by acids, and in biological systems by glycohydrolases. We wish to describe our preliminary results, which provide possible solutions to both the issues of poor yields and hydrolytic instability, namely the preparation of *C*- and *S*-glycosylated aryl porphyrins in high yields. Both yield and stability are crucial factors for the successful incorporation of sugar moieties into porphyrin combinatorial libraries, since the linkage must be stable both during the formation, subsequent reactions, and in the presence of a variety of functional groups.

For the *C*-glycoside series, we chose to exploit earlier work from our laboratory where the Ramberg–Backlund synthesis is used to link sugars to aglycones.<sup>9,10</sup> Thus, after screening several alternative functional equivalents of *p*-bromomethylbenzaldehyde, we settled on  $\alpha,\alpha'$ -dibromo-*p*-xylene as an inexpensive starting material. This was condensed with 1-thio- $\beta$ -D-glucose tetraacetate<sup>11</sup> to afford thioglucoside **1**. Then conversion to the silylated material **2** was accomplished by treatment of **1** with *tert*-butyldimethylsilanol in the presence of silver triflate and 2,6-di-*tert*-butylpyridine.<sup>12</sup> The replacement

of the acetyl groups with benzyl ethers to obtain **3** was achieved by deprotection of **2** with sodium methoxide, followed by benzylation. The sulfide **3** was oxidized to the sulfone **4** with monoperoxyphthalic acid (MMPP). The resulting sulfone was employed in the Ramberg–Bäcklund synthesis of the *exo*-glucal **5** under Chan's conditions.<sup>13</sup> The two isomers (*Z*)-**5** and (*E*)-**5** in the ratio 8:2 were identified by NOE measurements, which showed an effect between H-1 and H-3 in the case of the *E*-isomer. In some cases the intermediate  $\alpha$ -bromosulfone **6** was isolated from the reaction mixture and then converted to the *exo*-glucal **5** by treatment with base. The hydrogenation of **5** with palladium 5% on alumina afforded the  $\beta$ -C-glucoside **7**, identified by the coupling constant  $J = 9.2$  Hz, indicating an anti-diaxial configuration of the anomeric H-2 with respect to H-3. The cleavage of the silyl ether **7**, followed by Swern<sup>14</sup> oxidation gave the aldehyde **9**, which was utilized in the synthesis of the porphyrins **10** and **11** in 15 and 53% yields respectively, under Lindsey conditions.<sup>7</sup> The materials obtained showed the expected spectroscopic data. The hydrogenolysis of the benzylic protecting groups yielded the porphyrins **12** and **13**, without reduction of the porphyrin. *S*-Glycosides were chosen as worthy conjugates because they are considered to be good mimics of *O*-glycosides with enhanced stability toward enzymatic hydrolysis.<sup>15</sup>

The second reaction sequence begins with 5,10,15,20-tetra-(perfluorophenyl)porphyrin which can be routinely made in gram quantities in high yields by the Lindsey procedure.<sup>7</sup> The reactivity of the *para*-fluoro group toward a large variety of nucleophilic substitution reactions has been well established.<sup>4,16</sup> The tetrakis(thiogalactosyl) and tetrakis(thioglucosyl) derivatives of this fluorinated porphyrin are formed in > 85 and 90% yields, respectively, by modifications of previously described procedures.<sup>16</sup> Specifically, the porphyrin is dissolved in amine-free DMF, five equiv. of the sodium salt of the thiosugar added, and the mixture stirred at rt for 4 h. Purification is accomplished on silica-gel using an ethanol–ethyl acetate gradient as eluent.


DNA photocleavage assays are widely used to evaluate the photoreactivity of porphyrin compounds, though DNA is not the primary site of Photofrin activity.<sup>1,5,6</sup> Thus we used standard plasmid photocleavage assays to evaluate the photoreactivity and compare these results with those for other porphyrins. The conditions used to evaluate the photodamage to plasmid DNA caused by porphyrin compounds varies, but our analysis (see

Table 1

Compound	Partition coeff. octanol–water	DNA photocleavage
5-glu-triPP <b>12</b>	68	Fair
Tetra glu PP <b>13</b>	30	Poor
Tetra thio glu F4PP	50	Poor
Tetra thio gal F4PP	38	Fair
DiMePy + MeOHP <sup>a</sup>	7	Excellent

<sup>a</sup> Ref. 17.

† Electronic supplementary information (ESI) available: experimental details. See <http://www.rsc.org/suppdata/cc/b0/b0084891/>



In conclusion, both the C and S linked glycoporphyrins can be synthesized in high yields exceeding 50% based on starting aldehydes using more efficient synthetic strategies. These compounds are stable to hydrolysis and show some photo-

The authors thank Margareta Sorenson and Dr Clifford Soll, and acknowledge the support of NIGMS SCORE Program 1S06GM60654, NSF CAREER 9732950 to C. M. D.; PSC-CUNY awards to C. M. D. and R. W. F.; NIH GM51216 to R. W. F. Chemistry department infrastructure is partially supported by RCMI grant NIH RR03037. R. W. F. dedicates this article to Professor Steven Weinreb on the occasion of his 60th birthday. ©QLT Phototherapeutics Inc.

- 1 R. Bonnett, P. Charlesworth, B. D. Djelal, S. Foley, J. J. McGarvey and T. G. Truscott, *J. Chem. Soc.*, 1999, 325; E. D. Sternberg, C. Bruckner and D. Dolphin, *Tetrahedron*, 1998, **54**, 4151; T. J. Dougherty and D. Kessel, *Rev. Contemp. Pharmacother.*, 1999, **10**, 19; B. R. Munson and R. J. Fiel, *Nucleic Acids Res.*, 1992, **20**, 1315.
- 2 K. Fujimoto, T. Miyata and Y. Aoyama, *J. Am. Chem. Soc.*, 2000, **122**, 3558.
- 3 P. Maillard, J. L. Guerquin-Kern and M. Momenteau, *Tetrahedron Lett.*, 1991, **32**, 4901.
- 4 A thioether link between tetra(tetrafluorophenyl)porphyrin and C-6 of glucose embedded in cyclodextrin has been reported: J. Yang and R. Breslow, *Angew. Chem., Int. Ed.*, 2000, **39**, 2692, and references therein.
- 5 C-glycosides: M. Cornia, M. Menozzi, E. Ragg, S. Mazzini, A. Scarafoni, F. Zanardi and G. Casiraghi, *Tetrahedron*, 2000, **56**, 3977, and references therein; M. Momenteau, P. Maillard, M.-A. De Bélinay, D. Carrez and A. Croisy, *J. Biomed. Opt.*, 1999, **4**, 298; N. Ono, M. Baugauchi and K. Maruyama, *Tetrahedron Lett.*, 1992, **33**, 1629.
- 6 S-glycosides linked to alkyl groups: I. Sylvain, R. Benhaddou, V. Carre, S. Cottaz, H. Driguez, R. Granet, M. Guilloton and P. Krausz, *J. Porphyrins Phthalocyanines*, 1999, **3**, 1 and references therein.
- 7 J. S. Lindsey, I. C. Schreiman, H. C. Hsu, P. C. Kearney and A. M. Marguerettaz, *J. Org. Chem.*, 1987, **52**, 827; F. Li, K. Yang, J. S. Tyhonas, K. A. MacCrum and J. S. Lindsey, *Tetrahedron*, 1997, **53**, 12 339.
- 8 T. Matsumoto, T. Hosoya and K. Suzuki, *Synlett*, 1991, 709; T. Matsumoto, T. Hosoya and K. Suzuki, *Tetrahedron Lett.*, 1990, **31**, 4629.
- 9 P. S. Belica and R. W. Franck, *Tetrahedron Lett.*, 1998, **39**, 8225; G. Yang, R. W. Franck, H.-S. Byun, R. Bittman, P. Samadder and G. Arthur, *Org. Lett.*, 1999, **1**, 2149.
- 10 F. K. Griffin, P. V. Murphy, D. E. Patterson and R. J. K. Taylor, *Tetrahedron Lett.*, 1998, **39**, 8179; M.-L. Alcaraz, F. K. Griffin, D. E. Patterson and R. J. K. Taylor, *Tetrahedron Lett.*, 1998, **39**, 8183; R. J. K. Taylor, F. K. Griffin and D. E. Paterson, *Angew. Chem., Int. Ed.*, 1999, **38**, 2939; A. D. Campbell, D. E. Paterson, T. M. Raynham and R. J. K. Taylor, *Chem. Commun.*, 1999, 1599.
- 11 D. Horton, in *Methods in Carbohydrate Chemistry*, eds. R. L. Whistler and M. L. Wolfrom, Academic Press Inc., New York, 1963, Vol. II, p. 433-437.
- 12 R. M. Burk, T. S. Gac and M. B. Roof, *Tetrahedron Lett.*, 1994, **35**, 8111.
- 13 T. L. Chan, S. Fong, Y. Li, T. O. Man and C. D. Poon, *Chem. Commun.*, 1994, 1771.
- 14 K. Omura and D. Swern, *Tetrahedron*, 1978, **34**, 1651.
- 15 B. D. Johnston and B. M. Pinto, *J. Org. Chem.*, 2000, **65**, 4607, and references therein; J. C. Wilson, M. J. Kiefel, D. I. Angus and M. von Itzstein, *Org. Lett.*, 1999, **1**, 443; for recent examples of thioglycoside stability toward a glycohydrolase.
- 16 S. J. Shaw, K. J. Elgie, C. Edwards and R. W. Boyle, *Tetrahedron Lett.*, 1999, 1595.
- 17 C. M. Drain, X. Gong, V. Ruta, C. E. Soll and P. F. Chicoineau, *J. Comb. Chem.*, 1999, **1**, 286.