

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/9888654>

Concise total synthesis of the aporphine alkaloid 7,7'-bisdehydro-O-methylisopiline by an InCl_3 mediated cycloisomerization reaction

ARTICLE in CHEMICAL COMMUNICATIONS · OCTOBER 2003

Impact Factor: 6.83 · DOI: 10.1039/B306880C · Source: PubMed

CITATIONS

71

READS

25

2 AUTHORS, INCLUDING:



Victor Mamane

University of Strasbourg

62 PUBLICATIONS 1,507 CITATIONS

SEE PROFILE

Concise total synthesis of the aporphine alkaloid 7,7'-bisdehydro-*O*-methylisopiline by an InCl_3 mediated cycloisomerization reaction

Alois Fürstner* and Victor Mamane

Max-Planck-Institut für Kohlenforschung, D-45470 Mülheim/Ruhr, Germany.


E-mail: fuerstner@mpi-muelheim.mpg.de; Fax: +49 208 306 2994; Tel: +49 208 306 2342

Received (in Cambridge, UK) 17th June 2003, Accepted 9th July 2003

First published as an Advance Article on the web 22nd July 2003

A novel InCl_3 mediated cycloisomerization reaction leading to 10-halophenanthrene derivatives constitutes the key step of the first total syntheses of *O*-methyldehydroisopiline **10** and 7,7'-bisdehydro-*O*-methylisopiline **11**, two prototype members of the aporphine family of alkaloids.

As part of our ongoing investigations on metal catalyzed skeletal rearrangements¹ we have recently developed a new entry into highly substituted phenanthrenes and related polycyclic arenes based on the cycloisomerization process depicted in Scheme 1 ($X = \text{H}$, alkyl).² PtCl_2 turned out to be the catalyst of choice, triggering the desired 6-*endo* cyclizations with high selectivity in all but one case.³



Scheme 1 Metal catalyzed cycloisomerization of *ortho*-alkynylated biphenyl derivatives.

Therefore we were surprised to find that the corresponding haloalkyne derivatives ($X = \text{Cl}$, Br) react rather poorly under these conditions. In addition to the expected 10-halophenanthrenes, significant amounts of the corresponding alkenylidene fluorenes are formed *via* the competing 5-*exo*-cyclization pathway. In an attempt to improve on this result, a set of different metal species was screened for catalytic activity. Among them, InCl_3 turned out to be optimal, effecting the desired transformation with good to excellent yields and high selectivity (Table 1).⁴

The resulting 10-halophenanthrenes are ideally suited for further elaboration. This is exemplified by the first total synthesis of *O*-methyldehydroisopiline **10** isolated from the


Table 1 InCl_3 catalyzed synthesis of 10-halophenanthrenes

Entry	R	X	Yield (%) ^a
1	Me	Cl	90
2	Me	Br	77
3	OMe	Cl	90
4	OMe	Br	59 ^b


^a Isolated yield unless stated otherwise. ^b GC yield.

leaves of the annonaceous plant *Gutteria ouregon*,⁵ and its symmetrical dimer **11**, a secondary metabolite of the tropical trees *Polyalthia bullata*⁶ and *Phoenicanthus obliqua*.⁷ These compounds are prototype members of the aporphine family, a large and rapidly growing class of isoquinoline alkaloids endowed with an impressive number of biological activities.⁸

Selective iodination of commercial bromotrimethoxybenzene **3** furnishes compound **4**⁹ which undergoes a selective activation of its C–I bond in the presence of a catalyst formed *in situ* from $\text{Pd}(\text{OAc})_2$ and tri-*o*-tolylphosphine. The resulting organopalladium species reacts with commercial *N*-vinylphthalimide in a standard Heck reaction¹⁰ to afford the corresponding enamide¹¹ which is chemoselectively hydrogenated in the presence of Crabtree's catalyst¹² without damaging the residual bromide function. The resulting compound **5** allows for a subsequent Suzuki coupling¹³ with commercial 2-formylbenzeneboronic acid to give the highly functionalised biphenyl derivative **6** in 94% yield. Conversion of its aldehyde group into the desired bromoalkyne **7** follows standard procedures¹⁴ and sets the stage for the envisaged carbocyclization to form the phenanthrene core. Gratifyingly, this key transformation worked exquisitely well in the presence of InCl_3 in toluene at 80 °C. The phthalimide protecting group in phenanthrene **8** thus formed was cleaved off by hydrazinol-



Scheme 2 Reagents and conditions: [a] I_2 , HgO , CH_2Cl_2 , r.t., 81%; [b] $\text{Pd}(\text{OAc})_2$ cat., $\text{P}(\text{o-tol})_3$ cat., *N*-vinylphthalimide, iPrNEt_2 , MeCN , 100 °C, 57%; [c] $[\text{Ir}(\text{COD})\text{Py}(\text{PCy}_3)]\text{PF}_6$ cat., H_2 (1 atm), CH_2Cl_2 , quant.; [d] 2-formylbenzeneboronic acid, $\text{Pd}(\text{OAc})_2$ cat., $\text{Cy}_2\text{P}(\text{o-biphenyl})$ cat., K_3PO_4 , toluene, 94%; [e] CBr_4 , PPh_3 , CH_2Cl_2 , 0 °C, 88%; [f] DBU, DMSO, 15 °C, 79%; [g] InCl_3 (1 eq.), toluene, 80 °C, 87%; [h] hydrazine, MeOH , reflux, quant.; [i] CuI , CsOAc , DMSO, 71%.



Scheme 3 Reagents and conditions: [a] $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$, tBuNH_2 , MeOH , 86%.

ysis to give compound **9** which is set up for a smooth intramolecular amination reaction in the presence of CuI and CsOAc as the promoters forging the heterocyclic ring.¹⁵ This high yielding step completes the first total synthesis of *O*-methyl-dehydroisopiline **10** (Scheme 2). The spectroscopic data[†] of this prototype 6a,7-dehydroaporphine derivative¹⁶ are in full accord with the proposed structure.¹⁷

Since 6a,7-dehydroaporphines in general are known to behave like enamines,¹⁸ it was anticipated that a selective activation of the 7-position in **10** might be possible, thus allowing direct conversion of this compound to the corresponding symmetrical dimer **11** (Scheme 3). While the use of $\text{PhI}(\text{OAc})_2$, $\text{Hg}(\text{OAc})_2$, I_2 , or air, which were previously recommended for such purposes,¹⁹ was unsuccessful in our hands leading either to no conversion or to a rapid degradation of the starting material, we were pleased to find that a combination of $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ and tBuNH_2 in MeOH effected the desired oxidative coupling in satisfactory yields.²⁰ The spectral data[†] of the resulting 7,7'-bisaporphine derivative **11** are in excellent agreement with those reported in the literature.^{6,17}

In summary, a straightforward entry into the dehydroaporphine series is described based on a highly productive sequence of metal-catalyzed and -mediated transformations relying on $\text{In}(3+)$, $\text{Pd}(0)$, $\text{Ir}(1+)$, $\text{Cu}(1+)$ and $\text{Cu}(2+)$ as the active components. Due to the flexibility inherent to this route and the fact that dehydroaporphines can be further elaborated into a host of other (natural) products, this approach provides ample opportunity for further exploration of this important class of bioactive natural products.

Notes and references

[†] Data of compound **10**: IR (KAP) 3374, 2933, 2832, 1623, 1391, 749 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 3.24 (t, $J = 6.1$ Hz, 2H), 3.47 (t, $J = 6.0$ Hz, 2H), 3.98 (s, 6H), 4.06 (s, 3H), 6.79 (s, 1H), 7.35 (dt, $J = 8.5$, 1.6 Hz, 1H), 7.44 (dt, $J = 7.8$, 1.1 Hz, 1H), 7.58 (dd, $J = 8$, 1.5 Hz, 1H), 9.40 (d, $J = 8.6$, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ 24.1, 40.7, 60.2, 60.9, 61.3, 105.1, 120.4, 121.2, 123.0, 125.1, 125.8, 126.3, 127.1, 133.6, 140.6, 146.4, 148.4, 151.1. MS (EI) m/z (rel. intensity): 309 ($[\text{M}^+]$, 100), 294 (26), 266 (11). Data of compound **11**: ^1H NMR (400 MHz, CDCl_3) δ 3.14–3.34 (m, 8H), 4.00 (s, 6H), 4.06 (s, 6H), 4.14 (s, 6H), 7.15 (dd, $J = 8.2$, 1.3 Hz, 2H), 7.22 (ddd, $J = 8.4$, 6.4, 1.2 Hz, 2H), 7.35 (ddd, $J = 8.8$, 6.4, 1.6, 2H), 9.57 (dd, $J = 8.6$, 0.6 Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 23.9, 40.6, 60.4, 60.9, 61.3, 120.2, 121.9, 123.1, 123.9, 125.5, 126.8, 127.3, 132.7, 139.6, 146.7, 148.6, 151.1. MS (EI) m/z (rel. intensity): 616 ($[\text{M}^+]$, 100), 308 (12), 294 (12).

1 (a) A. Fürstner, H. Szillat, B. Gabor and R. Mynott, *J. Am. Chem. Soc.*, 1998, **120**, 8305; (b) A. Fürstner, H. Szillat and F. Stelzer, *J. Am. Chem.*

- Soc.*, 2000, **122**, 6785; (c) A. Fürstner, F. Stelzer and H. Szillat, *J. Am. Chem. Soc.*, 2001, **123**, 11863.
- 2 A. Fürstner and V. Mamane, *J. Org. Chem.*, 2002, **67**, 6264.
- 3 Reviews on the use of PtCl_2 in organic synthesis: (a) M. Méndez and A. M. Echavarren, *Eur. J. Org. Chem.*, 2002, **15**; (b) G. C. Lloyd-Jones, *Org. Biomol. Chem.*, 2003, **1**, 215; (c) C. Aubert, O. Buisine and M. Malacria, *Chem. Rev.*, 2002, **102**, 813; (d) M. Méndez, V. Mamane and A. Fürstner, *Chemtracts – Org. Chem.*, 2003, **16**, 397; (e) for a pioneering study see: N. Chatani, N. Fukukawa, H. Sakurai and S. Murai, *Organometallics*, 1996, **15**, 901.
- 4 Reviews on InCl_3 in synthesis: (a) P. Cintas, *Synlett*, 1995, 1087; (b) K. K. Chauhan and C. G. Frost, *J. Chem. Soc., Perkin Trans. 1*, 2000, 3015; (c) B. C. Ranu, *Eur. J. Org. Chem.*, 2000, 2347.
- 5 D. Cortes, R. Hocquemiller, M. Lebouef, A. Cavé and C. Moretti, *J. Nat. Prod.*, 1986, **49**, 878.
- 6 J. D. Connolly, M. E. Haque and A. A. Kadir, *Phytochemistry*, 1996, **43**, 295.
- 7 (a) E. M. K. Wijeratne, B. D. Lankananda, Y. Tezuka, T. Nagaoka and A. A. L. Gunatilaka, *J. Nat. Prod.*, 2001, **64**, 1465; (b) for closely related dimers see: S. Kanokmedhakul, K. Kanokmedhakul, D. Yodbuddee and N. Phonkerd, *J. Nat. Prod.*, 2003, **66**, 616.
- 8 H. Guinaudeau, M. Lebouef and A. Cavé, *J. Nat. Prod.*, 1988, **51**, 389 and references cited therein.
- 9 K. Orito, T. Hatakeyama, M. Takeo and H. Sugimoto, *Synthesis*, 1995, 1273.
- 10 (a) I. Beletskaya and A. V. Cheprakov, *Chem. Rev.*, 2000, **100**, 3009; (b) A. de Meijere and F. E. Meyer, *Angew. Chem.*, 1994, **106**, 2473; A. de Meijere and F. E. Meyer, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 2379.
- 11 Some α -arylation product is also formed which is easily removed by flash chromatography.
- 12 R. H. Crabtree, H. Felkin, T. Fellebeen-Khan and G. E. Morris, *J. Organomet. Chem.*, 1979, **168**, 183.
- 13 (a) J. P. Wolfe, R. A. Singer, B. H. Yang and S. L. Buchwald, *J. Am. Chem. Soc.*, 1999, **121**, 9550; (b) review: A. Suzuki, *J. Organomet. Chem.*, 1999, **576**, 147.
- 14 (a) E. J. Corey and P. L. Fuchs, *Tetrahedron Lett.*, 1972, 3769; (b) V. Ratovelomana, Y. Rollin, C. Gébénne, C. Gosmini and J. Périchon, *Tetrahedron Lett.*, 1994, 4777.
- 15 K. Yamada, T. Kubo, H. Tokuyama and T. Fukuyama, *Synlett*, 2002, 231.
- 16 See the following for leading references on previous syntheses of 6a,7-dehydroaporphines and references cited therein: (a) M. P. Cava, I. Noguchi and K. T. Buck, *J. Org. Chem.*, 1973, **38**, 2394; (b) G. R. Lenz and F. J. Koszyk, *J. Chem. Soc., Perkin Trans. 1*, 1984, 1273; (c) C. Saá, E. Guitian, L. Castedo and J. M. Saá, *Tetrahedron Lett.*, 1985, **26**, 4559; (d) M. P. Cava, D. L. Edie and J. M. Saá, *J. Org. Chem.*, 1975, **40**, 3601; (e) N. C. Yang, G. R. Lenz and A. Shani, *Tetrahedron Lett.*, 1966, 2941; (f) K. Orito, S. Uchiito, Y. Satoh, T. Tatsuzawa, R. Harada and M. Tokuda, *Org. Lett.*, 2000, **2**, 307.
- 17 Note that *O*-methyldehydroisopiline **10** isolated from the natural source had not been obtained in analytically pure form. Its spectral data, however, allowed for an unambiguous determination of its structure, cf. ref. 5. In contrast, the synthetic material gave fully satisfactory data. It is important to mention, however, that **10** is unstable when exposed to air, as can be judged from a rather rapid coloration of the sample. Likewise, compound **11** is unstable in solution as can be seen from a characteristic color change from yellow to green.
- 18 A. Venkateswarlu and M. P. Cava, *Tetrahedron*, 1976, **32**, 2079.
- 19 (a) S. Ruchirawat and S. Predapitakkun, *Heterocycles*, 2001, **55**, 371; (b) M. Gerecke, R. Borer and A. Bossi, *Helv. Chim. Acta*, 1975, **58**, 185; (c) L. Castedo, R. Riguera, J. M. Saá and R. Suau, *Heterocycles*, 1977, **6**, 677.
- 20 M. Smrčina, Š. Vyskočil, B. Máca, M. Polášek, T. A. Claxton, A. P. Abbott and P. Kočovský, *J. Org. Chem.*, 1994, **59**, 2156.