

A New Chiral Synthesis of (—)-Anisomycin and its Demethoxy Analogue

J. Grant Buchanan,^{*a} Keith A. MacLean,^a Hans Paulsen,^b and Richard H. Wightman^a

^a *Department of Chemistry, Heriot-Watt University, Riccarton, Edinburgh EH14 4AS, U.K.*


^b *Institut für Organische Chemie und Biochemie der Universität Hamburg, D-2000 Hamburg 13, Federal Republic of Germany*

(—)-Anisomycin (**1a**) and its demethoxy-analogue (**1b**) have been prepared enantiospecifically and stereoselectively from D-ribose.

The antibiotic anisomycin (**1a**), with relative¹ and absolute² configuration as shown in Scheme 2, has been isolated from the culture filtrates of various species of *Streptomyces*; it

possesses promising antiprotozoal activity, and has been shown to block ribosomal peptide synthesis.³

Early syntheses of anisomycin were non-stereoselective and




Scheme 1. i, ArCH_2MgCl , THF; ii, NaIO_4 ; iii, NaBH_4 , EtOH; iv, $(\text{CF}_3\text{SO}_2)_2\text{O}$, $\text{C}_6\text{H}_5\text{N}$; v, PhCOCl , $\text{C}_6\text{H}_5\text{N}$; vi, *p*- $\text{MeC}_6\text{H}_4\text{SO}_2\text{Cl}$, $\text{C}_6\text{H}_5\text{N}$; vii, KOH , MeOH; viii, excess of MeSO_2Cl , $\text{C}_6\text{H}_5\text{N}$.

proceeded in poor overall yield.⁴ Since then a much more efficient chiral synthesis of (–)-anisomycin (**1a**) has been reported,⁵ and very recently Schumacher and Hall have announced a highly efficient and stereospecific synthesis of the racemate and two analogues including the racemic demethoxy-compound (**1b**).⁶ We now report a new chiral and stereoselective synthesis of (–)-anisomycin (**1a**) and analogue (**1b**), proceeding from D-ribose in *ca.* 10% overall yield.

Treatment of 2,3-O-isopropylidene-D-ribose (**2**)⁷ with a large excess of the appropriate Grignard reagent in tetrahydrofuran (THF) gave (**3a**) (77%) and (**3b**) (70%) (see Scheme 1). In each case a single stereoisomer was produced, which was assigned the D-*allo*-configuration shown on the basis of earlier work,⁸ and this was confirmed as described below. Periodate oxidation of (**3a,b**) gave the crystalline hemiacetals (**4a,b**) readily reduced with borohydride to diols (**5a,b**). Treatment of (**5b**) with trifluoromethanesulphonic anhydride caused cyclisation to the tetrahydrofuran derivative (**6**), whilst monobenzoylation, and subsequent base treatment gave the all-*cis*-isomer (**7**). Examination of ^{13}C n.m.r. spectra of (**6**) and (**7**) confirmed the stereochemistry of the initial Grignard reaction; on the basis of earlier work,^{9,10} one would predict upfield shifts for C-1, C-2, and C-3, as well as the carbons of the isopropylidene group, in the all-*cis*-arrangement (**7**) as compared with β -isomer (**6**). Such an upfield shift was observed for all six carbon atoms.

Diols (**5a,b**) were readily converted into dimesylates (**8a,b**); we had hoped that reaction of (**8a,b**) with ammonia or a primary amine would give an all-*cis* pyrrolidine, but these dimesylates were surprisingly unreactive with a variety of nitrogen nucleophiles,[†] and an alternative route was sought (Scheme 2).



Scheme 2. i, $\text{NH}_2\text{OH}\cdot\text{HCl}$, $\text{C}_6\text{H}_5\text{N}$; ii, excess of MeSO_2Cl , $\text{C}_6\text{H}_5\text{N}$; iii, LiAlH_4 , ether; iv, HCl , MeOH, H_2O , reflux; v, HBr , HOAc, 10°C ; vi, KOH , MeOH, H_2O ; vii, allyl alcohol, HClO_4 , CHCl_3 , 60°C ; viii, PhCH_2Br , Et_3N , CHCl_3 ; ix, Ac_2O , $\text{C}_6\text{H}_5\text{N}$; x, Pd/C , MeOH, dil. HCl , reflux; xi, Pd/C , H_2 , MeOH, HCl .

Hemiacetals (**4a,b**) were converted into the oximes (**9a,b**) (95%, mixture of *E*- and *Z*-isomers, *ca.* 70:30). Treatment of these with mesyl chloride in pyridine formed the *O*-mesyl nitriles (**10a,b**), which without purification were reduced with lithium aluminium hydride to give pyrrolidines (**11a,b**) isolated as their crystalline hydrochlorides [60% from (**4a,b**)]. Hydrolysis, and subsequent treatment of the diol hydrochlorides with HBr in glacial acetic acid¹¹ gave, presumably *via* the 3,4-acetoxonium ion, mixtures of *trans*-bromoacetates in which (**12a,b**) predominated;‡ treatment of the mixed bromoacetates with base gave epoxides (**13a,b**) [68% from (**11a,b**)], (**13a**) proving identical with material produced from (–)-anisomycin.¹²

Regioselective acid-catalysed ring opening with allyl alcohol[‡] then gave alcohols (**14a,b**) (*ca.* 65%); *N*-benzoylation and subsequent *O*-acetylation yielded (**15a,b**), which were deprotected by palladium-catalysed rearrangement and hydrolysis

† Examination of a molecular model indicates considerable hindrance to either the incoming nucleophile, or the departing leaving group, for displacement at C-5 of (**8**). Forcing conditions produced 1,2-elimination products.

‡ The regioselectivity in this reaction is predictable from results obtained during structural work on anisomycin (refs. 1a, 2, and 12); the regioselectivity is lost if the nitrogen is substituted (refs. 4b and 4c).

of the allyl group, followed by hydrogenolysis, to give (–)-anisomycin (**1a**), identical with natural material, and the demethoxy-analogue (**1b**) [70% from (**14a,b**)]. The synthetic (–)-anisomycin (**1a**) had identical activity towards *Trichomonas vaginalis* compared with the natural product, whilst (**1b**) had about one sixth of this activity.

We thank S.E.R.C. (Studentship to K. A. M.) and Nuffield Foundation (One-Year Science Research Fellowship to R. H. W.) for financial support, and Dr. K. Richardson (Pfizer Central Research) for providing the biological data and a generous gift of natural anisomycin.

Received, 14th February 1983; Com. 211

References

- 1 (a) J. R. Schaefer and P. J. Wheatley, *J. Org. Chem.*, 1968, **33**, 166; (b) K. Butler, *ibid.*, p. 2136.
- 2 C. M. Wong, *Can. J. Chem.*, 1968, **46**, 1101.
- 3 A. Jimenez and D. Vazquez, in 'Antibiotics,' ed. F. E. Hahn, Springer Verlag, Berlin, 1979, p. 1, and references therein.
- 4 (a) C. M. Wong, J. Buccini, and J. TeRaa, *Can. J. Chem.*, 1968, **46**, 3091; (b) C. M. Wong, J. Buccini, I. Chang, J. TeRaa, and R. Schwenk, *ibid.*, 1969, **47**, 2421; (c) S. Oida and E. Ohki, *Chem. Pharm. Bull.*, 1969, **17**, 1405; (d) I. Felner and K. Schenker, *Helv. Chim. Acta*, 1970, **53**, 754.
- 5 J. P. H. Verheyden, A. C. Richardson, R. S. Bhatt, B. D. Grant, W. L. Fitch, and J. G. Moffatt, *Pure Appl. Chem.*, 1967, **50**, 1363.
- 6 D. P. Schumacher and S. S. Hall, *J. Am. Chem. Soc.*, 1982, **104**, 6076.
- 7 N. A. Hughes and P. R. H. Speakman, *Carbohydr. Res.*, 1965, **1**, 171.
- 8 J. G. Buchanan, A. D. Dunn, and A. R. Edgar, *J. Chem. Soc., Perkin Trans. 1*, 1975, 1191.
- 9 H. Ohru, G. H. Jones, J. G. Moffatt, M. L. Maddox, A. T. Christensen, and S. K. Byram, *J. Am. Chem. Soc.*, 1975, **97**, 4602.
- 10 J. G. Buchanan, A. R. Edgar, D. I. Rawson, P. Shahidi, and R. H. Wightman, *Carbohydr. Res.*, 1982, **100**, 75.
- 11 B. T. Golding, D. R. Hall, and S. Sakrikar, *J. Chem. Soc., Perkin Trans. 1*, 1973, 1214.
- 12 J. J. Beereboom, K. Butler, F. C. Pennington, and I. A. Solomons, *J. Org. Chem.*, 1965, **30**, 2334.