

Phosphorus oxychloride as an efficient coupling reagent for the synthesis of esters, amides and peptides under mild conditions†

Cite this: *RSC Advances*, 2013, 3, 16247

Received 11th June 2013,

Accepted 12th July 2013

DOI: 10.1039/c3ra42887g

www.rsc.org/advances

Hu Chen,^{*ab} Xunfu Xu,^a Liu Liu,^a Guo Tang^{*a} and Yufen Zhao^a

A mild method is described for the conversion of carboxylic acids into esters, amides, as well as peptides without racemization through carboxyl activation by the reagent combination of POCl₃ and DMAP. Long chain alcohols could be converted to the corresponding ester in good yields. ³¹P NMR spectrum was used to detect phosphorus-containing intermediates in ongoing reactions directly, and a possible mechanism has been proposed based on these results.

Esters and amides, particularly peptides, have been widely used as chemicals and pharmaceuticals.¹ Activation of the carboxylic groups is required before the reaction can occur. A large number of methods are available for activation of carboxylic acids, including pre-activation of the carboxyl group with dicyclohexylcarbodiimide (DCC),² Yamaguchi acid chloride,³ thionyl chloride,⁴ dimethylsulfamoyl chloride,⁵ mixed anhydride,⁶ triphenylphosphine,⁷ 1-tosylimidazole,⁸ O-alkylisoureas,⁹ imidazole carbamates and ureas,¹⁰ iodosodilactone,¹¹ T3P,¹² other organocatalysts¹³ and coupling agents.¹⁴ Unfortunately, two major disadvantages of most activating reagents invented to date are that they generate substantial amounts of undesired by-products and the potential loss of chiral integrity at the carboxyl residue undergoing activation. Moreover, the preparation of these coupling reagents is often very difficult and requires harsh conditions. Therefore, improved methods for the synthesis of ester and amide bonds, particularly peptide couplings are in great demand.

As one of the most common and diversely used industrial chemicals, phosphorus oxychloride (POCl₃) is widely used as a dehydrating agent in the laboratory, a liquid phosphorus source in diffusion processes in the semiconductor industry, and an important material in the manufacture of herbicides, insecticides,

plasticizers, oil additives, and flame retardants. POCl₃ has been used for amide synthesis in recent years.¹⁵ However, these methods still suffered from some drawbacks such as high temperature, long reaction time, and often moderate yields. Herein, we wish to report that the combination of POCl₃ and DMAP efficiently mediates the esterification and amidation of carboxylic acid under mild conditions. The method provides an efficient method for the activation of carboxylic acid.


We began our study by examining the reaction of 3-phenylpropanoic acid (**1a**) with phenylmethanol (**2a**). To our satisfaction, when a mixture of **1a** (0.50 mmol), **2a** (0.60 mmol), Et₃N (1.2 mmol) and a catalytic amount of DMAP (0.15 mmol) in CH₂Cl₂ (3.0 mL) at room temperature with POCl₃ (0.50 mmol) for 2 h, benzyl 3-phenylpropanoate (**3a**) was obtained in 96% yield. Many solvents (THF, toluene, dichloroethane, and acetonitrile) are effective, obtaining **3a** in all cases more than 90% yields. Inorganic bases such as K₃PO₄, K₂CO₃ and Na₂CO₃ were less effective, giving **3a** in 60–70% yields. Organic bases such as *N*-methylmorpholine and pyridine also gave high yields. This is likely due to the greater solubility of these bases in organic solvents, leading to the rapid formation of the mixed carboxylic-phosphoric anhydride. However, no product **3a** was observed in the absence of base. Without DMAP, the reaction was very sluggish and gave **3a** in 32% yield after 12 h.

With the optimized conditions in hand, a variety of alcohols, phenols and carboxylic acids were examined (Table 1). The desired esters were obtained in good to excellent yields regardless of primary alcohols (**3a–3d**) or secondary alcohol *L*-menthol also performed well and yielded the corresponding optically pure ester product **3f** with retention of configuration around the alkoxy carbon. Furthermore, phenol and 1-naphthol also proceeded smoothly, affording the corresponding products in good yields (**3g**, **3h**). On the other hand, both aliphatic and aromatic carboxylic acids reacted with phenylmethanol to give products **3k–3n** in good to excellent yields. Moreover, the reactions are completely compatible with the presence of double and triple bonds, both terminal and internal, and no products of addition over these multiple bonds were observed (**3b**, **3c**, and **3q**). Additionally, α,β -unsaturated

^aDepartment of Chemistry, College of Chemistry and Chemical Engineering, and the Key Laboratory for Chemical Biology of Fujian Province, Xiamen University, Xiamen, Fujian 361005, China. E-mail: t12g21@xmu.edu.cn; Fax: +86 592 2185780; Tel: +86 592 2185780

^bDepartment of Chemistry and Chemical Engineering, Hefei Normal University, Hefei 230601, China. E-mail: hchen808@yahoo.cn


† Electronic supplementary information (ESI) available. See DOI: 10.1039/c3ra42887g

Table 1 Esterification promoted by POCl₃ and DMAP^a

^a Reaction conditions: acid (0.5 mmol), alcohols or phenol (0.6 mmol), Et₃N (1.2 mmol), POCl₃ (0.5 mmol), DMAP (0.15 mmol), solvent (3 mL), rt, 2 h. Isolated yields. ^b 8 h.

carboxylic acid and heterocyclic substituted carboxylic acid were also successfully employed in the reaction with phenylmethanol under the standard conditions (3i, 3k). *tert*-Butyl alcohol was also examined. Unfortunately, only trace amounts of desired ester bond-forming products were detected.

It is worth noting that long chain aliphatic acids and long chain alcohols could be converted to the corresponding ester (3o, 3p), although it is less reactive than others.¹⁶ Finally, relatively hindered alcohol was efficiently esterification with our method in a 76% yield (3q). Although several methods have been described for the esterification of these steroidal structures, normally using strong basic or acidic media and high temperatures, our procedure proceeds under milder conditions and contribute a


Table 2 Amidation and peptide coupling promoted by POCl₃ and DMAP^a

^a Reaction conditions: acid or *N*-protected amino acid (0.5 mmol), amine or methyl ester amino acid (0.6 mmol), Et₃N (1.2 mmol), POCl₃ (0.5 mmol), DMAP (0.15 mmol), solvent (3 mL), rt, 2 h. Isolated yields.

precise control over the individual reactivity of functional groups within a complex molecular architecture, which constitutes an important objective in the synthesis of complex natural products.

After the successful construction of an ester bond, we next turned to change the nucleophile to amines, which are generally thought to be easier to acylate than alcohol. To our satisfaction, various aliphatic or aromatic primary amines and secondary amines effectively reacted with 3-phenylpropanoic acid (1a) to provide the desired amides 5a–f in good to excellent yields (Table 2). Terminal alkene group was tolerated under the present condition (5c). Besides, hindered amines also worked well to give corresponding products 5b and 5c. Then, aromatic acids with different functional groups were studied and afforded desired amide products 5j–k in high to excellent yields with butylamine.

Success in the construction of amides inspired us to explore the applications of the present catalysis system in peptide



Scheme 1 Possible reaction mechanism.

coupling reactions. Accordingly, *N*-Cbz-phenylalanine was treated with methyl ester of leucine in the presence of Et_3N as a base and a catalytic amount of DMAP. POCl_3 was added to the above solution, and the reaction mixture was stirred at room temperature for 2 h. To our immense satisfaction, the product formed was found to be the expected dipeptide **5l** in a 95% yield, and its specific rotation was in good agreement with the reported value [**5l**; $[\alpha]_{\text{D}}^{20} -24.7$ (*c* 3.1, MeOH) (lit.¹⁷ $[\alpha]_{\text{D}}^{20} -24.7$ (*c* 3.1, MeOH))]. The diastereomeric purity was further verified by HPLC analysis of **5l** and a diastereomeric mixture of (D,L)- and (L,L)-**5l** [prepared by coupling Leu-OMe with racemic Cbz-phe], which clearly showed **5l** to be free from any (D,L) isomer, thereby indicating the nonracemizing nature of the above coupling protocol. A variety of amino acid combinations were similarly subjected to dipeptide formation affording the products **5l-r** in good yields and high diastereomeric purity.

A possible mechanism is proposed as shown in Scheme 1. POCl_3 is initially activated by DMAP to form the phosphorylpyridinium intermediate **6**¹⁸ that then undergo the addition of the Cbz-Phe to give the mixed anhydride **7**. Once the mixed anhydride is formed, it reacts rapidly with DMAP to form acylpyridinium intermediate **8**, which activates the amino acid carbonyl toward nucleophilic attack. At last, the activated species **8** is attacked by Leu-OMe to give the product **5l**, DMAP and PO_2Cl_2^- .

To further study the reaction mechanism, we decided to use ^{31}P NMR spectrum analysis, which has been successfully employed in studying a number of phosphorus-containing reactions.¹⁹ The formation of **5l** was monitored by ^{31}P NMR spectroscopy, as shown in Fig. 1. The initial ^{31}P NMR spectrum studies indicated Et_3N was used as a base only, which prompted the use of excess DMAP instead of Et_3N , reducing the sets of ^{31}P NMR spectrum peaks. The starting POCl_3 in CH_2Cl_2 showed signal in the ^{31}P NMR spectrum at $\delta = 4.7$ ppm. After DMAP (2.4 eq.) was added to the solution of POCl_3 , the peak of POCl_3 was decreasing while a new peak at -15.8 ppm corresponding to intermediate **6** emerged (A in Fig. 1). When Cbz-Phe (**1k**) was added to the mixture, two sets of new signals appear at -5.8 and -10.4 ppm (B in Fig. 1), which indicated the forming of the mixed phosphoric-carboxylic anhydride **7**^{19b,20} that subsequently reacted with DMAP to give the more reactive intermediate **8**. Although no spectro-




Fig. 1 ^{31}P NMR spectra for the synthesis of **5l**. A: POCl_3 (1.0 mmol), DMAP (2.4 mmol), CH_2Cl_2 (1 mL). B: Cbz-Phe (1.5 mmol) in CH_2Cl_2 (2 mL) was added to A. C: Leu-OMe (1.5 mmol) in CH_2Cl_2 (2 mL) was added to B.

scopic data for intermediate **8** are available, PO_2Cl_2^- (-5.8 ppm) has been previously studied by ^{31}P NMR,²¹ showing chemical shifts at -7.5 ppm in MeCN, similar to that observed for our proposed intermediate. However, addition of Leu-OMe resulted in a peak (-10.4 ppm) that rapidly decreased (C in Fig. 1), meanwhile the peak at -5.8 ppm gradually increased, indicating the existence of the intermediate **7** again. In addition, a new peak at -19.5 ppm was assigned as the by-product **9**, which was proved by a mix of acetylpyridinium intermediate **6** with Leu-OMe.

In summary, we have developed a mild, safe, and economic one-pot protocol for the conversion of carboxylic acids into esters, amides, as well as peptides without racemization through carboxyl activation by the reagent combination of POCl_3 and DMAP. The method eliminates some commonly encountered problems, such as separation of product from reagent derived coproducts, racemization, poor recovery of product, high cost, instability of the reagent, etc. The “*in situ*” activation of the carboxylic acid makes our method an excellent alternative to the known acylation procedures, useful for the synthesis of peptides, particularly complex natural products. The suggested possible mechanism is supported by the ^{31}P NMR spectrum analyses.

Acknowledgements

We acknowledge financial support from the Chinese National Natural Science Foundation (21173178, 21232005), Program for Changjiang Scholars and Innovative Research Team in University, Collaborative Innovation Center of Chemistry for Energy Materials, and Anhui (KJ2012B145, 2012rcjj03).

Notes and references

- (a) J. Otera and J. Nishikido, *Esterification: Methods, Reactions, and Applications*, Wiley-VCH, Weinheim, 2nd ed., 2010; (b) V. R. Pattabiraman and J. W. Bode, *Nature*, 2011, **480**, 471–479.
- (a) B. Neises and W. Steglich, *Angew. Chem., Int. Ed. Engl.*, 1978, **17**, 522–524; (b) A. Hassner and V. Alexanian, *Tetrahedron Lett.*, 1978, **19**, 4475–4478.
- (a) J. Inanaga, K. Hirata, H. Saeki, T. Katsuki and M. Yamaguchi, *Bull. Chem. Soc. Jpn.*, 1979, **52**, 1989–1993; (b) I. Dhimitruka and J. SantaLucia, *Org. Lett.*, 2006, **8**, 47–50.
- F. Matsuda, S. Itoh, N. Hattori, M. Yanagiya and T. Matsumoto, *Tetrahedron*, 1985, **41**, 3625–3631.

- 5 K. Wakasugi, A. Nakamura, A. Iida, Y. Nishii, N. Nakatani, S. Fukushima and Y. Tanabe, *Tetrahedron*, 2003, **59**, 5337–5345.
- 6 (a) J. R. Vaughan and R. L. Osato, *J. Am. Chem. Soc.*, 1951, **73**, 5553–5555; (b) E. Valeur and M. Bradley, *Tetrahedron*, 2007, **63**, 8855–8871.
- 7 (a) G. Tang, G. J. Zhou, F. Ni, L. M. Hu and Y. F. Zhao, *Chinese Chem. Lett.*, 2005, **16**, 385–388; (b) S. P. Morcillo, L. Á. Cienfuegos, A. J. Mota, J. Justicia and R. Robles, *J. Org. Chem.*, 2011, **76**, 2277–2281; (c) T. Taniguchi, D. Hirose and H. Ishibashi, *ACS Catal.*, 2011, **1**, 1469–1474.
- 8 M. N. S. Rad, S. Behrouz, M. A. Faghihi and A. Khalafi-Nezhad, *Tetrahedron Lett.*, 2008, **49**, 1115–1120.
- 9 A. Chighine, S. Crosignani, M. C. Arnal, M. Bradley and B. Linclau, *J. Org. Chem.*, 2009, **74**, 4753–4762.
- 10 S. T. Heller and R. Sarpong, *Org. Lett.*, 2010, **12**, 4572–4575.
- 11 J. Tian, W. C. Gao, D. M. Zhou and C. Zhang, *Org. Lett.*, 2012, **14**, 3020–3023.
- 12 J. R. Dunetz, Y. Xiang, A. Baldwin and J. Ringling, *Org. Lett.*, 2011, **13**, 5048–5051.
- 13 H. W. Horn, G. O. Jones, D. S. Wei, K. Fukushima, J. M. Lecuyer, D. J. Coady, J. L. Hedrick and J. E. Rice, *J. Phys. Chem. A*, 2012, **116**, 12389–12398.
- 14 (a) J. K. Twibanire and T. B. Grindley, *Org. Lett.*, 2011, **13**, 2988–2991; (b) A. El-Faham and F. Albericio, *Chem. Rev.*, 2011, **111**, 6557–6602; (c) E. Valeur and M. Bradley, *Chem. Soc. Rev.*, 2009, **38**, 606–631; (d) C. A. G. N. Montalbetti and V. Falque, *Tetrahedron*, 2005, **61**, 10827–10852; (e) K. Thalluri, K. C. Nadimpally, M. P. Chakravarty, A. Paul and B. Mandal, *Adv. Synth. Catal.*, 2013, **355**, 448–462.
- 15 (a) K. Sakamoto, K. Sato, A. Shigenaga, K. Tsuji, S. Tsuda, H. Hibino, Y. Nishiuchi and A. Otaka, *J. Org. Chem.*, 2012, **77**, 6948–6958; (b) A. Bach, N. Stühr-Hansen, T. S. Thorsen, N. Bork, I. S. Moreira, K. Frydenvang, S. Padrah, S. B. Christensen, K. L. Madsen, H. Weinstein, U. Gether and K. Stromgaard, *Org. Biomol. Chem.*, 2010, **8**, 4281–4288; (c) Y. D. Gong, M. K. Jeon, H. La and D. C. Ha, *Synlett*, 2007, 1431–1435.
- 16 (a) K. Mantri, K. Komura and Y. Sugi, *Synthesis*, 2005, 1939–1944; (b) N. Ieda, K. Mantri, Y. Miyata, A. Ozaki, K. Komura and Y. Sugi, *Ind. Eng. Chem. Res.*, 2008, **47**, 8631–8638.
- 17 M. Ueda and H. Oikawa, *J. Org. Chem.*, 1985, **50**, 760–763.
- 18 F. Sartillo-Piscio, L. Quintero, C. Villegas, E. Santacruz-Juarez and C. A. Parrodi, *Tetrahedron Lett.*, 2002, **43**, 15–17.
- 19 (a) S. Kong, W. Fan, G. Wu and Z. Miao, *Angew. Chem., Int. Ed.*, 2012, **51**, 8864–8867; (b) Q. L. Luo, L. Lv, Y. Li, J.-P. Tan, W. Nan and Q. Hui, *Eur. J. Org. Chem.*, 2011, 6916–6922; (c) H. A. Bronstein and C. K. Luscombe, *J. Am. Chem. Soc.*, 2009, **131**, 12894–12895.
- 20 R. Kluger and L. L. Cameron, *J. Am. Chem. Soc.*, 2002, **124**, 3303–3308.
- 21 (a) D. Amsallem, H. Gornitzka, A. Baceiredo and G. Bertrand, *Angew. Chem., Int. Ed.*, 1999, **38**, 2201–2203; (b) Z. Voitenko, M. R. Mazieres, M. Sanchez and J. G. Wolf, *Tetrahedron*, 2001, **57**, 1059–1066.