

# Phorboxazole Synthetic Studies. 2. Construction of a C(20–28) Subtarget, a Further Extension of the Petasis–Ferrier Rearrangement


Amos B. Smith, III,\* Kevin P. Minbiole, Patrick R. Verhoest, and  
Thomas J. Beauchamp

Department of Chemistry, Laboratory for Research on the Structure of Matter, and  
Monell Chemical Senses Center, University of Pennsylvania,  
Philadelphia, Pennsylvania 19104

smithab@sas.upenn.edu

Received July 19, 1999

## ABSTRACT



In this, the second of two Letters, we describe the efficient assembly of (+)-**4**, a C(20–28) subtarget for the total synthesis of phorboxazoles A (**1**) and B (**2**). The synthesis was achieved in 12 linear steps (20% overall yield) via Petasis–Ferrier rearrangement of an *E:Z* mixture of trisubstituted enol ethers (**15**) to assemble the C(22–26) *cis*-tetrahydropyran. A mechanism for the observed diastereoconvergence of **15** is proposed. In addition, a new tactic for the synthesis of enol ethers (e.g., **15**) based on the elegant work of Julia is described.

Phorboxazoles A (**1**) and B (**2**), rare marine macrolides comprised of three tetrahydropyrans, two oxazoles, and a 21-membered macrolactone, display extraordinary cancer cell growth inhibition, and as such have attracted considerable interest in the synthetic community.<sup>1–3</sup> In the preceding Letter,<sup>4</sup> we outlined a strategy for the construction of **1** and **2**, in conjunction with an efficient synthesis of the C(3–19) subtarget (–)-**5**, exploiting an extension of the Petasis–Ferrier rearrangement<sup>5,6</sup> to assemble the C(11–15) tetrahy-

dropyran ring (Scheme 1). The success of the Petasis–Ferrier rearrangement encouraged us to explore a similar tactic for construction of the fully substituted C(22–26) tetrahydropyran moiety present in the C(20–28) subtarget **4**.

Our modification of the Petasis–Ferrier rearrangement permits direct conversion of a 4-alkylidenyl-1,3-dioxane (**9**) to the corresponding *cis*-tetrahydropyranone (**10**) when Me<sub>2</sub>AlCl is employed as the Lewis acid. This protocol avoids the Meerwein–Ponndorf–Verley reduction of the initially derived tetrahydropyranone by *i*-Bu<sub>3</sub>Al, as observed by Petasis. To extend the Petasis–Ferrier rearrangement to assemble a fully substituted tetrahydropyran such as **4** (Scheme 1), trisubstituted enol ether **7** would be required. No information on the stereochemical outcome for trisubstituted enol ethers however was available. To explore this stereochemical issue, we prepared and rearranged (–)-**11**,<sup>7</sup> a model of enol ether **7** (Scheme 2).

(1) Searle, P. A.; Molinski, T. F. *J. Am. Chem. Soc.* **1995**, *117*, 8126.

(2) (a) Searle, P. A.; Molinski, T. F.; Brzezinski, L. J.; Leahy, J. W. *J. Am. Chem. Soc.* **1996**, *118*, 9422. (b) Molinski, T. F. *Tetrahedron Lett.* **1996**, *37*, 7879.



(3) Forsyth, C. J.; Ahmed, F.; Cink, R. D.; Lee, C. S. *J. Am. Chem. Soc.* **1998**, *120*, 5597. For references to other synthetic efforts, see ref 4.

(4) Smith, A. B., III.; Verhoest, P. R.; Minbiole, K. P.; Lim, J. J. *Org. Lett.* **1999**, *1*, 909.


(5) Petasis, N. A.; Lu, S. P. *Tetrahedron Lett.* **1996**, *36*, 141.

(6) Ferrier, R. J.; Middleton, S. *Chem. Rev.* **1993**, *93*, 2779.

Scheme 1




Scheme 2



Although Petasis suggests the rearrangement proceeds via a chair conformation, we rationalized that a least motion mechanism involving a boat conformation (**i**  $\rightarrow$  **ii**, Scheme 3) held promise of delivering the required stereochemical outcome (e.g., **14**). In the event, however, Petasis–Ferrier rearrangement of (–)-**11** furnished only the all equatorial tetrahydropyran (+)-**12** (58% unoptimized); the configuration of (+)-**12** was secured via 1D-NOE experiments.<sup>8</sup> Presumably, Lewis acid complexation at the enol ether oxygen of (–)-**11** triggers ring opening, reversibly liberating the


Scheme 3



aluminum enolate; rotation of the enolate by 180° (**i**  $\rightarrow$  **iii**) and reclosure via a chair conformation would afford (+)-**12**.

Armed with this insight, we envisioned the oxygen-transposed enol ether **15** to be an appropriate substrate for the construction of **4** (Scheme 4). Rearrangement involving


Scheme 4



bond rotation of 180° would lead via a chair conformation to **16** possessing the requisite axial methyl at C(23).


Assembly of **15** began with aldol condensation of the boron enolate derived from Evans oxazolidinone (+)-**17**<sup>9</sup> with aldehyde **8**;<sup>7c</sup> hydrolysis (LiOH, H<sub>2</sub>O<sub>2</sub>) afforded  $\beta$ -hydroxy acid (+)-**18**<sup>10</sup> in 84% yield for the two steps (Scheme 5). Bis-silylation with hexamethyldisilazane (HMDS),<sup>11</sup> followed by TMSOTf<sup>12</sup>-promoted condensation with alde-

Scheme 5




hyde **19**<sup>13</sup> furnished dioxanone (+)-**20**<sup>10</sup> in 66% yield, along with 19% of the C(26) epimer, the latter readily removed by flash chromatography.<sup>14</sup> The configuration of (+)-**20** was again determined by 1D-NOE experiments. Unfortunately, ethylidenation via the Takai protocol<sup>7d</sup> failed to yield **15**, furnishing instead the C(23) epimeric enol ether as a *E/Z* mixture (**21**). Related olefination strategies were equally unsuccessful.<sup>15</sup>

Undaunted, we explored the Julia protocol for olefination of sulfones with electrophilic carbenoids.<sup>16</sup> This protocol calls for  $\alpha$ -alkylation of a sulfone (**22a**) with  $\alpha$ -halo Grignard reagents (**23**); subsequent elimination furnishes the alkene (**24a**; Scheme 6). We reasoned that a similar reaction with



sulfone **22b** ( $R_2 = \text{OR}$ ) would afford **24b**, contingent on preferential expulsion of phenyl sulfinate over the alkoxide.<sup>17</sup>

(7) (a) Enol ether (–)-**11** was prepared by condensation of  $\beta$ -hydroxy acid (+)-**6**<sup>7b</sup> and aldehyde **8**,<sup>7c</sup> hydrogenation of the resultant dioxanone, and Takai ethylidenation.<sup>7d</sup> (b) (+)-**6** was prepared according to Oppolzer's protocol: Oppolzer, W.; Lienard, P. *Tetrahedron Lett.* **1993**, 34, 4321. (c) Boeckman, R. K., Jr.; Charette, A. B.; Asberom, T.; Johnston, B. H. *J. Am. Chem. Soc.* **1987**, 109, 7553. (d) Okazoe, T.; Takai, K.; Oshima, K.; Utimoto, K. *J. Org. Chem.* **1987**, 52, 4410.



(8) That **12** is not the result of epimerization of the axial methyl group was demonstrated via exposure of **14** to the reaction conditions; no epimerization occurred.

(9) Evans, D. A.; Bartroli, J.; Shih, T. L. *J. Am. Chem. Soc.* **1981**, 103, 2127.

(10) The structure assigned to each new compound is in accord with its infrared, 500 MHz <sup>1</sup>H NMR, and 125 MHz <sup>13</sup>C NMR spectra, as well as appropriate ion identification by high-resolution mass spectrometry.

(11) (a) Harada, T.; Yoshida, T.; Kagamihara, Y.; Oku, A. *J. Chem. Soc. Chem. Commun.* **1993**, 1367. (b) Seebach, D.; Imwinkelried, R.; Stucky, G. *Helv. Chim. Acta* **1987**, 70, 448.

(12) Initial difficulties in the scale-up of this reaction suggested that TfOH is the actual catalyst. Advantageous water, more pronounced on smaller scale, may generate TfOH in situ from TMSOTf (as well as TMS<sub>2</sub>O). Large-scale reactions do not proceed until catalytic TfOH (2–4 mol %) is added. Yields and diastereoselectivity were similar with the added TfOH.

(13) (a) Lange, T.; van Loon, J.-D.; Tykwinski, R. R.; Schreiber, M.; Diederich, F. *Synthesis*, **1996**, 537. (b) For an improved general route to  $\alpha,\beta$ -acetylenic aldehydes, see: Journet, M.; Cai, D.; DiMichele, L. M.; Larsen, R. D. *Tetrahedron Lett.* **1998**, 39, 6427.


(14) Although all attempts to epimerize the undesired C(26) isomer to (+)-**20** were unsuccessful (e.g., with TMSOTf), hydrolysis of the epimer (LiOH, H<sub>2</sub>O/THF) afforded (+)-**18** in 97% yield.

(15) Horikawa, Y.; Watanabe, M.; Fujiwara, T.; Takeda, T. *J. Am. Chem. Soc.* **1997**, 119, 1127.

(16) De Lima, C.; Julia, M.; Verpeaux, J.-N. *Synlett* **1992**, 133.

(17)  $\alpha$ -Amido sulfones in a similar reaction, see: Alonso, D. A.; Alonso, E.; Najera, C.; Ramon, D. J.; Yus, M. *Tetrahedron Lett.* **1997**, 53, 4835.


**Scheme 7**



Toward this end, reduction of dioxanone (+)-**20** (DIBAL) followed by in situ acylation of the alkoxide (Ac<sub>2</sub>O, DMAP) afforded acetal (+)-**25**<sup>10</sup> (Scheme 7).<sup>18</sup> Treatment of (+)-**25** with PhSTMS in the presence of ZnI<sub>2</sub><sup>19</sup> then led to the corresponding sulfide which upon oxidation (*m*CPBA) generated sulfone (+)-**26**<sup>10</sup> in 68% yield for the two steps. Deprotonation of (+)-**26** with *n*-BuLi and exposure to Grignard **27**<sup>16,20</sup> furnished the desired enol ether **15** in excellent yield (95%), albeit with no *E/Z* selectivity. Careful flash chromatography permitted separation of the *E* and *Z* diastereomers; again the stereochemistry was secured by NOE experiments.

To our delight, treatment of the 1:1 mixture of enol ethers (**15**) with Me<sub>2</sub>AlCl afforded *only the desired tetrahydropyran* (+)-**16**<sup>10,21</sup> in a 91% yield (Scheme 8). The individual diastereomers also rearranged to tetrahydropyran (+)-**16** in similarly high yields.

**Scheme 8**



Although (+)-**15Z** presumably rearranges through a chair transition state as anticipated in Scheme 3, formation of (+)-**16** from (+)-**15E** implies that the unfavorable 1,3-diaxial interactions in transition state **ii** (Scheme 9) preclude a chair


(18) Dahanukar, V. H.; Rychnovsky, S. D. *J. Org. Chem.* **1996**, 61, 8317.

(19) Evans, D. A.; Trotter, B. W.; Côté, B.; Coleman, P. J. *Angew. Chem., Int. Ed. Engl.* **1997**, 36, 24, 2741.

(20) Simpson, M. M. *Bull. Soc. Chem. Fr.* **1879**, 31, 411.

(21) The stereochemistry of (+)-**16** was secured by NOE experiments and coupling constant analysis.

Scheme 9



conformation and instead lead to a boatlike transition state (**iii**), wherein the *re* face of the enolate approaches the oxocarbenium species. Presumably, the small steric requirements of the alkyne as well as the propargylic stabilization of the oxocarbenium ion<sup>22</sup> lower the transition state energy of **ii** and **iii** comparably. However, the 1,3 diaxial interactions in **ii** evidently outweigh the energetic penalty of adopting the boat conformation in **iii**.


An alternative mechanism would entail *E* to *Z* isomerization of the enol ether followed by rearrangement through a chair conformation.<sup>23</sup> All attempts, however, to observe **15Z** upon interruption of the Petasis–Ferrier rearrangement failed to provide evidence for this isomerization.

With an efficient synthesis of tetrahydropyranone (+)-**16** in hand, four steps remained to complete subtarget **4** (Scheme 10): reduction of (+)-**16** with NaBH<sub>4</sub> (91% yield, 15:1 dr) followed by protection of the alcohol (Bz<sub>2</sub>O, >99% yield) provided benzoate (+)-**29**.<sup>10</sup> Desilylation (TBAF, 96% yield) and Dess–Martin<sup>24</sup> oxidation then furnished aldehyde (+)-**4**<sup>10</sup> in 93% yield.

In summary, the Petasis–Ferrier rearrangement has been extended to permit assembly of the fully substituted C(22–

(22) For an example of an acetylene stabilizing an oxocarbenium ion in an acetal opening, see: Denmark, S. E.; Almstead, N. G. *J. Am. Chem. Soc.* **1991**, *113*, 8089.

Scheme 10



26) tetrahydropyran in subtarget (+)-**4**. The synthesis proceeded in 12 linear steps and 20% overall yield. In addition, we have developed a new tactic for enol ether synthesis, extending the elegant work of Julia. Finally, a mechanistic rationale for the diastereoconvergent Petasis–Ferrier rearrangement of *E* and *Z* trisubstituted enol ethers is presented. Studies to assemble the phorboxazole macrocycle and ultimately phorboxazoles A (**1**) and B (**2**) continue in our laboratory.

**Acknowledgment.** Support was provided by the National Institutes of Health (National Cancer Institute) through grant CA-19033.

**Supporting Information Available:** Spectroscopic and analytical data for **4**, **6**, **11–13**, **15**, **16**, **18**, **20**, **25**, **26**, and **29** and selected experimental procedures. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL990829M

(23) For an example of silyl enol ether isomerization, see: Duffy, J. L.; Yoon, T. P.; Evans, D. A. *Tetrahedron Lett.* **1995**, *36*, 9245.

(24) Dess, D. B.; Martin, J. C. *J. Am. Chem. Soc.* **1991**, *113*, 7277.