

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/224878804>

# ChemInform Abstract: Concise Enantioselective Construction of a Bridged Azatricyclic Framework via Domino Semipinacol–Schmidt Reaction.

ARTICLE *in* CHEMICAL COMMUNICATIONS · MAY 2012

Impact Factor: 6.83 · DOI: 10.1039/c2cc31906c · Source: PubMed

---

CITATIONS

7

---

READS

44

## 3 AUTHORS, INCLUDING:



Manohar Puppala

Indian Institute of Technology Madras

3 PUBLICATIONS 21 CITATIONS

SEE PROFILE



Sundarababu Baskaran

Indian Institute of Technology Madras

80 PUBLICATIONS 1,208 CITATIONS

SEE PROFILE

Cite this: *Chem. Commun.*, 2012, **48**, 5778–5780

www.rsc.org/chemcomm

## COMMUNICATION

Concise enantioselective construction of a bridged azatricyclic framework *via* domino semipinacol–Schmidt reaction†

Manohar Puppala, Annamalai Murali and Sundarababu Baskaran\*

Received 15th March 2012, Accepted 17th April 2012

DOI: 10.1039/c2cc31906c

A  $\text{TiCl}_4$ -promoted domino semipinacol–Schmidt reaction of oxaspiropentane-azide provides an easy access to bridged azatricyclic ring systems, which possess the azaquaternary center, present in the immunosuppressant FR901483 and platelet aggregation inhibitor daphlongeranine B.

A number of alkaloids possessing linearly and angularly fused as well as the bridged aza-polycyclic frameworks are widespread in nature (Fig. 1).<sup>1–3</sup> The stemoamide family of alkaloids possess a linearly fused aza-polycyclic framework, whereas angularly fused aza-polycyclic ring systems are widespread in the family of stenine and stemonamine alkaloids.<sup>1</sup> Likewise, structurally diverse bridged aza-polycyclic ring systems are very common in *Daphniphyllum*<sup>2</sup> and *Lycopodium*<sup>3</sup> alkaloids. In recent years, biologically significant alkaloids bearing an intriguing bridged azatricyclic core coupled with the presence of an azaquaternary center have evoked considerable interest among the synthetic chemists.<sup>4,5</sup>

The immunosuppressant FR901483 (**1**) possesses an unusually novel bridged azatricyclic skeleton formed by the spiro fusion of the morphan structural motif (2-azabicyclo[3.3.1]nonane) and pyrrolidine ring (Fig. 2). As a consequence of its potent biological activity and remarkable structural framework, a number of approaches towards the synthesis of FR901483 (**1**) as well as its bridged azatricyclic core, 5-azatricyclo[6.3.1.0<sup>1,5</sup>]dodecane, have been reported.<sup>6,7</sup> Similarly, daphlongeranine B (**2**), a member of the *Daphniphyllum* alkaloids, is a novel platelet aggregation inhibitor having an unprecedented bridged hexacyclic framework coupled with an azaquaternary center whose synthesis has not been realized so far.<sup>8</sup>




Fig. 1 Prevalent aza-structural motifs present in natural products.

Department of Chemistry, Indian Institute of Technology Madras, Chennai, 600036, India. E-mail: sbhaskar@iitm.ac.in; Fax: +91-44-22570545; Tel: +91-44-22574218

† Electronic supplementary information (ESI) available: Representative experimental procedure and characterization of reaction products. CCDC 840274. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c2cc31906c






Fig. 2 Biologically active alkaloids having a bridged azatricyclic core coupled with an azaquaternary center.

Several synthetic approaches for the construction of linearly<sup>9</sup> as well as angularly fused azatricyclic ring systems containing an azaquaternary center have been reported, which include a semipinacol–Schmidt based rearrangement,<sup>10a</sup> a tandem Prins–Schmidt cyclization,<sup>10b</sup> and a nitron based intramolecular dipolar cycloaddition,<sup>10c</sup> however the stereoselective construction of a bridged azatricyclic system having an azaquaternary center is synthetically quite demanding and only a few methods have been devised to achieve this structural motif which includes sequential formal [4 + 3] cycloaddition followed by stereocontrolled enolate chemistry,<sup>5a</sup> an enoxysilane *N*-sulfonyliminium ion cyclization,<sup>5b</sup> and highly diastereoselective formal [3 + 3] cycloaddition followed by transannular Mannich reaction.<sup>5c</sup>

Herein, we report an exceptionally simple and efficient approach for the construction of a bridged azatricyclic ABC-core of FR901483 (**1**) and daphlongeranine B (**2**) using domino semipinacol–Schmidt cyclization as a key step.<sup>11–13</sup> The retrosynthetic analysis is shown in Scheme 1. Oxaspiropentane-azide **5**, a key intermediate in the domino semipinacol–Schmidt cyclization reaction, can be readily synthesized using the Trost spiroannulation<sup>14</sup> protocol from the corresponding azido-ketone **6**,



Scheme 1 Retrosynthetic analysis of a bridged azatricyclic ABC-core of alkaloids **1** and **2**.




**Scheme 2** Synthesis of oxaspiropentane-azide **5**. Reagents: (a)  $\text{CH}_2(\text{CO}_2\text{Me})_2$ ,  $\text{K-O}^t\text{Bu}$ , THF, RT, 92–95%; (b)  $(\text{CH}_2\text{OH})_2$ , PTSA, toluene, reflux; (c)  $\text{NaCl}$ –DMSO, 145 °C, 62–65% over two steps; (d)  $\text{LiAlH}_4$ , THF, RT; (e)  $\text{MsCl}$ ,  $\text{Et}_3\text{N}$ ,  $\text{CH}_2\text{Cl}_2$ , 0 °C; (f)  $\text{NaN}_3$ , DMF, 65 °C; (g) PPTS, acetone– $\text{H}_2\text{O}$  reflux, 48–50% over four steps; (h) ylide,  $\text{KOH}$ , DMSO, 25 °C, 90–94%.

which in turn can be prepared in a few steps starting from cycloalkenone **8** (Scheme 2).

Azido-ketone **6a** was stereoselectively converted to the corresponding *syn*-oxaspiropentane-azide **5a** in excellent yield as a single diastereomer using the Trost spiroannulation reaction. Under similar reaction conditions, azido-ketone **6b** gave a 1 : 1 inseparable mixture of *syn*- and *anti*-oxaspiropentane-azide **5b** in 90% yield, whereas azido-ketone **6c** furnished a 5 : 1 mixture of *syn*- and *anti*-oxaspiropentane-azide **5c**, respectively, in 92% yield (Scheme 2).

When exposed to  $\text{BF}_3\cdot\text{OEt}_2$ , the *syn*-oxaspiropentane-azide **5a** in DCM at  $-78^\circ\text{C}$  underwent domino semipinacol–Schmidt rearrangement to give the corresponding bridged azatricyclic lactam **4a** in 25% yield (Scheme 3).



**Scheme 3** Domino semipinacol–Schmidt reaction of *syn*-oxaspiropentane-azide **5a**.

Encouraged by this preliminary observation, this novel cyclization was carried out with different Lewis acids and the results obtained are summarized in Table 1.

Among the Lewis acids screened,  $\text{TiCl}_4$  was found to be the most efficient catalyst to bring about this domino transformation and furnished the corresponding bridged azatricyclic ABC-core **4a** of immunosuppressant FR901483 (**1**) in a stereoselective manner, which on subsequent reduction with  $\text{LiAlH}_4$  furnished the known bridged azatricyclic system **3a**, whose mass and NMR spectral data are found to be in complete agreement with the literature values.<sup>7a</sup> Similarly, the mixture of *syn*- and *anti*-oxaspiropentane-azide **5b** on

**Table 1** Domino semipinacol–Schmidt reaction of *syn*-oxaspiropentane-azide **5a** with different Lewis acids<sup>a</sup>


Entry	Lewis acid	Time/h	Yield of <b>4a</b> <sup>b</sup> (%)
1	TMSOTf	4	59
2	$\text{EtAlCl}_2$	6	62
3	$\text{TiCl}_4$	4	78

<sup>a</sup> Reactions were performed using 2.5 equiv. of Lewis acid. <sup>b</sup> Isolated yield.

**Table 2**  $\text{TiCl}_4$ -promoted domino semipinacol–Schmidt reaction of oxaspiropentane-azide **5**<sup>a</sup>

Entry	Oxaspiropentane-azide <b>5</b>	Ratio of <b>5</b> <sup>b</sup> ( <i>syn/anti</i> )	Azatricyclic lactam <b>4</b> <sup>c</sup> (% yield)	Azatricyclic amine <b>3</b> <sup>c</sup> (% yield)
1		1 : 0		
2		1 : 1		
3		5 : 1		

<sup>a</sup> Reactions were performed using 2.5 equiv. of  $\text{TiCl}_4$ . <sup>b</sup> *Syn*- and *anti*-mixture was used in the reaction. <sup>c</sup> Yield is shown in parentheses. <sup>d</sup> Isolated yield based on the *syn*-isomer.



**Fig. 3** ORTEP-diagram of thiolactam derivative (±)-**10b** of the ABC ring system of daphlongeranine B.


exposure to  $\text{TiCl}_4$  resulted in the corresponding bridged azatricyclic ABC-core **4b** of daphlongeranine B (**2**) in 80% yield, based on the *syn*-isomer (Table 2).<sup>15</sup> The structure and the relative stereochemistry of the cyclized product **4b** was unambiguously confirmed by single crystal X-ray analysis of the corresponding thiolactam derivative (±)-**10b** (Fig. 3). Interestingly, this is the first stereoselective approach for the construction of the bridged azatricyclic ABC-core of daphlongeranine B (**2**). Under similar cyclization conditions, the *syn*- and *anti*-mixture of oxaspiropentane-azide **5c** afforded the corresponding bridged azatricyclic lactam **4c** in 80% yield based on the *syn*-isomer. Reduction of cyclized azatricyclic lactams **4b** and **4c** afforded the corresponding bridged azatricyclic amines **3b** and **3c**, respectively, in excellent yields (Table 2).

The scope of this novel stereoselective transformation was further explored in the enantioselective construction of the bridged azatricyclic ABC-core of FR901483. The asymmetric

**Table 3** Enantioselective synthesis of the bridged azatricyclic system

S. no.	Michael adduct <b>9</b>	ee <sup>a</sup> (%)	Azatricyclic lactam <b>4</b> <sup>b</sup>	ee <sup>a</sup> (%)
1		99		99
	(-)- <b>9a</b> [α] <sub>D</sub> <sup>25</sup> -3.4 (c 0.1, CHCl <sub>3</sub> )		(-)- <b>4a</b> [α] <sub>D</sub> <sup>26</sup> -38.0 (c 1.0, CHCl <sub>3</sub> )	
2		99		99
	(-)- <b>9c</b> [α] <sub>D</sub> <sup>25</sup> -41.0 (c 1.0, CHCl <sub>3</sub> )		(-)- <b>4c</b> [α] <sub>D</sub> <sup>23</sup> -32.0 (c 1.0, CHCl <sub>3</sub> )	

<sup>a</sup> % ee calculated using chiral HPLC. <sup>b</sup> HPLC analysis was done on the corresponding thiolactam derivative.

**Scheme 4** Plausible mechanism for the domino semipinacol-Schmidt reaction of *syn*-oxaspiropentane-azide.

Michael addition of dimethyl malonate with cyclohexenone and cycloheptenone in the presence of Shibasaki (*S*)-ALB catalyst<sup>16</sup> furnished the corresponding adducts (–)-**9a** and (–)-**9c**, respectively, in 99% ee. Following a similar sequence of reactions as shown in Scheme 2, the Michael adducts were further converted to the azatricyclic lactam (–)-**4a** and (–)-**4c**, respectively, in good yields (Table 3). The azatricyclic lactam (–)-**4a** is the enantiomer of the ABC core of FR901483.<sup>17</sup>

A plausible mechanism for the formation of a bridged azatricyclic framework from *syn*-oxaspiropentane-azide **5** via domino semipinacol-Schmidt cyclization is depicted in Scheme 4.

In summary, a novel and general approach for the stereo- and enantioselective construction of bridged azatricyclic ring systems having an azaquaternary center has been developed based on a domino semipinacol-Schmidt reaction. This new method has provided an elegant entry for the compact synthesis of the ABC-core of the biologically significant alkaloids such as FR901483 and daphlongeranine B. Since our approach is simple and effective, it can be readily implemented in the stereo- and enantioselective synthesis of natural products possessing bridged aza-polycyclic frameworks.

We thank DST (New Delhi) for financial support and DST-FIST (New Delhi) for the infrastructure facilities. M. P. (SRF) thanks CSIR (New Delhi) for research fellowship. Authors thank Mr V. Ramkumar for single crystal X-ray analysis.

## Notes and references

- For reviews on aza-polycyclic systems, see: (a) R. A. Pilli, G. B. Rosso and M. C. F. de Oliveira, *Nat. Prod. Rep.*, 2010, **27**, 1908; (b) J. Bonjoch, F. Diaba and B. Bradshaw, *Synthesis*, 2011, 993.
- For a recent review on the *Daphniphyllum* alkaloids: J. Kobayashi and T. Kubota, *Nat. Prod. Rep.*, 2009, **26**, 936.
- For recent reviews on the *Lycopodium* alkaloids, see: (a) Y. Hirasawa, J. Kobayashi and H. Morita, *Heterocycles*, 2009, **77**, 679; (b) M. Kitajima and H. Takayama, *Lycopodium Alkaloids: Isolation and Asymmetric Synthesis*, in *Topics in Current Chemistry*, ed. H.-J. Knölker, Springer, Berlin, 2011.
- (a) H. Li, X. Wang and X. Lei, *Angew. Chem., Int. Ed.*, 2012, **51**, 491; (b) X. M. Zhang, Y. Q. Tu, F. M. Zhang, H. Shao and X. Meng, *Angew. Chem., Int. Ed.*, 2011, **50**, 3916.
- (a) A. S. Kende, T. L. Smalley, Jr. and H. Huang, *J. Am. Chem. Soc.*, 1999, **121**, 7431; (b) M. H. Becker, P. Chua, R. Downham, C. J. Douglas, N. K. Garg, S. Hiebert, S. Jaroch, R. T. Matsuoka, J. A. Middleton, F. W. Ng and L. E. Overman, *J. Am. Chem. Soc.*, 2007, **129**, 11987; (c) B. B. Liao and M. D. Shair, *J. Am. Chem. Soc.*, 2010, **132**, 9594.
- For the total synthesis of FR901483, see: (a) B. B. Snider and H. Lin, *J. Am. Chem. Soc.*, 1999, **121**, 7778; (b) G. Scheffler, H. Seike and E. J. Sorensen, *Angew. Chem., Int. Ed.*, 2000, **39**, 4593; (c) M. Ousmer, N. A. Braun and M. A. Ciufolini, *Org. Lett.*, 2001, **3**, 765; (d) J. H. Maeng and R. L. Funk, *Org. Lett.*, 2001, **3**, 1125; (e) C. A. Carson and M. A. Kerr, *Org. Lett.*, 2009, **11**, 777.
- Approaches to the core structure of FR901483, see: (a) N. Yamazaki, H. Suzuki and C. Kibayashi, *J. Org. Chem.*, 1997, **62**, 8280; (b) D. J. Wardrop and W. Zhang, *Org. Lett.*, 2001, **3**, 2353; (c) J. E. Kropf, I. C. Meigh, M. W. P. Bebbington and S. M. Weinreb, *J. Org. Chem.*, 2006, **71**, 2046; (d) S. Kaden and H. U. Reissig, *Org. Lett.*, 2006, **8**, 4763; (e) D. B. Gotchev and D. L. Comins, *J. Org. Chem.*, 2006, **71**, 9393; (f) S. T. M. Simila and S. F. Martin, *J. Org. Chem.*, 2007, **72**, 5342.
- For recent reports on the construction of the core structure of *Daphniphyllum* alkaloids see: F. Sladojevich, I. N. Michaelides, B. Darses, J. W. Ward and D. J. Dixon, *Org. Lett.*, 2011, **13**, 5132 and references cited therein.
- For the construction of linearly fused azatricyclic ring systems, see: (a) R. A. Pilli and M. C. F. de Oliveira, *Nat. Prod. Rep.*, 2000, **17**, 117; (b) R. Alibès and M. Figueredo, *Eur. J. Org. Chem.*, 2009, 2421; (c) Y. Wang, L. Zhu, Y. Zhang and R. Hong, *Angew. Chem., Int. Ed.*, 2011, **50**, 2787.
- For the construction of angularly fused azatricyclic ring systems, see: (a) J. Aubé and G. L. Milligan, *J. Am. Chem. Soc.*, 1991, **113**, 8965; Y. M. Zhao, P. Gu, Y. Q. Tu, C. A. Fan and Q. Zhang, *Org. Lett.*, 2008, **10**, 1763; (b) A. M. Meyer, C. E. Katz, S. W. Li, D. V. Velde and J. Aubé, *Org. Lett.*, 2010, **12**, 1244; Z. H. Chen, Y. Q. Tu, S. Y. Zhang and F. M. Zhang, *Org. Lett.*, 2011, **13**, 724; (c) A. C. Flick, M. J. A. Cabellero, H. I. Lee and A. Padwa, *J. Org. Chem.*, 2010, **75**, 1992.
- For intramolecular Schmidt reaction of azido-epoxides, see: (a) P. G. Reddy, B. Varghese and S. Baskaran, *Org. Lett.*, 2003, **5**, 583; (b) P. G. Reddy and S. Baskaran, *J. Org. Chem.*, 2004, **69**, 3093; (c) P. G. Reddy, M. G. Sankar and S. Baskaran, *Tetrahedron Lett.*, 2005, **46**, 4559.
- S. Lang, A. R. Kennedy, J. A. Murphy and A. H. Payne, *Org. Lett.*, 2003, **5**, 3655.
- For domino reactions in organic synthesis, see: L. F. Tietze, G. Brasche and K. Gericke, *Domino Reactions in Organic Synthesis*, Wiley-VCH, Weinheim, 2006.
- B. M. Trost and M. J. Bogdanowicz, *J. Am. Chem. Soc.*, 1973, **95**, 5321.
- Due to geometrical constraints, only the *syn*-isomer is anticipated to undergo intramolecular Schmidt cyclization.
- S. Shimizu, K. Otori, T. Arai, H. Sasai and M. Shibasaki, *J. Org. Chem.*, 1998, **63**, 7547.
- In the asymmetric Michael addition step, use of Shibasaki (*R*)-ALB catalyst,<sup>16</sup> in place of (*S*)-ALB catalyst, would lead to the natural enantiomer of the ABC core of FR901483.