

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/267760564>

# ChemInform Abstract: One Pot Synthesis of 1,2,4,5-Tetrasubstituted-Imidazoles Catalyzed by Trityl Chloride in Neutral Media.

ARTICLE *in* RSC ADVANCES · NOVEMBER 2014

Impact Factor: 3.84 · DOI: 10.1039/C4RA10589C

CITATIONS

3

READS

99

## 5 AUTHORS, INCLUDING:



**Mohammad Ali Zolfigol**

Bu-Ali Sina University

574 PUBLICATIONS 7,701 CITATIONS

SEE PROFILE



**Ahmad Reza Moosavi-Zare**

Sayyed Jamaledin Asadabadi University

121 PUBLICATIONS 1,212 CITATIONS

SEE PROFILE



**Mohsen Shekouhy**

Shiraz University

41 PUBLICATIONS 721 CITATIONS

SEE PROFILE

## PAPER

CrossMark  
click for updatesCite this: *RSC Adv.*, 2014, 4, 60636

## One pot synthesis of 1,2,4,5-tetrasubstituted-imidazoles catalyzed by trityl chloride in neutral media†

Ahmad Reza Moosavi-Zare,<sup>\*a</sup> Zhila Asgari,<sup>b</sup> Abdolkarim Zare,<sup>\*c</sup>  
Mohammad Ali Zolfigol<sup>\*b</sup> and Mohsen Shekouhy<sup>d</sup>

Received 16th September 2014

Accepted 31st October 2014

DOI: 10.1039/c4ra10589c

www.rsc.org/advances

Trityl chloride (TrCl or Ph<sub>3</sub>CCl) efficiently catalyzes the one-pot multi-component condensation of benzil with aldehydes, primary amines and ammonium acetate under neutral and solvent-free conditions to give 1,2,4,5-tetrasubstituted imidazoles in high to excellent yields and in short reaction times. Mechanistically, it is attractive that trityl chloride promotes the reaction by *in situ* generation of trityl carbocation (Ph<sub>3</sub>C<sup>+</sup>).

Multi-component reactions (MCRs) play an important role in combinatorial chemistry because of their capability to synthesize target molecules with atom economy and high efficiency, by reaction of three or more reactants in one step. Furthermore, these reactions increase synthetic efficacy and simplicity with respect to conventional organic transformations.<sup>1</sup>

Imidazole derivatives are one of the most important classes of nitrogen-containing five-members heterocycles. For example, they are an essential component of various biological and pharmaceutical molecules, including histidine, histamine, biotin, losartan, olmesartan, eprosartan, miconazole, ketoconazole, clotrimazole and trifluoragrel.<sup>2</sup> Some pharmaceutical compounds, based on imidazole moiety, are displayed in Fig. 1. Moreover, imidazole derivatives are used as green solvents in form of ionic liquids,<sup>3</sup> and as N-heterocyclic carbenes in organometallic chemistry.<sup>4</sup> 1,2,4,5-Tetrasubstituted imidazoles are an important class of imidazoles which are prepared *via* the one-pot multi-component condensation of benzil with aldehydes, primary amines and ammonium acetate using acidic catalysts.<sup>5</sup> Although several catalysts for this transformation are known, development of newer catalysts with high novelty which can promote the reaction attract attention for their difference with the others, and effectiveness.

Currently, development of organocatalysts has attracted much attention in organic synthesis, especially from green chemistry point of view.<sup>6</sup> These catalysts have some advantages compared with acidic catalysts or precious metal, such as commercial availability, low cost, relative non-toxicity, air

stability, need to simple reaction conditions, and ability to promote a chemical reaction *via* different activation modes.<sup>7,8</sup> Considering the high importance of organocatalysts, recently, we have applied a neutral and attractive kind of these catalysts namely triarylmethyl chlorides (Ar<sub>3</sub>CCl) to promote the preparation of 12-aryl-8,9,10,12-tetrahydrobenzo[*a*]-xanthen-11-ones,<sup>7</sup> bis(indolyl)methanes,<sup>9</sup> 1-amidoalkyl-2-naphthols<sup>10</sup> and N-sulfonyl imines.<sup>11</sup> In the mentioned investigations, we have used Ar<sub>3</sub>CCl to catalyze the two and three-component reactions.




Fig. 1 The structure of losartan, olmesartan, eprosartan, miconazole, ketoconazole, clotrimazole and trifluoragrel as drugs.

<sup>a</sup>Department of Chemistry, University of SayyedJamaledinAsadabadi, Asadabad, 6541835583, Iran. E-mail: moosavizare@yahoo.com

<sup>b</sup>Faculty of Chemistry, Bu-Ali Sina University, Hamedan, 6517838683, Iran

<sup>c</sup>Department of Chemistry, Payame Noor University, PO Box 19395-3697, Tehran, Iran

<sup>d</sup>Department of Chemistry, College of Science, Shiraz University, 71454, Shiraz, Iran

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c4ra10589c



Scheme 1 The preparation of 1,2,4,5-tetrasubstituted imidazoles.

Table 1 Effect of the catalyst amount and temperature on the reaction between benzil, benzaldehyde, aniline and ammonium acetate

Mol% of TrCl	Temp. (°C)	Time (min)	Yield <sup>a</sup> (%)
7	90	30	61
10	50	30	37
10	80	30	64
10	90	30	79
10	100	30	79

<sup>a</sup> Isolated yield.

However, in the presented work, we employ trityl chloride (TrCl or Ph<sub>3</sub>CCl) as a homogeneous organocatalyst to promote a one-pot four-component reaction, *i.e.* the condensation of benzil with aldehydes, primary amines and ammonium acetate, for the first time (Scheme 1).

To optimize the reaction conditions, as a model reaction, the condensation of benzil (1 mmol) with benzaldehyde (1 mmol), aniline (1 mmol) and ammonium acetate (1 mmol) was studied in the presence of different molar ratios of trityl chloride at range of 50–100 °C under solvent-free conditions; the respective results are summarized in Table 1. As Table 1 indicates, the best results were obtained when the reaction was performed using 10 mol% of the organocatalyst at 90 °C. To confirm that heat can not thermodynamically promote the reaction in the absence

of catalyst; the model reaction was examined at 90 °C under catalyst-free conditions. In these conditions, the desired product was obtained in trace yield after 120 min. This observation clearly showed that heat can not promote the reaction without catalyst; thus, presence of catalyst (*e.g.* TrCl) is essential for the reaction.

After optimization of the reaction conditions, the efficiency and scope of the organocatalyst was evaluated by the reaction of benzil with various arylaldehydes, different primary amines and ammonium acetate using 10 mol% of TrCl at 90 °C; the corresponding results are summarized in Table 2. As it can be seen in Table 2, all reactions proceeded efficiently to afford the desired 1,2,4,5-tetrasubstituted imidazoles in high yields and in short reaction times. Thus, TrCl was efficient to catalyze the one-pot four-component reaction.

In a plausible mechanism which is supported by the literature (Scheme 2),<sup>1b,9–11</sup> at first, aldehyde is activated by Ph<sub>3</sub>C<sup>+</sup> generated from Ph<sub>3</sub>CCl; consequently, two resonance forms of the activated aldehyde (**I** and **II**) can produce.<sup>1b,9–11</sup> The activated aldehyde react with amine to give iminium intermediate **III**. Afterward, **III** reacts with ammonia (resulted from ammonium acetate) to afford **IV**, and this intermediate reacts with trityl alcohol (Ph<sub>3</sub>COH) to provide **V** and Ph<sub>3</sub>CCl. In the next step, intermediate **VI** produces by nucleophilic attack of the nitrogens of **V** to the carbonyl groups of benzil. Finally, 1,2,4,5-tetrasubstituted imidazole is synthesized by removing two molecules of H<sub>2</sub>O from **VI**.


To confirm the generation of **I** and **II** as resonance structures of the activated aldehyde, benzaldehyde was treated with trityl chloride at room temperature, and IR, <sup>1</sup>H NMR, and UV spectra of the mixture was run;<sup>1b,10</sup> the obtained results include:

IR (nujol): ν<sub>max</sub> (cm<sup>−1</sup>) of the carbonyl group of benzaldehyde (1705) was decreased to 1697 in the mixture of benzaldehyde and trityl chloride (Fig. S1†).

Table 2 The synthesis of 1,2,4,5-tetrasubstituted imidazoles using TrCl (10 mol%) under solvent-free conditions at 90 °C

Entry	Ar	R	Time (min)/yield <sup>a</sup> (%)	M.p. (°C) found/reported
1a	C <sub>6</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub>	30/79	218–220/218–221 (ref. 5a)
1b	C <sub>6</sub> H <sub>5</sub>	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub>	32/80	156–158/156–159 (ref. 5a)
1c	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub>	35/76	165–167/165–168 (ref. 5a)
1d	4-Cl-C <sub>6</sub> H <sub>4</sub>	4-Cl-C <sub>6</sub> H <sub>4</sub>	20/81	188–190/189–191 (ref. 4b)
1f	3-NO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	60/78	146–147/145–147 (ref. 4b)
1g	4-NO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	30/81	214–217/215–217 (ref. 4b)
1h	4-Cl-C <sub>6</sub> H <sub>4</sub>	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub>	30/79	158–160/161–163 (ref. 4b)
1i	4-Cl-C <sub>6</sub> H <sub>4</sub>	4-F-C <sub>6</sub> H <sub>4</sub>	20/78	197–200/198–201 (ref. 4b)
1j	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	60/75	193–195/194–196 (ref. 4b)
1k	3-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub>	40/75	131–133/130–132 (ref. 5a)
1l	4-Cl-C <sub>6</sub> H <sub>4</sub>	C <sub>6</sub> H <sub>5</sub>	30/79	148–150/148–151 (ref. 5a)
1m	4-OH-C <sub>6</sub> H <sub>4</sub>	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub>	35/78	134–137/135–138 (ref. 5a)
1n	4-CH <sub>3</sub> O-C <sub>6</sub> H <sub>4</sub>	C <sub>6</sub> H <sub>5</sub> CH <sub>2</sub>	45/75	158–160/158–161 (ref. 5a)
1o	4-OH-C <sub>6</sub> H <sub>4</sub>	C <sub>6</sub> H <sub>5</sub>	35/81	280–283/282–284 (ref. 5a)
1p	4-CN-C <sub>6</sub> H <sub>4</sub>	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	30/82	197–200/198–201 (ref. 4b)
1q	4-OH-C <sub>6</sub> H <sub>4</sub>	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	35/82	232–234/233–235 (ref. 5a)
1r	2-Thienyl	4-CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	20/78	198–200/199–202 (ref. 4b)
1s	2-Thienyl	4-OH-C <sub>6</sub> H <sub>4</sub>	3/92	198–200/198–201 (ref. 4b)

<sup>a</sup> Isolated yield.



Scheme 2 The proposed mechanism for the synthesis of 1,2,4,5-tetrasubstituted imidazoles using trityl chloride.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ): chemical shift (ppm) of the  $\text{HC}=\text{O}$  of benzaldehyde (9.78) was increased to 10.04 in the mixture of  $\text{PhCHO}$  and  $\text{TrCl}$  (Fig. S2†).

UV: maximum absorption of benzaldehyde and trityl chloride were 240 and 222 nm, respectively. However,  $\lambda_{\text{max}}$  of the mixture of  $\text{PhCHO}$  and  $\text{TrCl}$  was observed in 245 nm (Fig. S3†).

These observations showed that aldehyde in the presence of trityl chloride give resonance forms **I** and **II**.

Although  $\text{TrOH}$  produced during the reaction; but it can not catalyze the reaction like  $\text{TrCl}$ . To confirm this point, we tested the model reaction in the presence of  $\text{TrOH}$  as catalyst in which the product yield was only 23% after 50 min. This result verified that  $\text{TrOH}$  is changed to  $\text{TrCl}$  during the reaction.

Thus, the mechanism is confirmed by the above observations besides the literature.<sup>1b,7,9–11</sup>

Catalytic application of other trityl sources such as monomethoxytrityl chloride ( $\text{Ph}_2(p\text{-MeOC}_6\text{H}_4)\text{CCl}$ ,  $\text{MMTCl}$ ), dimethoxytrityl chloride ( $\text{Ph}(p\text{-MeOC}_6\text{H}_4)_2\text{CCl}$ ) and triphenylmethanol ( $\text{TrOH}$ ) on the model reaction, was also investigated under




Fig. 2 Catalytic activity of different triarylmethyl sources (10 mol%) on the reaction between benzil (1 mmol), benzaldehyde (1 mmol), aniline (1 mmol) and ammonium acetate (1 mmol) at 90 °C.

solvent-free conditions at 90 °C. The results are shown in Fig. 2. As Fig. 2 indicates, higher yield and shorter reaction time were obtained using trityl chloride. Furthermore, turn over numbers (TON) and turn over frequencies (TOF) on the effectiveness of these catalysts for the synthesis of the imidazoles were calculated (Fig. 2). The TON and TOF values showed that  $\text{TrCl}$  is more efficient than  $\text{MMTCl}$ ,  $\text{DMTCl}$  and  $\text{TrOH}$ .

## Conclusions

In summary, we have applied  $\text{TrCl}$  as a neutral and homogeneous organocatalyst for the one-pot four-component reaction of aldehydes with amines (aliphatic or aromatic), benzil and ammonium acetate at 90 °C in solvent-free conditions to furnish 1,2,4,5-tetrasubstituted imidazoles. The promising points for the presented catalytic process include generality, efficacy, high yield, simplicity, cleaner reaction profile, relatively short reaction times and good agreement with the green chemistry protocols.<sup>12</sup>

## Acknowledgements

The authors gratefully acknowledge support of this work by the Research Affairs Office of Bu-Ali Sina University (Grant number 32-1716 entitled development of chemical methods, reagents and molecules), University of Sayyed Jamaledin Asadabadi, Payame Noor University, and Center of Excellence in Development of Environmentally Friendly Methods for Chemical Synthesis (CEDEFMCS), for providing support to this work.

## Notes and references

- (a) *Multicomponent Reactions*, ed. J. Zhu and H. Bienayme, Wiley, Weinheim, 2005; (b) A. Khazaei, M. A. Zolfigol, A. R. Moosavi-Zare, F. Abi, A. Zare, H. Kaveh, V. Khakyzadeh, M. Kazem-Rostami, A. Parhami and H. Torabi-Monfared, *Tetrahedron*, 2013, **69**, 212; (c) A. R. Moosavi-Zare, M. A. Zolfigol, S. Farahmand, A. Zare, A. R. Pourali and R. Ayazi-Nasrabadi, *Synlett*, 2014, **25**, 193; (d) M. A. Zolfigol, A. Khazaei, A. R. Moosavi-Zare, A. Zare and V. Khakyzadeh, *Appl. Catal.*, A, 2011, **400**, 70; (e) A. R. Moosavi-Zare, M. A. Zolfigol and M. Daraei, *Synlett*, 2014, **25**, 1173.
- (a) S. A. Laufer, W. Zimmermann and K. J. Ruff, *J. Med. Chem.*, 2004, **47**, 6311; (b) C. Leister, Y. Wang, Z. Zhao and C. W. Lindsley, *Org. Lett.*, 2004, **6**, 1453.
- P. Wasserscheid and T. Welton, *Ionic Liquids in Synthesis*, Wiley-VCH, Weinheim, 2008.
- (a) L. H. Lee, M. Bang and C. S. Pak, *Tetrahedron Lett.*, 2005, **46**, 7139; (b) A. Hasaninejad, A. Zare, M. Shekouhi and J. Ameri Rad, *J. Comb. Chem.*, 2010, **12**, 844.
- (a) M. A. Zolfigol, A. Khazaei, A. R. Moosavi-Zare, A. Zare, Z. Asgari, V. Khakyzadeh and A. Hasaninejad, *J. Ind. Eng. Chem.*, 2013, **19**, 721; (b) A. Y. Usyatinsky and Y. L. Khmelnitsky, *Tetrahedron Lett.*, 2000, **41**, 5031; (c) S. Kantevari, S. V. N. Vuppapapati, D. O. Biradar and L. Nagarapu, *J. Mol. Catal. A: Chem.*, 2007, **266**, 109; (d) M. Kidwai, P. Mothra, V. Bansal, R. K. Somvanshi, A. S. Ethayathulla, S. Dey and T. P. Singh, *J. Mol. Catal. A: Chem.*, 2007, **265**, 177; (e) B. Sadeghi, B. B. F. Mirjalili and M. M. Hashemi, *Tetrahedron Lett.*, 2008, **49**, 2575; (f) S. D. Sharma, P. Hazarika and D. Konwar, *Tetrahedron Lett.*, 2008, **49**, 2216; (g) A. R. Moosavi-Zare, M. A. Zolfigol, V. Khakyzadeh, C. Böttcher, M. H. Beyzavi, A. Zare, A. Hasaninejad and R. Luque, *J. Mater. Chem. A*, 2014, **2**, 770.
- A. Dondoni and A. Massi, *Angew. Chem., Int. Ed.*, 2008, **47**, 4638.
- A. Khazaei, M. A. Zolfigol, A. R. Moosavi-Zare, A. Zare, M. Khojasteh, Z. Asgari, V. Khakyzadeh and A. Khalafi-Nezhad, *Catal. Commun.*, 2012, **20**, 54.
- S. Verma, L. S. Jain and B. Sain, *Tetrahedron Lett.*, 2010, **51**, 6897.
- A. Khalafi-Nezhad, A. Parhami, A. Zare, A. R. Moosavi Zare, A. Hasaninejad and F. Panahi, *Synthesis*, 2008, 617.
- A. Khazaei, M. A. Zolfigol, A. R. Moosavi-Zare, A. Zare, A. Parhami and A. Khalafi-Nezhad, *Appl. Catal.*, A, 2010, **386**, 179.
- A. Khalafi-Nezhad, A. Parhami, A. Zare, A. Nasrollahi Shirazi, A. R. Moosavi-Zare and A. Hasaninejad, *Can. J. Chem.*, 2008, **86**, 456.
- General procedure for the synthesis of 1,2,4,5-tetrasubstituted imidazoles: a mixture of compounds consisting of benzil (1 mmol), aldehyde (1 mmol), primary amine (1 mmol), ammonium acetate (1 mmol) and TrCl (10 mol%) in a 10 mL round-bottomed flask connected to a reflux condenser, was stirred in an oil-bath (90 °C). After completion of the reaction, as monitored by TLC, the reaction mixture was cooled to room temperature, and then petroleum ether (5 mL) was added to the crude reaction mixture, and filtered to separate the catalyst which will be recycled for other reactions. Then, the pure product was obtained by recrystallization of the reaction mixture in ethanol (90%) or ethyl acetate.