

4,5-Dihydrothiepin 1,1-Dioxide

By WILLIAM L. MOCK

(Department of Chemistry, Carnegie-Mellon University, Pittsburgh, Pennsylvania, 15213)


Summary The reaction between sulphur dioxide and two equivalents of vinyldiazomethane yields 4,5-dihydrothiepin 1,1-dioxide: a probable mechanism is discussed.

THE reaction of a diazoalkane with a sulphenel¹ (produced either by dehydrohalogenation of a sulphonyl halide or from another diazoalkane molecule and sulphur dioxide) at present provides the only general preparative route to thiiran 1,1-dioxides (equation 1). In the course of searching for new synthetic routes to medium-sized ring sulphones, we have examined the reaction between vinyldiazomethane and sulphur dioxide.

Passage of gaseous sulphur dioxide into an ethereal solution of vinyldiazomethane² results in rapid disappearance of the characteristic red colour of the latter and evolution of N₂. Work-up of the resulting solution by evaporation and chromatography of the residue on silicic acid yields a crystalline sulphone, which, after sublimation (100°/0.1 mm) and recrystallization from benzene-hexane, melts at 111.5–112.5°.

The composition of this new sulphone is C₆H₈SO₂ as shown by elemental analysis and mass spectroscopy. I.r. absorption bands at 7.8 and 9.0 μm establish the presence of the sulphonyl group. Catalytic hydrogenation (Pd-C) results in the uptake of two equivalents of hydrogen and yields thiepan 1,1-dioxide. Treatment of the new substance in benzene under reflux with triethylamine as catalyst, effects quantitative conversion into 2,7-dihydrothiepin 1,1-dioxide,† which has been prepared by 1,6-cycloaddition of

sulphur dioxide to *cis*-hexatriene.³ Hence, the new material is probably an isomeric (and thermodynamically less stable) dihydrothiepin dioxide. Its relatively simple n.m.r. spectrum suggests that it is 4,5-dihydrothiepin 1,1-dioxide (II).




There is a four proton multiplet at δ 2.70, assigned to the γ-methylene groups at positions 4 and 5 shown by decoupling experiments to be adjacent to the high-field pair of a set of four olefinic protons at δ 6.58 and 6.44. These are assigned, respectively, to the α- and β-hydrogens, which appear to be strongly coupled to each other (*J* 11.0 ± 0.1 Hz).

A possible mechanism for the formation of the observed product (II) involves the steps of the ordinary diazoalkane-sulphur dioxide reaction (equation 2), producing *cis*-

† When the isomerization is followed by n.m.r. no intermediates can be detected.

divinylthiiran 1,1-dioxide (I) as a transient intermediate.[‡] On the basis of close analogies,⁴ rapid Cope rearrangement to 4,5-dihydrothiepin 1,1-dioxide would be anticipated. Alternative mechanisms involving mesoionic intermediates or various types of conjugate addition cannot be ruled out. If the suggested mechanism is tentatively accepted, a new mode of reactivity is revealed for the highly unstable thiiran dioxide ring, which normally fragments to an olefin and sulphur dioxide.

This work was supported by the National Science Foundation.



(Received, July 27th, 1970; Com. 1234.)

[‡] The *trans*-isomer, if formed and if not in equilibrium with the *cis*-isomer, may contribute to the tar which comprises the main reaction product.

¹ G. Optiz and K. Fischer, *Angew. Chem.*, 1965, **77**, 41; G. Hesse, E. Reichold, and S. Majmudar, *Chem. Ber.*, 1957, **90**, 2106; G. Hesse and S. Majmudar, *ibid.*, 1960, **93**, 1129.

² C. D. Hurd and S. C. Lui, *J. Amer. Chem. Soc.*, 1935, **57**, 2656; D. W. Adamson and J. Kenner, *J. Chem. Soc.*, 1935, 286. We prepared the diazoalkane from *N*-allyl-*N*-nitrosotoluene-*p*-sulphonamide and base.

³ W. L. Mock, *J. Amer. Chem. Soc.*, 1967, **89**, 1281.

⁴ E. Vogel, *Angew. Chem. Internat. Edn.*, 1962, **1**, 53; 1963, **2**, 1; E. L. Stogryn, M. H. Gianni, and A. J. Passannante, *J. Org. Chem.*, 1964, **29**, 1275; R. A. Braun, *ibid.*, 1963, **28**, 1383; E. L. Stogryn and S. J. Brois, *ibid.*, 1965, **30**, 88.