

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51477606>

Total Synthesis of (-)-CP2-Disorazole C-1

ARTICLE *in* ORGANIC LETTERS · AUGUST 2011

Impact Factor: 6.36 · DOI: 10.1021/ol2015994 · Source: PubMed

CITATIONS

10

READS

57

4 AUTHORS, INCLUDING:



John C Schmitz

University of Pittsburgh

56 PUBLICATIONS 756 CITATIONS

SEE PROFILE

Published in final edited form as:

Org Lett. 2011 August 5; 13(15): 4088–4091. doi:10.1021/ol2015994.

Total synthesis of (-)-CP₂-disorazole C₁

Chad D. Hopkins[†], John C. Schmitz^{#,§}, Edward Chu^{‡,§}, and Peter Wipf^{†,§}

Peter Wipf: pwipf@pitt.edu


[†]Department of Chemistry, University of Pittsburgh, Pittsburgh, PA 15260

[#]VACT Cancer Research Laboratory, VACT Healthcare System, West Haven, CT 06516

[‡]Division of Hematology/Oncology, Department of Medicine and University of Pittsburgh Cancer Institute

[§]Molecular Therapeutics Drug Discovery Program Pittsburgh, PA 15232, USA

Abstract



The total synthesis of a bis-cyclopropane analog of the antimitotic natural product (-)-disorazole C₁ was accomplished in 23 steps and 1.1% overall yield. A vinyl cyclopropane cross-metathesis reaction generated a key (*E*)-alkene segment of the target molecule. IC₅₀ determinations of (-)-CP₂-disorazole C₁ in human colon cancer cell lines indicated low nanomolar cytotoxic properties. Accordingly, this synthetic bioisostere represents the first biologically active disorazole analog not containing a conjugated diene or polyene substructure element.

The disorazoles comprise a family of ~30 closely related polyketide microtubule disruptors isolated since 1994 from the fermentation broth of the myxobacterium *Sorangium cellulosum* by Jansen, Reichenbach, Höfle and co-workers.^{1,2} Members of the disorazole class have displayed anti-cancer activity in the nano- and picomolar range against a variety of transformed cell lines, including multi-drug resistant cells.³ To date, only (-)-disorazole C₁ has yielded to total synthesis,⁴ although several simplified analogs and segments have been reported.⁵ As part of our investigations of structure-activity relationships (SAR) of biologically active natural products,^{2d,6} we designed a cyclopropane analog of (-)-disorazole C₁ (i.e., CP₂-disorazole C₁), with the goal to replace the labile⁷ (*E,Z,Z*)-triene subunit with an isosteric and biologically similarly effective moiety. The choice of absolute configuration for the *cis*-cyclopropanes in CP₂-disorazole C₁ reflected the stereochemistry observed for the epoxide present in disorazole A₁.^{2a}

Retrosynthetically, the C₂-symmetrical macrodiolide CP₂-disorazole C₁ (**1**) could be constructed from two key fragments, **15** and **9** (Scheme 1). Diene **15** is derived from β-hydroxy ester **12** which can be obtained from an asymmetric Mukaiyama aldol reaction of

Correspondence to: Peter Wipf, pwipf@pitt.edu.

Supporting Information **Available:** Experimental procedures and spectral data for all new compounds, including copies of ¹H and ¹³C NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>

crotonaldehyde. By taking advantage of a vinyl cyclopropane cross-metathesis reaction,⁸ our plan was to access alkene **9** from the readily available cyclopropane **5** and the oxazole precursor **3**. This approach was realized in excellent yields beginning with the straightforward preparation of ester **3** in two steps from the known⁹ alkyne **2** (Scheme 2).

The vinyl cyclopropane **5** was prepared in 74% yield in a Wittig condensation from known aldehyde **4**^{5d} (Scheme 3). Cross-metathesis reaction of alkenes **5** and **3**¹⁰ in the presence of a 2nd generation ruthenium catalyst¹¹ provided alkene **6** in 70% yield as an inseparable 10:1 mixture of *E/Z*-stereoisomers. *O*-Methylation of **6** with silver oxide¹² and methyl iodide, followed by enzymatic hydrolysis¹³ with pig liver esterase (PLE) and subsequent coupling of the carboxylic acid with serine methyl ester yielded amide **8**. A two-step cyclodehydration/oxidation procedure¹⁴ led to oxazole **9**. Oxidative deprotection of the PMB ether followed by further oxidation of the resulting primary alcohol under Dess-Martin conditions afforded the desired aldehyde **10** in excellent yield.

After considerable experimentation, we found that Kiyooka's conditions were best suited for the asymmetric Mukaiyama aldol reaction between the sterically hindered silyl ketene acetal **11** and crotonaldehyde (Scheme 4).^{5a,15}

The α,α -dimethylated aldol product **12** was obtained in 79% yield and 93% *ee* as determined by chiral HPLC analysis. Protection of the secondary alcohol in **12** with 3,4-dimethoxybenzyl (DMB) bromide¹⁶ followed by ester reduction with LAH and oxidation of the primary alcohol under Swern conditions provided aldehyde **13**. Among contemporary allylation protocols,¹⁷ we found that the Duthaler-Hafner allylation¹⁸ of aldehyde **13** gave alcohol **14** in the highest (87%) yield and with a satisfactory *dr*=4.8:1.0, favoring the desired *anti*-configuration of the 1,3-diol. The minor isomer was present in diminishing amounts in subsequent intermediates until it was finally completely removed in the Wittig reaction of **17** and **10**.

At first, the advancement of homoallylic alcohol **14** to the desired phosphonium salt **17** proved problematic. Upon formation of the alkyl iodide at C12, an *in situ* S_N2-displacement of the iodide by the electron-rich DMB ether at C16 afforded the corresponding tetrahydropyran as the major product.¹⁹ Although small quantities of the desired alkyl iodide were isolated, immediate formation of the tetrahydropyran occurred upon heating. Consequently, TES protection of the alcohol at C14 followed by the exchange of the DMB with a TBS ether at C16 was used to access the orthogonally silyl-protected diol **15** (Scheme 4). Stepwise Johnson-Lemieux oxidation followed by NaBH₄ reduction afforded the primary alcohol **16**. Iodination and phosphonium salt generation at elevated temperature in a sealed vessel proceeded now without incident to furnish **17** in good overall yield. The Wittig condensation of the freshly formed phosphonium salt with aldehyde **10** in a vigorously degassed THF solution yielded 86% of the desired *Z*-alkene as a single stereoisomer by ¹H NMR analysis. Finally, selective TES deprotection with PPTs in the presence of the allylic TBS ether proceeded in 65% yield to afford the advanced segment **18**.²⁰

Originally, our end game strategy had envisioned the direct *cyclo*-dimerization of the carboxylic acid corresponding to **18**. However, all efforts to realize this *cyclo*-dimerization resulted in the exclusive formation of the 15-membered macrolactone arising from the direct lactonization at the C14 hydroxyl group, i.e. the formation of the cyclic monomer.²¹ In order to circumvent this undesired reaction, a stepwise approach was adopted (Scheme 5). Hydrolysis of ester **9** followed by DCC-mediated coupling to the secondary alcohol **18** provided the bis-cyclopropane **19** in 89% yield. Oxidative PMB deprotection with DDQ and subsequent Dess-Martin oxidation of the resulting primary alcohol led to the requisite cyclopropyl aldehyde in good yield. Condensation of this aldehyde and the ylide generated

from phosphonium salt **17** after vigorous degassing afforded the *seco*-dimer **20** in 74% yield as a single alkene stereoisomer.

Selective TES ether deprotection of **20** revealed the secondary alcohol at C14' (Scheme 5). With all functional groups properly installed, the stage was now set for the final lactone formation. Selective saponification of the methyl ester in the presence of the internal ester linkage of **20** with barium hydroxide followed by Shiina macrolactonization²² with 2-methyl-6-nitrobenzoic anhydride (MNBA) and DMAP provided the desired 30-membered macrocycle in 55% yield over the two steps. Finally, double TBS ether deprotection in the presence of HF-pyridine proceeded in 64% yield to the desired (-)-CP₂-disorazole C₁.

Table 1 illustrates our preliminary analysis of the antiproliferative properties of this novel disorazole analog. In three human colon cancer cell lines, **1** displayed low IC₅₀ values ranging from 25-50 nM, and therefore proved to be only slightly less active than its parent, (-)-disorazole C₁.^{3a-d,23} Furthermore, **1** was equipotent to the clinically used anticancer vinca alkaloid vincristine in these colon cancer cells. In contrast, cyclomonomer **21** was inactive at concentrations up to 50 μM in all three cell lines, thereby serving as a negative control and confirming the importance of the 3-D architecture of the disorazoles.^{3d}

In summary, we have developed an efficient route to (-)-CP₂-disorazole C₁ **1**, a bis-cyclopropane analog of the antimetabolic natural product disorazole C₁.²⁴ The target agent was obtained in 23 steps and 1.1% overall yield for the longest linear sequence. It contains 10 stereogenic carbons, 4 more than disorazole C₁, and 6 double bonds, 2 less than the natural product. Most importantly, by replacing the central *cis*-alkenes at C9-C10/C9'-C10' of disorazole C₁ with *cis*-cyclopropane rings, **1** no longer has a labile conjugated (*E,Z,Z*)-triene; yet, it shows nearly the same potent level of low nanomolar *in vitro* cytotoxicity in human colon cancer cell lines as the natural myxobacterium metabolite. Further biological studies of **1** will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


Acknowledgments

This work was supported in part by the NCI (CA78039) and the NSF (CHE-0910560). The authors thank Mr. Tianmin Chen (VACT Cancer Research Laboratory, West Haven, CT) for biological assays, and Profs. John Lazo (University of Virginia, Charlottesville, VA), Billy Day (University of Pittsburgh, Pittsburgh, PA) and Andreas Vogt (University of Pittsburgh, Pittsburgh, PA) for stimulating discussions and collaborative studies on antitubulin agents.


References

1. Jansen R, Irschik H, Reichenbach H, Wray V, Höfle G. Liebigs Ann Chem. 1994;759.
2. For reviews of disorazole chemistry and biology, see: (a) Hopkins CD, Wipf P. Nat Prod Rep. 2009; 26:585. [PubMed: 19387496] (b) Weissman KJ, Mueller R. Nat Prod Rep. 2010; 27:1276. [PubMed: 20520915] (c) Kretschmer M, Menche D. Synlett. 2010:2989. (d) Wipf P, Graham TH, Xiao J. Pure Appl Chem. 2007; 79:753.
3. (a) Lazo JS, Reese CE, Vogt A, Vollmer LL, Kitchens CA, Günther E, Graham TH, Hopkins CD, Wipf P. J Pharmacol Exp Ther. 2010; 332:906. [PubMed: 20008956] (b) Xu FL, Rbaibi Y, Kiselyov K, Lazo JS, Wipf P, Saunders WS. BMC Chem Biol. 2010; 10:1. [PubMed: 20181182] (c) Tierno MB, Kitchens CA, Petrik B, Graham TH, Wipf P, Xu FL, Saunders WS, Raccor BS, Balachandran R, Day BW, Stout JR, Walczak CE, Ducruet AP, Reese CE, Lazo JS. J Pharm Exp Ther. 2009; 328:715. (d) Wipf P, Graham TH, Vogt A, Sikorski RP, Ducruet AP, Lazo JS. Chem Biol Drug Des. 2006; 67:66. [PubMed: 16492150] (e) Hearn BR, Arslania RL, Fu H, Liu F, Gramajo H, Myles DC.


- J Nat Prod. 2006; 69:148. [PubMed: 16441089] (f) Elnakady YA, Sasse F, Lünsdorf H, Reichenbach H. Biochem Pharm. 2004; 67:927. [PubMed: 15104246]
4. Wipf P, Graham TH. J Am Chem Soc. 2004; 126:15346. [PubMed: 15563138]
5. (a) Schaeckel R, Hinkelmann B, Sasse F, Kalesse M. Angew Chem Int Ed. 2010; 49:1619.(b) Niess B, Hartung IV, Haustedt LO, Hoffmann HMR. Eur J Org Chem. 2006:1132.(c) Hearn BR, Arslanian RL, Fu H, Liu F, Gramajo H, Myles DC. J Nat Prod. 2006; 69:148. [PubMed: 16441089] (d) Haustedt LO, Panicker SB, Kleinert M, Hartung IV, Eggert U, Niess B, Hoffmann HMR. Tetrahedron. 2003; 59:6967.(e) Hartung IV, Eggert U, Haustedt LO, Niess B, Schaefer PM, Hoffmann HMR. Synthesis. 2003:1844.(f) Hartung IV, Niess B, Haustedt LO, Hoffmann HMR. Org Lett. 2002; 4:3239. [PubMed: 12227758] (g) Hillier MC, Price AT, Meyers AI. J Org Chem. 2001; 66:6037. [PubMed: 11529729]
6. (a) Wang Z, McPherson PA, Raccor BS, Balachandran R, Zhu G, Day BW, Vogt A, Wipf P. Chem Biol Drug Des. 2007; 70:75. [PubMed: 17683369] (b) Powis G, Wipf P, Lynch SM, Birmingham A, Kirkpatrick DL. Mol Cancer Ther. 2006; 5:630. [PubMed: 16546977] (c) Wipf P, Minion DJ, Halter RJ, Berggren MI, Ho CB, Chiang GG, Kirkpatrick L, Abraham R, Powis G. Org Biomol Chem. 2004; 2:1911. [PubMed: 15227545] (d) Wipf P, Fritch PC, Geib SJ, Seifler AM. J Am Chem Soc. 1998; 120:4105.
7. Examples documenting the lability of trienes include: (a) Burke CP, Haq N, Boger DL. J Am Chem Soc. 2010; 132:2157. [PubMed: 20108904] (b) Hicks JD, Roush WR. Org Lett. 2008; 10:681. [PubMed: 18205377] (c) Trost BM, Frederiksen MU, Papillon JPN, Harrington PE, Shin S, Shireman BT. J Am Chem Soc. 2005; 127:3666. [PubMed: 15771479] (d) Fuwa H, Sasaki M, Satake M, Tachibana K. Org Lett. 2002; 4:2981. [PubMed: 12182604]
8. Nicola T, Brenner M, Donsbach K, Kreye P. Org Proc Res Dev. 2005; 9:513.
9. (a) Salit AF, Meyer C, Cossy J, Delouvrié B, Hennequin L. Tetrahedron. 2008; 64:6684.(b) Salit AF, Meyer C, Cossy J. Synlett. 2007:934.
10. Attempts at performing the cross-metathesis of **5** with the methyl ether of **3** or the oxazole analog of **3** afforded complex mixtures.
11. Trnka TM, Grubbs RH. Acc Chem Res. 2001; 34:18. [PubMed: 11170353]
12. Stronger bases led to a rapid β -elimination of the C6 methyl ether.
13. Mahato TK, Babu S, Basak A. Biotech Lett. 1993; 15:1147. Alternative conditions, including LiOH, KOH, TMSOK, (Bu₃Sn)₂O, and Me₃SnOH, led to significant β -elimination at C6.
14. Phillips AJ, Uto Y, Wipf P, Reno MJ, Williams DR. Org Lett. 2000; 2:1165. [PubMed: 10804580]
15. Kiyooka S, Kaneko Y, Komura M, Matsuo H, Nakano M. J Org Chem. 1991; 56:2276.
16. Use of the more electron rich 3,4-DMB ether suppressed overoxidation of the allylic alcohol at C16 during DDQ deprotection.
17. Lu Z, Ma S. Angew Chem, Int Ed. 2008; 47:258.
18. (a) Hafner A, Duthaler RO, Marti R, Rihs G, Rothe-Streit P, Schwarzenbach F. J Am Chem Soc. 1992; 114:2321.(b) Cossy J, Willis C, Bellosta V, BouzBouz S. J Org Chem. 2002; 67:1982. [PubMed: 11925201]
19. Jarosz S, Gajewska A, Luboradzki R. Tetrahedron: Asymmetry. 2008; 19:1385.
20. NOE analysis confirmed the *cis*-cyclopropane stereochemistry.
21. Varying mixtures of the *cyclo*-monomer **21** and its corresponding C16 TBS-protected derivative were isolated and their structures confirmed by MALDI-TOF MS. Even upon attempted stepwise coupling of the two monomers, **21** was formed by an apparent *in situ* deprotection of the TES group of the acid component.




22. (a) Shiina I, Kubota M, Oshiumi H, Hashizume M. *J Org Chem.* 2004; 69:1822. [PubMed: 15058924] (b) Schweitzer D, Kane JJ, Strand D, McHenry P, Tenniswood M, Helquist P. *Org Lett.* 2007; 9:4619. [PubMed: 17915890]
23. Disorazole C₁ was not available for a direct comparison in these preliminary assays. However, reference 3c reports the IC₅₀ of disorazole C₁ and vincristine in HCT116 cells as 1.09±0.41 nM and 5.62±0.33 nM, respectively. Not surprisingly, there is some variability in the common standard vincristine in these two assays. However, in both batches of HCT116 cells, disorazole C₁ and **1** are more potent than vincristine by a factor of 5 and 1.2, respectively, making disorazole C₁ ca. 4-fold more active than its cyclopropane analog **1**.
24. For cyclopropane analogs of natural products, see, for example: (a) Cachoux F, Isarno T, Wartmann M, Altmann KH. *Synlett.* 2006:1384.(b) Wipf P, Reeves JT, Balachandran R, Day BW. *J Med Chem.* 2002; 45:1901. [PubMed: 11960501]




Scheme 1. Retrosynthetic approach for CP₂-disorazole C₁ 1




Scheme 2. Synthesis of ester 3



Scheme 3. Cross-metathesis and synthesis of aldehyde 10



Scheme 4. Synthesis of hydroxy ester segment 18



Scheme 5. Segment condensation and Shiina macrolactonization

Table 1 1

Cytotoxic activity of **1** and cyclic monomer **21** in human colon cancer cell lines. Vincristine is used as a reference compound.²³

cell line	1 IC ₅₀ [nM] ^b	21 IC ₅₀ [μM] ^b	vincristine IC ₅₀ [nM] ^b
RKO	28.0 ± 9.2	>50	18.6 ± 7.6
HCT116	28.3 ± 11.6	>50	35.2 ± 11.9
H630	49.5 ± 25.0	>50	68.0 ± 16.3

^a Cell proliferation and viability was quantified by the WST-1 assay (Roche Applied Science; Indianapolis, IN).

^b Values represent the mean±S.D. from 4-5 independent determinations.