

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51213155>

In Vitro and In Vivo Osteogenic Activity of Largazole

ARTICLE *in* ACS MEDICINAL CHEMISTRY LETTERS · MARCH 2011

Impact Factor: 3.12 · DOI: 10.1021/ml1002794 · Source: PubMed

CITATIONS

13

READS

24

9 AUTHORS, INCLUDING:



Hyung-Keun You

Wonkwang University

63 PUBLICATIONS 178 CITATIONS

SEE PROFILE



Seong Hwan Kim

Korea Research Institute of Chemical Techno...

119 PUBLICATIONS 731 CITATIONS

SEE PROFILE

Published in final edited form as:

ACS Med Chem Lett. 2011 March 10; 2(3): 248–251. doi:10.1021/ml1002794.

In Vitro and In Vivo Osteogenic Activity of Largazole

Su-Ui Lee[†], Han Bok Kwak[‡], Sung-Hee Pi[§], Hyung-Keun You[§], Seong Rim Byeon[#], Yongcheng Ying[#], Hendrik Luesch[¶], Jiyong Hong^{#,*}, and Seong Hwan Kim^{†,*}

[†] Laboratory of Chemical Genomics, Korea Research Institute of Chemical Technology, Daejeon 305-600, Korea

[‡] Department of Anatomy, School of Medicine, Wonkwang University, Iksan, Chonbuk 570-749, Korea

[§] Department of Periodontology, School of Dentistry, Wonkwang University, Iksan, Chonbuk 570-749, Korea

[#] Department of Chemistry, Duke University, Durham, NC 27708

[¶] Department of Medicinal Chemistry, University of Florida, Gainesville, FL 32610

Abstract

Due to their capability of modifying chromatin structure and thereby regulating gene transcription, histone deacetylases (HDACs) have been reported to play important roles in osteogenesis and considered a promising potential therapeutic target for bone diseases, including osteoporosis. We showed that the novel marine-derived HDAC inhibitor largazole exhibits *in vitro* and *in vivo* osteogenic activity. Largazole significantly induced the expression of ALP and OPN. The osteogenic activity of largazole was mediated through the increased expression of Runx2 and BMPs. Importantly, largazole showed *in vivo* bone-forming efficacy in the mouse calvarial bone formation assay and the rabbit calvarial bone fracture healing model. The dual action of largazole to stimulate bone formation and inhibit bone resorption would be a useful feature in drug development for bone-related disorders.

Keywords

largazole; osteogenic activity; histone deacetylases; Runx2; bone morphogenetic protein

The cyclic depsipeptide largazole (Figure 1A), isolated from a cyanobacterium of the genus *Symploca* by Luesch and co-workers, is a marine natural product with novel chemical scaffold.¹ Largazole possesses highly differential growth-inhibitory activity, preferentially targeting transformed over non-transformed cells. The intriguing structure and biological activity of largazole have attracted strong interest from the synthetic chemistry community to establish synthetic routes to largazole and to investigate its potential as a cancer therapeutic. Luesch and Hong reported the first total synthesis of largazole, identified histone deacetylases (HDACs) as the molecular targets, and demonstrated antitumor activity in a xenograft mouse model.^{2,3} Following our initial report, additional total and formal syntheses, further studies on its HDAC inhibition, and structure–activity relationship studies have appeared in the literature.^{4–17}

Corresponding Authors: hwan@krikt.re.kr; jiyong.hong@duke.edu.

Supporting Information Available. General experimental procedures. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Osteogenesis is a complex process associated with dramatic changes in gene expression.¹⁸ Due to their capability of modifying chromatin structure and thereby modulating gene transcription, HDACs have been reported to play important roles in osteogenesis and considered a promising potential therapeutic target for bone diseases, including osteoporosis.¹⁹ Several HDAC inhibitors, including trichostatin A,²⁰ sodium butyrate,²⁰ and valproic acid²¹ have been tested for their osteogenic activity, but none have been shown effective in *in vivo* models for further development as potential therapeutics for bone-related diseases. Moreover, it has been reported that long-term treatment with valproic acid increases the incidence of osteoporosis and bone fracture.^{22,23} Herein, we report the *in vitro* and *in vivo* osteogenic activity and the significant potential of largazole in bone formation, repair and regeneration.

Murine pluripotent mesenchymal precursor C2C12 cells were used as an *in vitro* model of osteogenesis. The C2C12 cells have been known to be committed into osteoblasts with the induction of several osteogenic bone morphogenetic protein (BMP) isoforms such as BMP-2, 4, 6, 7, and 9.²⁴ Before evaluating the osteogenic activity of largazole, we examined the effect of largazole on viability of C2C12 cells and found that largazole did not exhibit a significant level of cytotoxicity up to 100 nM (see the Supporting Information for details).

Next, we examined the effects of largazole on the inhibition of HDAC activity and the accumulation of acetylated histones in C2C12 cells to confirm its HDAC inhibitory activity. Largazole decreased the total HDAC activity in C2C12 cells (Figure 1B)²⁵ and significantly increased the levels of acetylated histones in a dose-dependent manner (Figure 1C).

The effect of largazole on the commitment of C2C12 cells into osteoblasts was determined by the expression level of alkaline phosphatase (ALP), an early phase marker of osteoblast differentiation. Up to 50 nM, largazole increased the expression of ALP in a dose-dependent manner (Figure 1D). In addition, largazole significantly induced the expression of ALP and osteopontin (OPN) at the transcript level (see the Supporting Information for details).

Since Runx2 has been shown to be critical in osteogenesis and regulates the expression of bone-specific genes such as ALP and OPN,²⁶ we examined the effect of largazole on the activation and mRNA expression of Runx2 using the p6×osteoblast-specific *cis*-acting element 2-luciferase reporter²⁷ and real-time PCR, respectively (Figure 2). Relative to the DMSO control, largazole at 50 nM significantly increased the activity and mRNA level of Runx2 by 29-fold and 3-fold, respectively (Figure 2A and 2B). Since HDACs have been identified as co-repressor proteins to affect Runx2 activity,²⁸ these results suggested that inhibition of HDACs by largazole increases Runx2 activation, potentiates osteoblast differentiation, and increases bone formation.

From the data showing that the increased expression of ALP induced by largazole was inhibited by noggin, a specific inhibitor of BMPs (see Figure 1D), we hypothesized that the osteogenic activity of largazole could result from its potential to increase the expression of BMPs. To test this hypothesis, we evaluated the effect of largazole on the mRNA expression levels of BMPs (Figure 2B). Largazole significantly induced the expressions of BMP-2, 4, 6, 7, and 9, supporting our hypothesis that the osteogenic activity of largazole is mediated through the increase of the expression and/or activity of Runx2 and BMPs. Recently, BMP-2 and BMP-7 with strong osteogenic activity have been approved for clinical applications including spinal fusion, fracture healing, and dental tissue engineering.^{29,30} Therefore, anabolic agents such as largazole that stimulate BMP expression or the BMP-signaling pathway would be useful for the treatment of osteoblast-related diseases by bone formation or regeneration.^{31,32} In addition, we found that largazole inhibits the formation of multinucleated osteoclasts (see the Supporting Information for details), suggesting that

largazole might elicit the dual action to stimulate bone formation and suppress bone resorption.

Next, we examined *in vivo* osteogenic activity of largazole using two independent *in vivo* models, the mouse calvarial bone formation assay and the rabbit calvarial bone fracture healing model.

In the mouse calvarial bone formation assay, when collagen sponges soaked with largazole were implanted in calvarial bones, largazole induced woven bone formation over the periosteum of the calvarial bones (Figure 3). The woven bone formation at the lower concentration (10 μ M, Figure 3B) was more significant than that at the higher concentration (50 μ M, Figure 3C). This data suggested that largazole may show the biphasic effect which has been observed with other osteogenic compounds.^{33–35} Further studies are required to determine the optimal dose for the highest *in vivo* efficacy.

We further evaluated *in vivo* bone-forming activity of largazole in the rabbit calvarial bone fracture healing model (Figure 4). In this study, macroporous biphasic calcium phosphate (MBCP) was used as a basic scaffold to induce bone formation. While an incomplete bone formation was observed with MBCPs alone (Figure 4A), newly-formed bones in direct contact with MBCPs mixed with largazole were observed (Figure 4B and 4C). Since there is a significantly unmet clinical need for small molecules to improve the biological properties of the scaffold and its osteoconductivity, this data clearly shows the great potential of largazole for improvement of the property of bone graft substitutes in bone defect reconstruction.

In summary, we showed that the HDAC inhibitor largazole exhibits *in vitro* and *in vivo* osteogenic activity. Largazole significantly induced the expression of ALP and OPN. The osteogenic activity of largazole was mediated through the increased expression of Runx2 and BMPs. More importantly, largazole showed *in vivo* bone-forming efficacy in the mouse calvarial bone formation assay and the rabbit calvarial bone fracture healing model. The dual action of largazole to stimulate bone formation and inhibit bone resorption would be a useful feature in drug development for bone-related disorders. Taken together, largazole shows the great clinical potential as a novel treatment for bone-related diseases in addition to its already proven potential as an antitumor agent.³

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Professors Hyun-Mo Ryoo and Zang Hee Lee (Department of Cell and Developmental Biology, School of Dentistry and Dental Research Institute, Seoul National University, Seoul, Korea) for the p6 \times OSE2-luc reporter vector and helpful discussion.

Funding Sources: This work has been supported by grants from the Korea Healthcare Technology R&D Project, the Ministry of Health, Welfare & Family Affairs, Republic of Korea (A084242 to S.L. and S.H.K.) and the National Institutes of Health (R01CA138544-01 to J.H. and H.L.).

References

1. Taori K, Paul VJ, Luesch H. Structure and Activity of Largazole, a Potent Antiproliferative Agent from the Floridian Marine Cyanobacterium *Symploca* sp. J Am Chem Soc. 2008; 130:1806–1807. [PubMed: 18205365]

2. Ying Y, Taori K, Kim H, Hong J, Luesch H. Total Synthesis and Molecular Target of Largazole, a Histone Deacetylase Inhibitor. *J Am Chem Soc.* 2008; 130:8455–8459. [PubMed: 18507379]
3. Liu Y, Salvador LA, Byeon S, Ying Y, Kwan JC, Law BK, Hong J, Luesch H. Anticancer Activity of Largazole, a Marine-Derived Tunable Histone Deacetylase Inhibitor. *J Pharmacol Exp Ther.* 2010; 335:351–361. [PubMed: 20739454]
4. Bowers A, West N, Taunton J, Schreiber SL, Bradner JE, Williams RM. Total Synthesis and Biological Mode of Action of Largazole: A Potent Class I Histone Deacetylase Inhibitor. *J Am Chem Soc.* 2008; 130:11219–11222. [PubMed: 18642817]
5. Seiser T, Kamena F, Cramer N. Synthesis and Biological Activity of Largazole and Derivatives. *Angew Chem Int Ed.* 2008; 47:6483–6485.
6. Nasveschuk CG, Ungermannova D, Liu X, Phillips AJ. A Concise Total Synthesis of Largazole, Solution Structure, and Some Preliminary Structure Activity Relationships. *Org Lett.* 2008; 10:3595–3598. [PubMed: 18616341]
7. Ghosh AK, Kulkarni S. Enantioselective Total Synthesis of (+)-Largazole, a Potent Inhibitor of Histone Deacetylase. *Org Lett.* 2008; 10:3907–3909. [PubMed: 18662003]
8. Ying Y, Liu Y, Byeon SR, Kim H, Luesch H, Hong J. Synthesis and Activity of Largazole Analogues with Linker and Macrocyclic Modification. *Org Lett.* 2008; 10:4021–4024. [PubMed: 18707106]
9. Ren Q, Dai L, Zhang H, Tan W, Xu Z, Ye T. Total Synthesis of Largazole. *Synlett.* 2008; 2379–2383.
10. Numajiri Y, Takahashi T, Takagi M, Shin-ya K, Doi T. Total Synthesis of Largazole and Its Biological Evaluation. *Synlett.* 2008; 2483–2486.
11. Bowers AA, Greshock TJ, West N, Estiu G, Schreiber SL, Wiest O, Williams RM, Bradner JE. Synthesis and Conformation–Activity Relationships of the Peptide Isosteres of FK228 and Largazole. *J Am Chem Soc.* 2009; 131:2900–2905. [PubMed: 19193120]
12. Bowers AA, West N, Newkirk TL, Troutman-Youngman AE, Schreiber SL, Wiest O, Bradner JE, Williams RM. Synthesis and Histone Deacetylase Inhibitory Activity of Largazole Analogs: Alteration of the Zinc-Binding Domain and Macrocyclic Scaffold. *Org Lett.* 2009; 11:1301–1304. [PubMed: 19239241]
13. Chen F, Gao AH, Li J, Nan FJ. Synthesis and Biological Evaluation of C7-Demethyl Largazole Analogues. *Chem Med Chem.* 2009; 4:1269–1272. [PubMed: 19431162]
14. Wang B, Forsyth CJ. Total Synthesis of Largazole - Devolution of a Novel Synthetic Strategy. *Synthesis.* 2009; 2873–2880.
15. Souto JA, Vaz E, Lepore I, Pöppler AC, Franci G, Álvarez R, Altucci L, de Lera ÁR. Synthesis and Biological Characterization of the Histone Deacetylase Inhibitor Largazole and C7- Modified Analogues. *J Med Chem.* 2010; 53:4654–4667. [PubMed: 20491440]
16. Zeng X, Yin B, Hu Z, Liao C, Liu J, Li S, Li Z, Nicklaus MC, Zhou G, Jiang S. Total Synthesis and Biological Evaluation of Largazole and Derivatives with Promising Selectivity for Cancers Cells. *Org Lett.* 2010; 12:1368–1371. [PubMed: 20184338]
17. Xiao Q, Wang LP, Jiao XZ, Liu XY, Wu Q, Xie P. Concise Total Synthesis of Largazole. *Journal of Asian Natural Products Research.* 2010; 12:940–949. [PubMed: 21061215]
18. Katagiri T, Takahashi N. Regulatory Mechanisms of Osteoblast and Osteoclast Differentiation. *Oral Dis.* 2002; 8:147–159. [PubMed: 12108759]
19. Jensen ED, Nair AK, Westendorf JJ. Histone Deacetylase Co-Repressor Complex Control of Runx2 and Bone Formation. *Crit Rev Eukaryot Gene Expr.* 2007; 17:187–196. [PubMed: 17725488]
20. Schroeder TM, Westendorf JJ. Histone Deacetylase Inhibitors Promote Osteoblast Maturation. *J Bone Miner Res.* 2005; 20:2254–2263. [PubMed: 16294278]
21. Cho HH, Park HT, Kim YJ, Bae YC, Suh KT, Jung JS. Induction of Osteogenic Differentiation of Human Mesenchymal Stem Cells by Histone Deacetylase Inhibitors. *J Cell Biochem.* 2005; 96:533–542. [PubMed: 16088945]
22. Guo CY, Ronen GM, Atkinson SA. Long-Term Valproate and Lamotrigine Treatment May Be a Marker for Reduced Growth and Bone Mass in Children with Epilepsy. *Epilepsia.* 2001; 42:1141–1147. [PubMed: 11580761]

23. Boluk A, Guzelipek M, Savli H, Temel I, Ozişik HI, Kaygusuz A. The Effect of Valproate on Bone Mineral Density in Adult Epileptic Patients. *Pharmacol Res.* 2004; 50:93–97. [PubMed: 15082034]
24. Cheng H, Jiang W, Phillips FM, Haydon RC, Peng Y, Zhou L, Luu HH, An N, Breyer B, Vanichakarn P, Szatkowski JP, Park JY, He TC. Osteogenic Activity of the 14 Types of Human Bone Morphogenetic Proteins (BMPs). *J Bone Joint Surg Am.* 2003; 85A:1544–1552. [PubMed: 12925636]
25. Largazole reduced the expressions of HDACs 2, 3, 4, and 5, but not the expression of HDAC 1 (see the Supporting Information for details).
26. Franceschi RT, Xiao G. Regulation of the Osteoblast-Specific Transcription Factor, Runx2: Responsiveness to Multiple Signal Transduction Pathways. *J Cell Biochem.* 2003; 88:446–454. [PubMed: 12532321]
27. Kim HJ, Park HD, Kim JH, Cho JY, Choi JY, Kim JK, Kim HJ, Shin HI, Ryoo HM. Establishment and Characterization of a Stable Cell Line to Evaluate Cellular Runx2 Activity. *J Cell Biochem.* 2004; 9:1239–1247. [PubMed: 15048877]
28. Westendorf JJ. Transcriptional Co-Repressors of Runx2. *J Cell Biochem.* 2006; 98:54–64. [PubMed: 16440320]
29. McKay WF, Peckham SM, Badura JM. A Comprehensive Clinical Review of Recombinant Human Bone Morphogenetic Protein-2 (INFUSE Bone Graft). *Int Orthop.* 2007; 31:729–734. [PubMed: 17639384]
30. White AP, Vaccaro AR, Hall JA, Whang PG, Friel BC, McKee MD. Clinical Applications of BMP-7/OP-1 in Fractures, Nonunions and Spinal Fusion. *Int Orthop.* 2007; 31:735–741. [PubMed: 17962946]
31. Nakashima M, Reddi AH. The Application of Bone Morphogenetic Proteins to Dental Tissue Engineering. *Nat Biotechnol.* 2003; 21:1025–1032. [PubMed: 12949568]
32. Seeherman H, Wozney JM. Delivery of Bone Morphogenetic Proteins for Orthopedic Tissue Regeneration. *Cytokine Growth Factor Rev.* 2005; 16:329–345. [PubMed: 15936978]
33. Yeo H, Beck LH, McDonald JM, Zayzafoon M. Cyclosporin A Elicits Dose-Dependent Biphasic Effects on Osteoblast Differentiation and Bone Formation. *Bone.* 2007; 40:1502–1516. [PubMed: 17392048]
34. Jeong JG, Kim YS, Min YK, Kim SH. Low Concentration of 3-Carene Stimulates the Differentiation of Mouse Osteoblastic MC3T3-E1 Subclone 4 Cells. *Phytother Res.* 2008; 22:18–22. [PubMed: 17685387]
35. Lee SU, Shin HK, Min YK, Kim SH. Emodin Accelerates Osteoblast Differentiation through Phosphatidylinositol 3-Kinase Activation and Bone Morphogenetic Protein-2 Gene Expression. *Int Immunopharmacol.* 2008; 8:741–747. [PubMed: 18387517]




Figure 1.

Largazole exhibited HDAC inhibitory and osteogenic activities in C2C12 cells. (A) Structure of largazole. (B) Effect of largazole on the total HDAC activity in C2C12 cells. Cells (4×10^3 cells/well) were cultured in a 96-well plate for 24 h and pre-treated with largazole for 1 h before incubation with deacetylase Lysyl substrate of the HDAC Fluorescent Activity Assay kit. HDAC activity was measured according to the manufacturer's protocol. (C) Effect of largazole on the level of hyperacetylated histones in C2C12 cells. Cells (2×10^6 cells/plate) were cultured in a 100-mm plate for 24 h and treated with largazole. After 24 h, protein extracts were prepared and western blot analysis was performed. Actin was used as a loading control. (D) Effect of largazole on osteoblast differentiation in C2C12 cells. The effect of largazole on osteoblast differentiation was evaluated by ALP staining. Cells (4×10^3 cells/well) were cultured in a 96-well plate for 24 h and treated with largazole for 6 days. Medium was changed every 3 days. Noggin (a specific BMP inhibitor) was co-treated with largazole.




Figure 2.

Effect of largazole on the Runx2 activation and mRNA induction of Runx2 and BMPs in C2C12 cells. (A) Runx2 activation was measured by luciferase reporter assay using 6×OSE2-luc plasmid-transfected C2C12 cells. The cells (4×10^3 cells/well) were cultured in a 96-well plate for 1 day and incubated with DMEM containing 5% FBS in the presence or absence of largazole for 1 day before luciferase activity assay. (B) The mRNA levels were evaluated by quantitative real-time PCR in cells (1×10^6 cells/60-mm plate) treated with largazole for 6 days. rhBMP-2 (300 ng/ml) was used as the reference of osteoblastogenesis inducer (^a $P < 0.05$; ^b $P < 0.01$; ^c $P < 0.001$).




Figure 3.

Largazole exhibited bone-forming activity in mouse calvaria. Collagen sponges containing 5 μ l of PBS vehicle (A) or largazole (B, 10 μ M; C, 50 μ M) were placed onto mouse calvarial bones. After a 3- week implantation, the mice were sacrificed. Calvarial bones were removed, fixed, decalcified, embedded in paraffin, and sectioned. Sections were stained with H&E and photographed at 200 \times magnification. Arrows indicated the region of newly-forming woven bone.




Figure 4.

Largazole exhibited bone-regenerating activity in rabbit calvaria. Bone graft substitute (10 mg), MBCP (indicated by a black asterisk (*); Biomatlante, France) was mixed with 50 μ l of PBS (A) or largazole (B, 100 nM; C, 250 nM), and defects were filled with these mixtures. Rabbits were sacrificed at 4 weeks after surgery. The bone (indicated by 'white +') grows into and around bone graft substitute. Box images in the top were magnified in the bottom.