

Erratum: Infrared Detection of the Formyl Radical HCO

George E. Ewing, Warren E. Thompson, and George C. Pimentel

Citation: *The Journal of Chemical Physics* **34**, 1067 (1961); doi: 10.1063/1.1731644

View online: <http://dx.doi.org/10.1063/1.1731644>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/34/3?ver=pdfcov>

Published by the AIP Publishing

Articles you may be interested in

[Vibronic structure in triatomic molecules: The hydrocarbon flame bands of the formyl radical \(HCO\). A theoretical study](#)

J. Chem. Phys. **108**, 7202 (1998); 10.1063/1.476138

[Photofragment excitation spectroscopy of the formyl \(HCO/DCO\) radical: Linewidths and predissociation rates of the \$\tilde{A}\(A'\)\$ state](#)

J. Chem. Phys. **94**, 1796 (1991); 10.1063/1.459953

[Two-photon resonance REMPI detection of the formyl radical](#)

J. Chem. Phys. **84**, 5334 (1986); 10.1063/1.449944

[The electronic structure of the formyl radical HCO](#)

J. Chem. Phys. **71**, 3697 (1979); 10.1063/1.438824

[Infrared Detection of the Formyl Radical HCO](#)

J. Chem. Phys. **32**, 927 (1960); 10.1063/1.1730819

A promotional banner for AIP Applied Physics Reviews. The background is a blue gradient with a bright light source on the right and several blue spheres of varying sizes. On the left, there is a small image of a book cover for 'AIP Applied Physics Reviews' showing a technical diagram. The text 'NEW Special Topic Sections' is prominently displayed in white. Below this, in orange, it says 'NOW ONLINE' followed by 'Lithium Niobate Properties and Applications: Reviews of Emerging Trends'. The AIP logo and 'Applied Physics Reviews' are in the bottom right corner.

NEW Special Topic Sections

NOW ONLINE
Lithium Niobate Properties and Applications:
Reviews of Emerging Trends

AIP Applied Physics Reviews

Erratum: Distribution Functions of a Fluid in an External Potential Field. Application to Physical Adsorption

[J. Chem. Phys. **33**, 464 (1960)]

WILLIAM A. STEELE AND MARVIN ROSS

Department of Chemistry, The Pennsylvania State University, University Park, Pennsylvania

(Received January 4, 1961)

A FACTOR of $\frac{1}{2}$ was omitted from one of the terms in Eq. (5.2). This equation should read

$$A_c - A_0 = \omega_1 - \omega_2 / 2kT + 0 / (kT)^2. \quad (5.2)$$

This error was carried over into several following equations. In particular, the last terms on the right-hand sides of Eqs. (5.9), (6.9), and (6.10) should be multiplied by $\frac{1}{2}$, and Eq. (6.11) should be altered to read

$$\mu_{ac} - \mu_{lc} = \bar{u}_s(L_z) [1 + \rho_0 k_0 \bar{u}_s(L_z) / 2 + \dots]. \quad (6.11)$$

This error thus invalidates the conclusion that the effect of the first-order change in the density of an adsorbed film in a weak external field upon the chemical potential is cancelled by the second-order change in the configurational free energy.

The third term in Eq. (14) should have a prime on the subscript of (V) ; it should read

$$(V)_{ia; ja'}^2. \quad (14)$$

The first sum in Eq. (20) is over capital K ; the second to the last term in Eq. (20) should have

$$\mathbf{q}_{i0a'} \text{ replace } \mathbf{q}_{j0a'}.$$

Equation (24) should read

$$\begin{aligned} \sum_K \nu_{OAK} \mathbf{u}_{OAK} \cdot \mathbf{u}_{AOK} &= \nu_{OA} \mathbf{u}_{OA} \cdot \mathbf{u}_{AO} = N \mu_{0a} \cdot \mu_{a0} \nu_{0a} \\ &- \sum_{i=1}^N \sum_{j \neq i} \sum_{a' \neq a} \frac{4(V)_{ia; ja'} \mathbf{u}_{i0a} \cdot \mathbf{u}_{j0a'} \nu_{j0a'} \nu_{0a}}{h(\nu_{j0a'}^2 - \nu_{0a}^2)} \\ &- 2 \sum_{i=1}^N \sum_{j \neq i} (V)_{ia; j0} \mathbf{u}_{i0a} \cdot (\mathbf{u}_{jaa} - \mathbf{u}_{i0}) / h \\ &- 2 \sum_{i=1}^N \sum_{j \neq i} \sum_{a' \neq a} \frac{(V)_{iaa'; j0} \mathbf{u}_{i0a} \cdot \mathbf{u}_{j0a'} \nu_{0a}}{h(\nu_{0a'} - \nu_{0a})} \\ &+ \sum_{i=1}^N \sum_{j \neq i} [(V)_{iaa; j0} - (V)_{i00; j0}] \mathbf{u}_{i0a} \cdot \mathbf{u}_{j0a} / h. \quad (24) \end{aligned}$$

Erratum: Optical and Other Electronic Properties of Polymers

[J. Chem. Phys. **33**, 1332 (1960)]

IGNACIO TINOCO, JR.

Chemistry Department, University of California, Berkeley, California

(Received January 16, 1961)

PROFESSOR R. Feynman, California Institute of Technology, has pointed out that Eq. (24) of this paper cannot be substituted into Eqs. (22) or (23) to give the correct oscillator strength or polarizability for a polymer. Instead of an expression for $\mathbf{u}_{0A} \cdot \mathbf{u}_{A0}$ we need one for $\nu_{0A} \mathbf{u}_{0A} \cdot \mathbf{u}_{A0}$. The new equation is given below, together with some corrected typographical errors. An analogous correction in a previous paper (J. Am. Chem. Soc. **1960**, 4785) leads to the deletion of the second term in Eqs. (1) and Eq. (A11) of that paper.

The last term in Eq. (4) should be

$$\dots \sum_{j \neq i} [(V)_{ia; j0} \psi_0^0 / h \nu_{0a}]. \quad (4)$$

Equation (5) should read

$$(V)_{ia; ja'} = \int \psi_{ia}^0 V_{a i, t j} \psi_{ja'}^0 d\tau. \quad (5)$$

Erratum: Infrared Detection of the Formyl Radical HCO

[J. Chem. Phys. **32**, 927 (1960)]GEORGE E. EWING, WARREN E. THOMPSON, AND
GEORGE C. PIMENTEL*Department of Chemistry, University of California, Berkeley, California*

(Received December 19, 1960)

AN error in our calculations of the thermodynamic functions of HCO has been brought to our attention.¹ We have omitted the statistical weight factor associated with the doublet character of the ground electronic state of this free radical. A temperature independent term of $R \ln 2 = +1.38$ cal/deg mole must be added to every number in the columns headed $-(F^\circ - H_0^\circ)/T$ and S° in Table IX. The enthalpy function and heat capacity computations are unchanged.

¹ Dr. J. Gordon, private communication.