

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/13640820>

# Magnone A and B, Novel Anti-PAF Tetrahydrofuran Lignans from the Flower Buds of *Magnolia fargesii*

ARTICLE *in* JOURNAL OF NATURAL PRODUCTS · JULY 1998

Impact Factor: 3.8 · DOI: 10.1021/np970445+ · Source: PubMed

CITATIONS

41

READS

34

8 AUTHORS, INCLUDING:



**Sei-Ryang Oh**

Korea Research Institute of Bioscience and ...

180 PUBLICATIONS 1,899 CITATIONS

SEE PROFILE



**Si Hyung Park**

Mokpo National University

55 PUBLICATIONS 614 CITATIONS

SEE PROFILE



**Hwa-Young Lee**

Soon Chun Hyang University Hospital

328 PUBLICATIONS 5,619 CITATIONS

SEE PROFILE



## 5 $\alpha$ ,7 $\alpha$ (H)-6,8-CYCLOEUDESMA-1 $\beta$ ,4 $\beta$ -DIOL FROM THE FLOWER BUDS OF *MAGNOLIA FARGESII*

KEUN YOUNG JUNG, DONG SEON KIM, SI-HYUNG PARK, IM SEON LEE, SEI RYANG OH, JUNG JOON LEE, EUN-HEE KIM†, CHAEJOON CHEONG† and HYEONG-KYU LEE\*

Natural Product Biosynthesis Research Unit, Korea Research Institute of Bioscience and Biotechnology, Yusong P.O. Box 115, Taejeon 305-600; † Magnetic Resonance Group, Korea Basic Science Institute, Taejeon 305-333, Korea

(Received in revised form 22 October 1997)

**Key Word Index**—*Magnolia fargesii*; Magnoliaceae; cycloedesmane-type sesquiterpene; 5 $\alpha$ ,7 $\alpha$ (H)-6,8-cycloedesma-1 $\beta$ ,4 $\beta$ -diol.

**Abstract**—From the Chinese crude drug *shin-i*, the flower buds of *Magnolia fargesii*, a new cycloedesmane-type sesquiterpene, 5 $\alpha$ ,7 $\alpha$ (H)-6,8-cycloedesma-1 $\beta$ ,4 $\beta$ -diol, was isolated. The structure was elucidated by means of various NMR techniques. This compound has biogenetic significance because it is a plausible intermediate of the oppositol-type compound such as homalomenol A reported from this plant. © 1998 Elsevier Science Ltd. All rights reserved


### INTRODUCTION

The dried flower buds of *Magnolia fargesii* has a history of use for the treatment of nasal congestion with headache, sinusitis, and allergic rhinitis [1]. Pharmacological studies have revealed that *shin-i* has uterus-stimulating, hypotensive, antifungal, and skeletal muscle contracting effects [2, 3]. In previous reports on chemical investigations of this plant, many kinds of essential oils, lignans, neolignans and sesquiterpenes have been found and pharmacological activities of these lignans from *shin-i*, Ca<sup>2+</sup> antagonistic activity and platelet activating factor (PAF) antagonistic activity were revealed [4–10].

Recently, we reported the structural elucidation of the four sesquiterpenes, oplopanone, oplodiol, homalomenol A and 1 $\beta$ ,4 $\beta$ ,7 $\alpha$ -trihydroxyeudesmane isolated from the ethyl acetate extracts of this plant besides seven known phenolic lignans with PAF antagonistic activity [11]. In a continuing study, we have isolated a new cycloedesmane-type sesquiterpene, 5 $\alpha$ ,7 $\alpha$ (H)-6,8-cycloedesma-1 $\beta$ ,4 $\beta$ -diol (**1**) from this plant.

### RESULTS AND DISCUSSION

An ethyl acetate extract of the flower buds of *M. fargesii* was fractionated by repeated column chro-



**1**

matography (silica gel and RP-18) to yield compound **1** as colorless needles.

Compound **1** revealed an absorption at 3401 cm<sup>-1</sup> in its IR spectrum suggesting the presence of a hydroxyl group. The EI mass spectrum of **1** exhibited a molecular ion peak at *m/z* 238 and the significant fragmentation ion peaks at *m/z* 220 [M–H<sub>2</sub>O]<sup>+</sup>, 202 [M–2H<sub>2</sub>O]<sup>+</sup>, 195 [M–C<sub>3</sub>H<sub>7</sub>]<sup>+</sup>, 177 [220–C<sub>3</sub>H<sub>7</sub>]<sup>+</sup> and/or [195–H<sub>2</sub>O]<sup>+</sup>, indicating the presence of at least two hydroxyl groups and a propyl group.

The <sup>1</sup>H NMR spectrum of **1** in CDCl<sub>3</sub> showed a methine proton attached a hydroxyl function at  $\delta$  3.33 (1H, *dd*, *J* = 10.9 and 4.4 Hz), two tertiary methyl proton at  $\delta$  1.30 (3H, *s*) and 1.17 (angular Me, *s*), and an isopropyl group at  $\delta$  0.93 (3H, *d*, *J* = 6.6 Hz), 0.97 (3H, *d*, *J* = 6.6 Hz) and 0.98 (1H, *m*), which was further confirmed by <sup>1</sup>H–<sup>1</sup>H COSY and HMBC spectra. The <sup>13</sup>C NMR and DEPT spectra suggested that the skeleton consisted of 15 carbons: four methyls, three methylenes, six methines and two quaternary carbons. From the HMQC spectrum of **1**, the chemi-

\*Author to whom correspondence should be addressed. Tel. +82-42-860-4413; Fax. +82-42-860-4608; e-mail: hykylee@kribb4680.kribb.re.kr.

Table 1.  $^1\text{H}$  and  $^{13}\text{C}$  NMR Assignments for  $5\alpha,7\alpha(\text{H})$ -6,8-cycloeudesma-1 $\beta$ ,4 $\beta$ -diol (**1**)

Position	$^1\text{H}$		$^{13}\text{C}$	
	$\delta_{\text{H}}^{\text{a}}$	$\delta_{\text{H}}^{\text{b}}$	$\delta_{\text{C}}^{\text{a}}$	$\delta_{\text{C}}^{\text{b}}$
1	3.33 (1H, <i>dd</i> , 10.9, 4.4)	3.27 (1H, <i>dd</i> , 11.4, 4.3)	78.04 ( <i>d</i> )	78.99
2	1.72 (1H, <i>m</i> )	1.76 (1H, <i>m</i> )	28.20 ( <i>t</i> )	28.89
	1.61 (1H, <i>m</i> )	1.49 (1H, <i>m</i> )		
3	1.66 (1H, <i>m</i> )	1.64 (1H, <i>m</i> )	39.57 ( <i>t</i> )	40.42
	1.35 (1H, <i>m</i> )	1.38 (1H, <i>m</i> )		
4			70.98 ( <i>s</i> )	71.52
5	0.71 (1H, <i>d</i> , 4.6)	0.69 (1H, <i>d</i> , 5.7)	60.61 ( <i>d</i> )	62.15
6	1.17 (1H, <i>m</i> )	1.32 (1H, <i>m</i> )	23.97 ( <i>d</i> )	25.37
7	0.52 (1H, <i>m</i> )	0.45 (1H, <i>ddd</i> , 8.8, 3.0, 3.0)	49.89 ( <i>d</i> )	51.12
8	1.18 (1H, <i>m</i> )	1.13 (1H, <i>m</i> )	24.37 ( <i>d</i> )	25.50
9	0.91 (1H, <i>m</i> )	0.83 (1H, <i>m</i> )	44.48 ( <i>t</i> )	45.81
	1.84 (1H, <i>m</i> )	1.82 (1H, <i>m</i> )		
10			58.03 ( <i>s</i> )	59.40
11	0.98 (1H, <i>m</i> )	0.88 (1H, <i>m</i> )	32.48 ( <i>d</i> )	33.94
12	0.93 (3H, <i>d</i> , 6.6)	0.96 (3H, <i>d</i> , 6.5)	21.87 ( <i>q</i> )	22.29
13	0.97 (3H, <i>d</i> , 6.6)	0.98 (3H, <i>d</i> , 6.5)	21.73 ( <i>q</i> )	22.29
14	1.17 (3H, <i>s</i> )	1.16 (3H, <i>d</i> , 0.9)	15.51 ( <i>q</i> )	16.23
15	1.30 (3H, <i>s</i> )	1.28 (3H, <i>s</i> )	30.28 ( <i>q</i> )	30.54

<sup>a</sup>Assignments were based on DEPT,  $^1\text{H}$ - $^1\text{H}$  COSY, HMQC and HMBC in  $\text{CDCl}_3$ .<sup>b</sup>Measured in  $\text{CD}_3\text{OD}$ .

cal shifts of protonated carbons were assigned as listed in Table 1. Characteristic  $^{13}\text{C}$  NMR signals of  $\delta$  78.04 (CH), 70.98 (C), 58.03 (C), 30.28 ( $\text{CH}_3$ ) and 15.51 ( $\text{CH}_3$ ) suggested a eudesmane skeleton with one secondary and one tertiary hydroxyl group. In addition, three methine carbon signals at  $\delta$  23.97, 24.37 and 49.89, which appeared in the significantly high field region, indicated the presence of a 1,2,3-trisubstituted cyclopropane ring [12]. In the HMBC correlations, two methyl proton signals of an isopropyl group were correlated to the carbon signals at 32.48 (C-11) and 49.89 (C-7), and the latter was correlated with the proton signals at  $\delta$  0.71 (H-5) and 1.84 (H-9). Also, in the cyclopropane ring moiety, the methine carbon signals at  $\delta$  23.97 (C-6) and 24.37 (C-8) were correlated with the proton signal of H-5 and H-9, respectively. The methyl proton signal at  $\delta$  1.17 (H-14) was long-range-correlated to the methine carbon signal at  $\delta$  60.61 (C-5) to which the proton signals at  $\delta$  3.33 (H-1) and 0.52 (H-7) were correlated. All the remaining proton signals were correlated to proximate carbon signals through two and/or three bond connections as shown Fig. 1. Thus, the structure of **1** could be assigned from the above evidence; its stereochemistry was determined as described below.

The configuration of the hydroxyl group at C-1 was confirmed to be in the equatorial position ( $\beta$ ) from the fact that the coupling constants (*dd*,  $J = 10.9$  and  $4.4$  Hz) of H-1 in the  $^1\text{H}$  NMR spectrum indicated its axial position [13]. Regarding the configuration of the hydroxyl at C-4, an axial orientation ( $\beta$ ) will result in deshielding of the methyl carbon (C-15) and a down-field shift in its resonance to approximately  $\delta$  30 [14,

15]. An equatorial orientation ( $\alpha$ ) results in the C-15 carbon having a chemical shift of approximately  $\delta$  25 [16, 17]. The corresponding signal observed in the  $^{13}\text{C}$  NMR spectrum is  $\delta$  30.28 and this value agrees well with the assignment of the axial orientation. The stereochemistry of C-7 was deduced through the observation of the NOESY spectrum of **1** in  $\text{CD}_3\text{OD}$  solution because the signals of H-6, H-8 and the methyl proton of C-14 in the  $^1\text{H}$  NMR spectrum in  $\text{CDCl}_3$  were very much overlapped. The NOESY spectrum of **1** showed strong interactions between the proton at  $\delta$  0.69 (H-5) and the proton at  $\delta$  0.45 (H-7), and no interaction of the proton at  $\delta$  1.16 (H-14) with H-7 and H-11. This result indicated that the configurations of the isopropyl moiety and the proton of C-7 are equatorial and axial orientations, respectively, as shown in the proposed structure (chair con-





Fig. 1. HMBC correlations for  $5\alpha,7\alpha(\text{H})$ -6,8-cycloeudesma-1 $\beta$ ,4 $\beta$ -diol (**1**). The arrow indicate correlations from proton to carbon.

Scheme 1. The possible biogenetic conversion of **1** into homalomenol A (**2**).

formation). All of these findings established that the structure of **1** was 5 $\alpha$ ,7 $\alpha$ (H)-6,8-cycloeudesma-1 $\beta$ ,4 $\beta$ -diol (**1**), a new cycloeudesmane-type sesquiterpene.

The reports of cycloeudesmane-type sesquiterpenes show that they are very rare in nature [18]. In the case of the 6,8-cycloeudesmane-type, Itokawa *et al.* [12] have reported the isolation of 1 $\beta$ -hydroxy-6,8-cyclo-4(15)eudsmene from the plant *Torilis japonica*. This compound was also obtained by the biomimetic reaction of epoxygermacrene-D with basic alumina [19]. Therefore, compound **1** has a biogenetic significance because it is a plausible intermediate of the oppositol-type of compound such as homalomenol A (**2**) [11] isolated from this plant (Scheme 1).

#### EXPERIMENTAL

Mps: Electrothermal 9100, uncorr. Optical rotation was recorded on Jasco DIP-370 digital polarimeter. The IR and UV spectra were recorded using a Magna 550 spectrophotometer in KBr pellet and a Milton Roy Spectronic 3000 Array spectrophotometer in MeOH soln, respectively. The EI-MS spectra were performed on a Hewlett-Packard 5889A. The  $^1\text{H}$  NMR (300 MHz) and  $^{13}\text{C}$  NMR (75 MHz) spectra were recorded on a Bruker DRX-300 spectrometer and the chemical shifts were referenced to TMS as the internal standard. HMBC and NOESY data were recorded on a Bruker DMX-600 spectrometer. CC was carried out on Kieselgel 60 (Merck No. 9385 and 7729) and LiChroprep RP-18.

#### Plant material

The dried flower buds of *Magnolia fargesii* Cheng was purchased from Il-Shin Pharm. Co. (Taejon, Korea) which imported the material from China. A voucher specimen is deposited in our laboratory (NDC-052).

#### Extraction and isolation

The dried and pulverized flower buds of *M. fargesii* (3 kg) were extracted with MeOH at room temp. for several days. The MeOH extracts were concentrated under red. press. to give a residue (225 g). The residue was partitioned between *n*-hexane (40 g), EtOAc (109 g), *n*-BuOH (20 g) and  $\text{H}_2\text{O}$ , in the usual order. The

EtOAc extract was loaded on silica gel CC eluted with a stepwise solvent gradient of MeOH in  $\text{CHCl}_3$  to afford nineteen subfractions. The subfr. 8 (3 g) was further purified by repeated RP-18 ( $\text{MeOH}-\text{H}_2\text{O}$ , 3:2) and  $\text{SiO}_2$  CC ( $\text{CHCl}_3$ -MeOH, 99:1) to give compound **1** (24 mg).

5 $\alpha$ ,7 $\alpha$ (H)-6,8-cycloeudesma-1 $\beta$ ,4 $\beta$ -diol (**1**). Colorless needles in MeOH; mp 146–147°.  $[\alpha]_{\text{D}}^{25} -4.0^\circ$  (*c* 0.25,  $\text{CHCl}_3$ ). IR  $[\nu]_{\text{max}}^{\text{KBr}}$   $\text{cm}^{-1}$ : 3401, 2949, 1015. EI-MS *m/z* (rel. int.): 238  $[\text{M}]^+$  (6), 220 (51), 205 (21), 202 (18), 195 (14), 187 (40), 177 (54), 159 (100).  $^1\text{H}$  NMR (300 MHz): Table 1.  $^{13}\text{C}$  NMR (75 MHz): Table 1.

#### REFERENCES

1. Namba, T., *Coloured Illustrations of Wakan-Yaku (The Crude Drugs in Japan, China and Neighbouring Countries)*, Vol. II. Hoikusha Publishing, Osaka, 1980, pp. 127–129.
2. Tang, W. and Eisenbrand, G., *Chinese Drugs of Plant Origin: Chemistry, Pharmacology, and Use in Traditional and Modern Medicine*. Springer-Verlag, Berlin, 1992, pp. 639–646.
3. Chen, C. C., Huang, Y. L., Chen, H. T., Chen, Y. P. and Hsu, H. Y., *Planta Medica*, 1988, **54**, 438.
4. Miyazawa, M., Kasahara, H. and Kameoka, H., *Phytochemistry*, 1992, **31**, 3666.
5. Miyazawa, M., Ishikawa, Y., Kasahara, H., Yamanaka, J. I. and Kameoka, H., *Phytochemistry*, 1994, **35**, 611.
6. Miyazawa, M., Kasahara, H. and Kameoka, H., *Phytochemistry*, 1996, **42**, 531.
7. Chen, C. C., Huang, Y. L., Chen, H. T., Chen, Y. P., Hsu, H. Y. and Kuo, Y. H., *Chemical and Pharmaceutical Bulletin*, 1988, **36**, 1791.
8. Kakisawa, H., Kusumi, T., Hsu, H. Y. and Chen, Y. P., *Bulletin of Chemical Society of Japan*, 1970, **43**, 3631.
9. Hwang, Y. L., Chen, C. C., Chen, Y. P., Hsu, H. Y. and Kuo, Y. H., *Planta Medica*, 1990, **56**, 237.
10. Pan, J. X., Hensens, O. D., Zink, D. L., Chang, M. N. and Hwang, S. B., *Phytochemistry*, 1987, **26**, 1377.
11. Jung, K. Y., Kim, D. S., Oh, S. R., Lee, I. S., Lee, J. J., Lee, H. K., Shin, D. H., Kim, E. H. and

- Cheong, C.-J., *Archives Pharmacal Research*, 1997, **20**, 363.
12. Itokawa, H., Matsumoto, H. and Mihashi, S., *Chemistry Letters*, 1983, 1253.
13. Nyasse, B., Tih, R. G., Sondengam, B. L., Martin, M. T. and Bodo, B., *Phytochemistry*, 1988, **27**, 179.
14. Herz, W. and Watanabe, K., *Phytochemistry*, 1983, **22**, 1457.
15. Sung, T. V., Steffan, B., Steglich, W., Klebe, G. and Adam, G., *Phytochemistry*, 1992, **31**, 3515.
16. Toyota, M. and Asakawa, Y., *Phytochemistry*, 1990, **29**, 3664.
17. Zhao, Y., Yue, J.-M., He, Y.-N., Lin, Z.-W. and Sun, H.-D., *Journal of Natural Products*, 1997, **60**, 545.
18. Connolly, J. D. and Hill, R. A., *Directory of Terpenoids, Vol. 1. Mono- and Sesquiterpenoids*, Chapman & Hall, London, 1991, pp. 394–397.
19. Yamamura, S., Niwa, M., Ito, M. and Saito, Y., *Chemistry Letters*, 1982, 1681.