

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/259243886>

H/D Kinetic Isotope Effect as a Tool to Elucidate the Reaction Mechanism of Methyl Radicals with Glycine in Aqueous Solutions

ARTICLE *in* THE JOURNAL OF PHYSICAL CHEMISTRY A · DECEMBER 2013

Impact Factor: 2.69 · DOI: 10.1021/jp4090052 · Source: PubMed

CITATION

1

READS

66

6 AUTHORS, INCLUDING:



Israel Zilbermann

Ben-Gurion University of the Negev

66 PUBLICATIONS 778 CITATIONS

SEE PROFILE



Eric Maimon

Ben-Gurion University of the Negev

42 PUBLICATIONS 240 CITATIONS

SEE PROFILE



Ronen Bar-Ziv

Ben-Gurion University of the Negev

10 PUBLICATIONS 20 CITATIONS

SEE PROFILE



Dan Meyerstein

Ariel University

350 PUBLICATIONS 5,076 CITATIONS

SEE PROFILE

H/D Kinetic Isotope Effect as a Tool to Elucidate the Reaction Mechanism of Methyl Radicals with Glycine in Aqueous Solutions

Guy Yardeni,^{*,†,‡} Israel Zilbermann,^{†,‡} Eric Maimon,^{†,‡} Lioubov Kats,[‡] Ronen Bar-Ziv,^{†,‡} and Dan Meyerstein^{*,‡,§}


[†]Chemistry Department, Nuclear Research Centre Negev, Beer-Sheva, 86000, Israel

[‡]Chemistry Department, Ben-Gurion University of the Negev, Beer-Sheva, 84105, Israel

[§]Biological Chemistry Department, Ariel University, Ariel, 44837, Israel

ABSTRACT: The H/D kinetic isotope effect (KIE) for the reaction of methyl radicals with glycine in aqueous solutions at pH 10.6 equals 16 ± 3 . This result proves that the methyl radical abstracts a hydrogen atom from the methylene group of glycine and not an electron from the unpaired couple on the nitrogen atom. The rate constant of the reaction of methyl radicals with glycine at pH 7.0 is orders of magnitude smaller than that at pH 10.6.

Large Kinetic Isotope Effect:



INTRODUCTION

The reactions of alkyl radicals and alkyl-peroxyl radicals with organic substrates are key steps in a variety of catalytic^{1–4} and physiological processes.^{5,6} The latter are usually initiated by hydroxyl radicals,⁷ which react in very fast reactions with most components of biological systems.⁸ The rate constants of the reactions of hydroxyl radicals with amines and amino acids are pH-dependent, the rate constants for RNH_2 being considerably higher than those for RNH_3^+ .⁹ This increase is usually attributed to the delocalization, reorganization energy of the radical $\text{H}_2\text{NCR}^1\text{R}^{2\bullet}/\text{H}_2\text{NCRCO}_2^{\bullet-}$.^{10,11} However it might also be due to an electron abstraction from the lone pair on the nitrogen of RNH_2 followed by an α H atom shift forming the same radical. Recent results point out that the reaction of hydroxyl radicals with glycine involves three very short-lived intermediates:¹² (a) $\text{H}_2\text{NCH}_2\text{CO}_2^{\bullet-}\cdots\text{HO}^\bullet$ that decomposes to yield the radicals $\text{H}_2\text{NCH}_2\text{CO}_2^{\bullet-}$ and $\bullet\text{NHCH}_2\text{CO}_2^-$; (b) $\text{HO}^\bullet\cdots\text{NH}_2\text{CH}_2\text{CO}_2^-$ or $\text{HO}^{\bullet\cdots} + \bullet\text{NH}_2\text{CH}_2\text{CO}_2^-$ that decomposes to yield the radicals $\text{NH}_2\text{CH}_2^\bullet$ and $\bullet\text{NHCH}_2\text{CO}_2^-$.¹² It seemed therefore of interest to measure the kinetics and mechanism of the simplest alkyl radical, methyl, with the simplest amino acid, glycine, in order of differentiating between the plausible mechanisms. It should be noted that $\bullet\text{CH}_3$ radicals differ from $\bullet\text{OH}$ radicals in two aspects: (a) They are weaker oxidizing agents as the C–H formed in their reactions is weaker than the O–H bond. (b) $\bullet\text{CH}_3$ radicals are hydrophobic and are not expected to form transients with hydrogen bonds, e.g., $\text{H}_2\text{NCH}_2\text{CO}_2^{\bullet-}\cdots\text{HO}^\bullet$. It was decided to measure the kinetic isotope effect (KIE) of the process, as it is well established that H atom abstraction from C–H groups by radicals involves a large KIE.^{13,14} In 1970, Moger et al. estimated that the rate constant of this reaction at pH 11.5 is $120 \text{ M}^{-1} \text{ sec}^{-1}$ and that the KIE = 10.5.¹⁵ These were clearly rough estimates as the results presented herein point out.


MATERIALS AND METHODS

Materials. All solutions were prepared from analytical grade chemicals. Glycine and D_2 -glycine, $\text{H}_2\text{NCD}_2\text{CO}_2\text{H}$, were purchased from Sigma-Aldrich. Dimethyl sulfoxide was purchased from Bio-Lab. The water used was heat distilled and then passed through a Millipore setup, the final resistance being above $15 \text{ M}\Omega \text{ cm}^{-1}$. Solutions were deaerated by $\text{N}_2\text{O}_{(\text{g})}$ purchased from “Maxima”. pH values were measured with a Hanna HI 9321 pH meter and adjusted with HClO_4 and/or NaOH solutions.

Irradiations. γ -Irradiations were carried out in a ^{60}Co γ source, G-220 Gammacell, with a dose rate of 13 Gy min^{-1} , which was determined by means of Fricke dosimetry using a G value of $G = 15.6$.^{16,17}

Gas Chromatographic Analysis. GC analyses were performed using a HP 5890 Series II gas chromatograph with a pack Q column and a flame ionization detector (FID).

Production of Radicals. $\bullet\text{CH}_3$ radicals were prepared by irradiating N_2O -saturated aqueous solutions containing $(\text{CH}_3)_2\text{SO}$ via the following reactions: The primary radicals formed when dilute aqueous solutions are irradiated with ionizing radiation are



In parentheses are given the relative yields of the primary products, expressed as G values.¹⁸ (G values are defined as the number of species of each product formed per 100 eV of

Received: September 9, 2013

Revised: November 28, 2013

Table 1. Yield of Products and Rate Constants for the Reaction of Methyl Radicals with Glycine and Deuterated Glycine^a

solution composition	G(CH ₄)	G(C ₂ H ₆)	G(total) = G(CH ₄) + 2G(C ₂ H ₆)	G(CH ₄)/G(C ₂ H ₆)	k ₇ (M ⁻¹ sec ⁻¹)
0.10 M (CH ₃) ₂ SO, N ₂ O sat., pH 7.0	2.35	1.86	6.06	1.27	
0.0010 M glycine, 0.10 M (CH ₃) ₂ SO, N ₂ O sat., pH 7.0	2.50	1.95	6.39	1.28	k < 3 × 10 ²
0.10 M (CH ₃) ₂ SO, N ₂ O sat., pH 10.6	2.39	2.04	6.47	1.18	
0.0010 M glycine, 0.10 M (CH ₃) ₂ SO, N ₂ O sat., pH 10.6	3.89	1.02	5.92	3.82	(2.0 ± 0.6) × 10 ⁴
0.0010 M D ₂ -glycine, 0.10 M (CH ₃) ₂ SO, N ₂ O sat., pH 10.6	2.82	1.83	6.47	1.55	(1.3 ± 0.4) × 10 ³
0.020 M (CH ₃) ₂ SO, N ₂ O sat., pH 10.6	0.99	2.98	6.95	0.33	
0.0010 M glycine, 0.020 M (CH ₃) ₂ SO, N ₂ O sat., pH 10.6	2.87	0.98	4.82	2.95	(2.5 ± 0.8) × 10 ⁴
0.0010 M D ₂ -glycine, 0.020 M (CH ₃) ₂ SO, N ₂ O sat., pH 10.6	1.39	2.49	6.36	0.56	(1.5 ± 0.4) × 10 ³

^aAll the solutions are N₂O-saturated and were irradiated to 200 Gy.



radiation absorbed by the solution). The distribution of these products in the solution after 1 × 10⁻⁷ s is homogeneous.¹⁸

In N₂O saturated solutions, [N₂O] = 2.2 × 10⁻² M,¹⁹ are irradiated, the hydrated electrons are transformed into •OH radicals via²⁰





Thus, at pH > 3, G(•OH) = 6.0 is obtained.²⁰

In the presence of (CH₃)₂SO the •OH radicals are converted into •CH₃ radicals via²¹



In the absence of other solutes the •CH₃ radicals thus formed are transformed into CH₄ and C₂H₆ via^{22,23}



The relative yields of CH₄ and C₂H₆ thus depend on the concentration of (CH₃)₂SO and the dose rate of irradiation.

RESULTS

Aqueous solutions of glycine and D₂-glycine were irradiated in the presence of (CH₃)₂SO in the ⁶⁰Co γ source, and the yields of the final products (methane and ethane) were measured; the results are summed up in Table 1.

The rate constant of the reaction of the methyl radical with glycine and D₂-glycine were calculated using the assumption that all the ethane is formed via reaction 6:

$$d[C_2H_6]/dt = k_6[\bullet CH_3]_{ss}^2 \quad (I)$$

Where [•CH₃]_{ss} is the steady-state concentration of •CH₃ during the irradiation.

$$d[CH_4]/dt = k_5[\bullet CH_3]_{ss}[(CH_3)_2SO] + k_7[\bullet CH_3]_{ss}[(\text{glycine})] \quad (II)$$


At steady state, i.e., during the irradiation, d[•CH₃]_{ss}/dt = 0. Therefore

$$0 = d[\bullet CH_3]_{ss}/dt = ((\text{dose rate, Gy min}^{-1})G(\bullet CH_3) \times 10^{-7}/60) - 2k[\bullet CH_3]_{ss}^2 - k_5[\bullet CH_3]_{ss}[(CH_3)_2SO] - k_7[\bullet CH_3]_{ss}[(\text{glycine})] \quad (III)$$

The dose rate of the ⁶⁰Co γ source was 13 Gy min⁻¹ and G(•CH₃) = 6.²¹


$$[C_2H_6]/[CH_4] = k_6[\bullet CH_3]_{ss}^2 / (k_5[\bullet CH_3]_{ss}[(CH_3)_2SO] + k_7[\bullet CH_3]_{ss}[(\text{glycine})]) = k_6[\bullet CH_3]_{ss} / (k_5[(CH_3)_2SO] + k_7[(\text{glycine})]) \quad (IV)$$

As k₅, k₆, and all the concentrations are known, measuring the yields of C₂H₆ and CH₄ enables the calculation of k₇.²⁴ (The pK_a of glycine is 9.6.)²⁴



The results clearly point out that the rate constants of reaction 7 at pH 10.6 are several orders of magnitude higher than those at pH 7.0. This result is in accord with an earlier report;¹⁵ however, the rate constants determined in this study are 2 orders of magnitude higher than those previously suggested.¹⁵ The redox potential for (NH₂CHCOO^{•-}/NH₂CH₂COO⁻) is 0.8 V, and the redox potential for (•NH₃CHCOO^{•-}/•NH₃CH₂COO⁻) is 1.5 V,²⁵ explaining the pH effect on k₇. The observation that k₇ for H₂NC(H₂/D₂)CO₂⁻ is orders of magnitude higher than that for •H₃NC(H₂/D₂)CO₂⁻ is in accord with the analogous reaction of •OH radicals with glycine and other amino acids and amines.^{26,27} This points out the effect of the delocalization, reorganization energy of the radical H₂NC(H/D)CO₂^{•-} and the effect of the -NH₃⁺/NH₂ groups on the bond dissociation energy of •H₃NC(H₂/D₂)CO₂⁻/H₂NC(H₂/D₂)CO₂⁻ on the rate of the HAT.

The large H/D KIE measured at pH 10.6, k_H/k_D = 16.0 ± 3.0, clearly proves that the rate-determining step of reaction 7 is the HAT and not the electron-transfer reaction



The large KIE is in accord with the large KIE reported for hydrogen atom abstraction from C–H groups.^{13,28} However, the KIE for reaction 7 is considerably smaller than that reported for the reaction of H atoms with methanol and acetate.¹³ This is somewhat surprising as k₇ is considerably smaller than the rate constants of the reactions of H• atoms with methanol and acetate, and the KIE for the reactions of H• atoms increases with the decrease in the rate constants of the reactions.¹³ One would expect a similar behavior for the reactions of •CH₃ and H• as the bond energies H–H and H–CH₃ are similar.^{29,30} Alternatively, it might be suggested that some of the radicals react via reaction 7', as happens for the •OH radicals,¹² and that the KIE for reaction 7 is considerably larger than 16. As •CH₃ radicals are weaker oxidizing agents, one expects that k₇/k_{7'} for •CH₃ radicals will be larger than that for •OH radicals as the radical H₂NC(H/D)CO₂⁻ is considerably more stable than •NH₂C(H₂/D₂)CO₂⁻.

Finally the large H/D KIE clearly points out that the HAT involves tunneling.³¹ For the slow reaction, e.g., reaction 7, clearly tunneling accelerates the rate constants at least by an order of magnitude, assuming that tunneling does not contribute

to deuterium abstraction reactions. This conclusion is surprising as the valence bond calculated activation energies, $\Delta E_{\text{VB}}^{\#}$, correlate linearly with the experimental activation energies, $\Delta G_{\text{exp}}^{\#}$ for HAT reactions.³²

CONCLUSIONS

1. Methyl radicals react with glycine via a mechanism that involves, at least mainly, a H^{\bullet} abstraction from the methylene group, as pointed out by the large KIE, and not, or only some, an e^{-} abstraction from the nitrogen amine.
2. The rate of this reaction is pH-dependent, increasing by several orders of magnitude above the last pK_{a} of glycine. Thus, probably even at physiological pH, that is, ca. 3 pH units below the pK_{a} of amino acids, alkyl radicals react with the alkaline form of amino acids present in the medium.

AUTHOR INFORMATION

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This study was supported by a grant from the Israeli Atomic Energy Committee and the Planning and Budgeting committee of the Israel Council of Higher Education.

REFERENCES

- (1) Cornlis, B.; Herrmann, W. A. *Applied Homogeneous Catalysis with Organometallic Compounds*; VCH: Weinheim, Germany, 1996.
- (2) Goldstein, S.; Meyerstein, D.; Czapski, G. The Fenton Reagents. *Free Radical Biol. Med.* **1993**, *15*, 435–445.
- (3) Goldstein, S.; Meyerstein, D. Comments on the Mechanism of the “Fenton-Like” Reaction. *Acc. Chem. Res.* **1999**, *32*, 547–550.
- (4) Simandi, L. I. *Catalytic Activation of Dioxygen by Metal Complexes*; Kluwer Academic Publishers: Dordrecht, The Netherlands, 1992.
- (5) Neta, P.; Huie, R. E.; Ross, A. B. Rate Constants for Reactions of Peroxyl Radicals in Fluid Solutions. *J. Phys. Chem. Ref. Data* **1990**, *19*, 413–513.
- (6) Meyerstein, D. Reactions of Aliphatic Carbon-Centered and Aliphatic Peroxyl Radicals with Transition Metal Complexes as a Plausible Source for Biological Damage Induced by Radical Processes. *Met. Ions Biol. Syst.* **1999**, *36*, 41–77.
- (7) Bauer, G.; Chatgililoglu, C.; Gebicki, J. L.; Gebicka, L.; Gescheidt, G.; Golding, B. T.; Goldstein, S.; Kaizer, J.; Merenyi, G.; Speier, G.; Wardman, P. Biologically Relevant Small Radicals. *Chimia* **2008**, *62*, 704–712.
- (8) Dorfman, L. M.; Adams, G. E. *Reactivity of the Hydroxyl Radical in Aqueous Solutions*; National Standard Reference Data System, U.S. Department of Commerce and National Bureau of Standards: United States, 1973.
- (9) Simic, M.; Neta, P.; Hayon, E. Pulse Radiolytic Investigation of Aliphatic Amines in Aqueous Solutions. *Int. Radiat. Phys. Chem.* **1971**, *3*, 309–320.
- (10) Rega, N.; Cossi, M.; Barone, V. Structure and Magnetic Properties of Glycine Radical in Solution at Different pH Values. *J. Am. Chem. Soc.* **1998**, *120*, 5723–5732.
- (11) Armstrong, D. A.; Rauk, A.; Yu, D. Solution Thermochemistry of the Radicals of Glycine. *J. Chem. Soc. Perkin Trans. 2* **1995**, 553–559.
- (12) Stefanic, I.; Ljubic, I.; Bonifacic, M.; Sabljic, A.; Asmus, K. D.; Armstrong, D. A. A Surprisingly Complex Aqueous Chemistry of the Simplest Amino Acid. A pulse Radiolysis and Theoretical Study on H/D Kinetic Isotope Effects in the Reaction of Glycine Anions with Hydroxyl Radicals. *Phys. Chem. Chem. Phys.* **2009**, *11*, 2256–2267.
- (13) Anbar, M.; Meyerstein, D. Isotope Effects in the Hydrogen Abstraction from Aliphatic Compounds by Radiolytically Produced Hydrogen Atoms in Aqueous Solutions. *J. Phys. Chem.* **1964**, *68*, 3184–3187.
- (14) Geimer, J.; Beckert, D.; Jenichen, A. Kinetic Isotope Effect in Hydrogen Abstraction from 2-propanol by Hydrogen and Deuterium Radicals. A Pulse Radiolysis Fourier Transform Electron Spin Resonance Study. *Chem. Phys. Lett.* **1997**, *280*, 353–358.
- (15) Moger, G.; Dobis, O. The Reaction of Free Methyl Radicals with Glycine in Aqueous Solution. *Magy. Kem. Foly.* **1970**, *76* (7), 328–332.
- (16) Hart, E. J. Radiation Chemistry of Aqueous Ferrous Sulfate-Cupric Sulfate Solutions; Effect of Gamma-Rays. *Radiat. Res.* **1955**, *2* (1), 33–46.
- (17) Fricke, H.; Hart, E. J. *Chemical Dosimetry in Radiation Dosimetry*; Attix, F. H., Roesch, W. C., Tochilin, E., Eds.; Academic Press: New York, 1966; p 167.
- (18) Matheson, M. S.; Dorfman, L. *Pulse Radiolysis*; MIT Press: Cambridge, MA, 1969.
- (19) Buxton, G. V.; Greenstock, C. L.; Helman, W. P.; Ross, A. B. Critical Review of Rate Constant for Reactions of Hydrated Electrons, Hydrogen Atoms and Hydroxyl Radicals ($\bullet\text{OH}/\bullet\text{O}^{-}$) in Aqueous Solutions. *J. Phys. Chem. Ref. Data* **1988**, *17*, 513–886.
- (20) Ross, A. B.; Mallard, W. G.; Helman, W. P.; Buxton, J. V.; Huie, R. E.; Neta, P. *NIST Standard References Database*, version 2.0; 1994.
- (21) Veltwisch, D.; Janata, E.; Asmus, K. D. Primary Processes in the Reaction of OH^{\bullet} -Radicals with Sulphoxides. *J. Chem. Soc. Perkin Trans. 2* **1980**, 146–153.
- (22) Rusonic, I.; Polat, H.; Cohen, H.; Meyerstein, D. Reaction of Methyl Radicals with Metal Powders Immersed in Aqueous Solutions. *Eur. J. Inorg. Chem.* **2003**, 4227–4233.
- (23) Hickel, B. Absorption Spectra and Kinetics of Methyl and Ethyl Radicals in Water. *J. Phys. Chem.* **1975**, *79*, 1054–1059.
- (24) Bonifacic, M.; Stefanic, I.; Li, H. G.; Armstrong, D. A.; Asmus, K. D. Glycine Decarboxylation: The Free Radical Mechanism. *J. Am. Chem. Soc.* **1998**, *120*, 9930–9940.
- (25) Armstrong, D. A.; Yu, D.; Rauk, A. Oxidative Damage to the Glycine α -Carbon Site in Proteins: An Ab-Initio Study of the C–H Bond Dissociation Energy and the Reduction Potential of the C-Centred Radical. *Can. J. Chem.* **1996**, *74*, 1192–1199.
- (26) Mossoba, M. M.; Rosental, I.; Riesz, P. Electron Spin Resonance of Spin-Trapped Radicals of Amines and Polyamines. Hydroxyl Radical Reactions in Aqueous Solutions and γ -Radiolysis in the Solid State. *Can. J. Chem.* **1982**, *60*, 1493–1500.
- (27) Bonifacic, M.; Armstrong, D. A.; Carmichael, I.; Asmus, K. D. β -Fragmentation and Other Reactions Involving Aminyl Radicals from Amino Acids. *J. Phys. Chem. B* **2000**, *104*, 643–649.
- (28) Klinker, E. J.; Shaik, S.; Hirao, H.; Que, L., Jr. Two-State Reactivity Model Explains Unusual Kinetic Isotope Effect Patterns in C–H Bond Cleavage by Nonheme Oxoiron(IV) Complexes. *Angew. Chem., Int. Ed.* **2009**, *48*, 1291–1295.
- (29) Branchaud, B. P. Free Radicals as a Result of Dioxygen Metabolism. *Met. Ions Biol. Syst.* **1999**, *36*, 79–102.
- (30) Neta, P.; Grodkowski, J.; Ross, A. B. Rate Constants for Reactions of Aliphatic Carbon-Centered Radicals in Aqueous Solution. *J. Phys. Chem. Ref. Data* **1996**, *25*, 709–1050.
- (31) Antoniou, D.; Schwartz, S. D. Large Kinetic Isotope Effects in Enzymatic Proton Transfer and the Role of Substrate Oscillations. *Proc. Natl. Acad. Sci. U.S.A.* **1997**, *94*, 12360–12365.
- (32) Lai, W.; Li, C.; Chen, H.; Shaik, S. Hydrogen-Abstraction Reactivity Patterns from A to Y: The Valence Bond Way. *Angew. Chem., Int. Ed.* **2012**, *51*, 5556–5578.