

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/236342680>

The Standard Enthalpy of Formation of the 1-Adamantyl Cation in the Gas Phase. An Experimental and ab Initio Re-assessment

ARTICLE *in* THE JOURNAL OF PHYSICAL CHEMISTRY A · NOVEMBER 1999

Impact Factor: 2.69 · DOI: 10.1021/jp991262u

CITATIONS

12

READS

26

8 AUTHORS, INCLUDING:



Juan Z Dávalos

Institute of Physical Chemistry Rocasolano

126 PUBLICATIONS 1,278 CITATIONS

SEE PROFILE



Jose-Luis M. Abboud

Spanish National Research Council

266 PUBLICATIONS 8,725 CITATIONS

SEE PROFILE

LETTERS


The Standard Enthalpy of Formation of the 1-Adamantyl Cation in the Gas Phase. An Experimental and ab Initio Re-assessment^{||}

H. Flores,[†] J. Z. Dávalos,[†] J.-L. M. Abboud,^{*,†} O. Castaño,^{*,‡} R. Gomperts,^{*,§} P. Jiménez,^{*,†} R. Notario,^{*,†} and M. V. Roux^{*,†}

Instituto de Química Física "Rocasolano", CSIC, E-28006 Madrid, Spain, Departamento de Química Física, Universidad de Alcalá, E-28871 Alcalá de Henares, Madrid, Spain, and Silicon Graphics, Inc., One Cabot Road, Hudson, Massachusetts 01749

Received: April 19, 1999; In Final Form: July 28, 1999


The standard molar enthalpy of formation of gaseous 1-adamantyl chloride (1Cl,g) was determined experimentally by means of thermochemical techniques. Ab initio quantum-chemical calculations at the G2-(MP2) level were performed on adamantyl cation (**1**⁺) and on 1Cl. Combination of these data with other experimental and computational results for adamantane (**1H**), isobutane (**2H**), *tert*-butyl cation (**2**⁺), and *tert*-butyl chloride (**2Cl**) provided purely experimental and purely computational, independent values for the standard enthalpy and Gibbs energy changes for reactions 1 and 2:



where X = Cl. This information was used to determine a value of $162.0 \pm 2.0 \text{ kcal mol}^{-1}$ for the standard molar enthalpy of formation of gaseous **1**⁺, $\Delta_f H_m^\circ(\mathbf{1}^+, \text{g})$.

Holmes and co-workers¹ have recently used appearance energies to determine the standard enthalpy of formation of 1-adamantyl cation (**1**⁺) in the gas phase, $\Delta_f H_m^\circ(\mathbf{1}^+, \text{g})$. The value they reported, $152 \pm 3 \text{ kcal mol}^{-1}$, significantly differs from the values obtained by other methods (in the range $158\text{--}164 \text{ kcal mol}^{-1}$).¹ Inasmuch as carbocations are important chemical species,^{2,3} this discrepancy has prompted us to reexamine the problem by means of experimental and computational techniques.

Equilibrium techniques, notably ICR (ion cyclotron resonance



* Authors to whom correspondence should be addressed.

[†] Instituto de Química Física "Rocasolano", CSIC.

[‡] Universidad de Alcalá.

[§] Silicon Graphics, Inc.

^{||} This work is dedicated to Prof. Louis Bellon.

spectroscopy⁴), FT ICR (Fourier transform ion cyclotron

TABLE 1: Experimentally Determined Standard Molar Energy of Combustion and Standard Molar Enthalpies of Combustion, Sublimation, and Formation in the Crystalline and Gaseous State at Temperature $T = 298.15$ K^{a,b} of 1-Adamantyl Chloride

$\Delta_c U_m^c$	-1398.92 ± 0.48
$\Delta_c H_m^d$	-1401.00 ± 0.48
$\Delta_{\text{sub}} H_m^e$	15.11 ± 0.17
$\Delta_f H_m^f(\text{cr})$	-57.55 ± 0.57
$\Delta_f H_m^g(\text{g})$	-42.45 ± 0.60

^a All values in kcal/mol, (1 cal = 4.184 J). ^b This work. ^c Standard molar energy of combustion. ^d Standard molar enthalpy of combustion. ^e Standard molar enthalpy of sublimation. ^f Standard molar enthalpy of formation in the crystalline state. ^g Standard molar enthalpy of formation in the gaseous state.

resonance spectroscopy⁵), and HPMS (high-pressure mass spectrometry⁶) have been used to determine the standard enthalpy and Gibbs energy changes for the hydride and halide exchange reactions between (1^+ ,g) and *tert*-butyl cation (2^+ ,g) in the gas phase, reactions 1 and 2: where X = Cl, Br.

Reliable experimental $\Delta_f H_m^g(\text{g})$ values are presently available for adamantane (1H), isobutane (2H), and 2^+ .⁷ In view of this, we have carried out the experimental determination of the standard enthalpy of formation of gaseous 1-adamantyl chloride, $\Delta_f H_m^g(\text{1Cl,g})$. *Combination of this datum with the other relevant experimental values involved in reaction 2 (X = Cl), leads to a new, purely experimental value of $\Delta_f H_m^g(1^+,g)$.* This value can be usefully compared to that obtained using the experimental thermodynamic data involved in reaction 1.

High-level ab initio methodologies, particularly the G2⁸ and G2(MP2)⁹ treatments have been shown to yield values of the changes in thermodynamic state functions for chemical reactions agreeing within 0.1 eV (2.3 kcal mol⁻¹) with the experimental values. In the context of the present study, we thought it worthwhile to carry out the study of 1H, 2H, 1Cl, 2Cl, 1^+ , and 2^+ at the G2(MP2) level. *This computational effort provides values of $\Delta H^\circ(1)$, $\Delta G^\circ(1)$, $\Delta H^\circ(2, X = \text{Cl})$ and $\Delta G^\circ(2, X = \text{Cl})$ fully independent from experiment.* They are used here in order to cross-check the experimental results.

$\Delta_f H_m^g(\text{1Cl,g})$ was determined in this work by combining the standard enthalpy of formation of crystalline 1Cl, namely $\Delta_f H_m^g(\text{1Cl,cr})$, with its standard enthalpy of sublimation, $\Delta_{\text{sub}} H_m^g(\text{1Cl})$. $\Delta_f H_m^g(\text{1Cl,cr})$ was obtained by combustion calorimetry, using a rotary-bomb calorimeter.¹⁰ $\Delta_{\text{sub}} H_m^g(\text{1Cl})$ was deduced from the temperature dependence of its vapor pressure (Clausius–Clapeyron)¹¹ over a 20 K temperature interval. The vapor pressure of the compound was measured by means of the Knudsen effusion technique.¹¹ Heat capacity measurements were carried out by differential scanning calorimetry.¹² Full experimental details are given as Supporting Information and Table 1 summarizes the results of the thermochemical study of 1Cl.

Table 2 collects the results of the G2(MP2) calculations. Full details are also given as Supporting Material.

$\Delta_f H_m^g(\text{g})$ values taken from the literature^{7,13} for adamantane, isobutane, *tert*-butyl chloride, and *tert*-butyl cation are, respectively, -32.0 ± 0.9 , -32.25 ± 0.50 , -42.99 ± 0.50 , and 169.9 ± 0.9 kcal mol⁻¹. The results of the combination of these data are summarized in Tables 3 and 4.

Examination of these results leads to the following conclusions: (i) The agreement between the experimental and calculated changes in thermodynamic state functions for reactions 1 and 2 (X = Cl) is very good, particularly in the case of the Gibbs energies. Notice that these are the primary experimental data for these reactions, the enthalpy changes being obtained from the corresponding van't Hoff plots (experimental uncertainties estimated at ± 1 –2 kcal mol⁻¹).

TABLE 2: G2(MP2) Thermodynamic Functions, in Hartree, for All the Species Used in This Work

compound	H_{298}	G_{298}
1-adamantyl cation, 1^+	-388.99432	-389.03284
adamantane, 1H	-389.88158 ^a	-389.91857
1-adamantyl chloride, 1Cl	-849.03829	-849.07974
<i>tert</i> -butyl cation, 2^+	-157.16933 ^b	-157.20664 ^b
isobutane, 2H	-158.07400	-158.10855
<i>tert</i> -butyl chloride, 2Cl	-617.22672 ^b	-617.26407 ^b

^a Value taken from Castaño, O.; Notario, R.; Abboud, J.-L. M.; Gomperts, R.; Palmeiro, R.; Frutos, L. M. *J. Org. Chem.*, submitted. ^b Value taken from Abboud, J.-L. M.; Castaño, O.; Herreros, M.; Leito, I.; Notario, R.; Sak, K. *J. Org. Chem.* **1998**, 63, 8995–8997.

TABLE 3: Determination of $\Delta_f H_m^g(1^+,g)$ on the Basis of Reaction 1^a

$\Delta G^\circ(1)$	$\Delta H^\circ(1)$	origin	$\Delta_f H_m^g(1^+,g)^d$
-8.2 ± 0.1 (300 K)	-7.6 ± 1.0	HPMS ^b	162.6 ± 3.0
-10.2	-10.9	G2(MP2) ^c	159.3 ± 2.8

^a All values in kcal mol⁻¹. ^b From ref 14. ^c This work. ^d Uncertainties taken as twice the standard deviation.

TABLE 4: Determination of $\Delta_f H_m^g(1^+,g)$ on the Basis of Reaction 2^a

$\Delta G^\circ(2, X = \text{Cl})$	$\Delta H^\circ(2, X = \text{Cl})$	origin	$\Delta_f H_m^g(1^+,g)^f$
-5.9 ± 0.1 (600 K)	-6.3 ± 1.0	HPMS ^b	164.1 ± 3.2
-6.5 ± 1.0 (300 K) ^c			
-5.5 ± 0.4 (333 K)		FT ICR ^d	
-6.6 (298 K)	-8.4	G2(MP2) ^e	162.0 ± 2.4

^a All values in kcal mol⁻¹. ^b From ref 14. ^c Calculated using the ΔH° and ΔS° values given in ref 14. ^d See ref 15. ^e This work. ^f Uncertainties taken as twice the standard deviation.

(ii) Using the G2(MP2) calculation on 1Cl, we obtain a purely computational estimate of $\Delta_f H_m^g(\text{1Cl,g})$, namely, -44.7 kcal mol⁻¹ found using atomization reaction and spin–orbit correction.¹⁷ This result compares favorably with the experimental value of -42.5 kcal mol⁻¹ found in this work.

(iii) The value of $\Delta G^\circ(2, X = \text{Cl})$ determined by FT ICR is quite consistent with that obtained by means of HPMS. This strongly supports the present treatment.

(iv) The average of the values of $\Delta_f H_m^g(1^+,g)$ based on purely experimental data is 163.4 kcal mol⁻¹. The corresponding average based on the computed values of $\Delta H^\circ(1)$ and $\Delta H^\circ(2, X = \text{Cl})$ is 160.7 kcal mol⁻¹. These results are also consistent in view of the experimental uncertainties involved. We consider that the average of these data, 162.0 ± 2.0 kcal mol⁻¹ can be taken as the final value of $\Delta_f H_m^g(1^+,g)$ as determined by the methods indicated in this study.

(v) This result is significantly different from the value $\Delta_f H_m^g(1^+,g) = 152 \pm 3$ kcal mol⁻¹ reported in ref 1.

(vi) A referee has pointed out that the combination of $\Delta_f H_m^g(1^+,g) = 162.0$ kcal mol⁻¹ with the available experimental ionization energy (determined by PES, photoelectron spectroscopy) of the 1-adamantyl radical (1^\bullet ,g), namely, 6.21 eV¹⁸ leads to a standard enthalpy of formation of gaseous 1^\bullet , $\Delta_f H_m^g(1^\bullet,g)$, of 19 kcal mol⁻¹. In turn, this makes the C(tertiary)–H bond dissociation energy in adamantane as high as 103 kcal mol⁻¹. This value is substantially larger than the corresponding value in 2H, 95.9 kcal mol⁻¹.¹⁹ On the other hand, the value 152 kcal mol⁻¹ for $\Delta_f H_m^g(1^+,g)$ leads to a dissociation energy of the C(tertiary)–H bond in 1H of 96.3 kcal mol⁻¹,¹⁹ this result seeming physically more sound.

In view of the above, we have performed high-level ab initio calculations on 1^\bullet and 2^\bullet .²⁰ They do confirm the referee's contention that both C–H linkages have rather similar bond

strengths. The situation is, however, more complicated. Thus, the value for $\Delta_f H_m^\circ(2^+, g)$ currently accepted by NIST^{7,21} and used here and in ref 1 is 169.9 kcal mol⁻¹. If we combine this datum with the value of $\Delta_f H_m^\circ(2^*, g)$ of 11.0 kcal mol⁻¹⁷ and the available PES ionization energies, namely, 6.70^{7,22} and 6.58 eV,^{7,23} we obtain bond strengths for the tertiary C–H bond in 2H of 99.2 and 102.0 kcal mol⁻¹, respectively. Therefore, the criterion of the PE ionization energy of 1• is to be considered with great circumspection.

Acknowledgment. This work was supported by Grant PB96-0927-C02-01 of the Spanish D.G.E.S. We thank a referee for valuable comments.

References and Notes

- (1) Aubry, C.; Holmes, J. L.; Walton, J. C. *J. Phys. Chem. A* **1998**, *102*, 1389–1393.
- (2) See, for example, (a) *Stable Carbocation Chemistry*; Surya Prakash, G. K., Schleyer, P. v. R., Eds.; Wiley: New York, 1997. (b) Abboud, J.-L. M.; Castaño, O.; Della, E. W.; Herreros, M.; Müller, P.; Notario, R.; Rossier, J.-C. *J. Am. Chem. Soc.* **1997**, *119*, 2262–2266, and references therein.
- (3) Corma, A.; García, H. *Top. Catal.* **1998**, *6*, 127–140.
- (4) (a) Lehman, T. A.; Bursey, M. M. *Ion Cyclotron Resonance Spectroscopy*; John Wiley & Sons: New York, 1976. (b) Staley, R. H.; Wieting, R. D.; Beauchamp, J. L. *J. Am. Chem. Soc.* **1977**, *99*, 5964–5972.
- (5) (a) Laukien, F. H.; Allemann, M.; Bischofberger, P.; Grossmann, P.; Kellerhals, H. P.; Kopf, P. In *Fourier Transform Mass Spectrometry. Evolution, Innovation and Applications*. Buchanan, M. V. Ed.; ACS Symposium Series 359; American Chemical Society: Washington, DC, 1987; Chapter 5. (b) *FT-ICR/MS Analytical Applications of Fourier Transform Ion Cyclotron Resonance Mass Spectrometry*; Asamoto, B., Ed.; VCH Publishers: New York, 1991.
- (6) (a) Kebarle, P. *Annu. Rev. Phys. Chem.* **1977**, *28*, 445–476. (b) For a recent review, see, for example, Kebarle, P. In *Techniques for the Study of Ion–Molecule Reactions*; Farrar, J. M., Saunders, W., Jr., Eds.; Wiley: New York, 1988; Chapter 5.
- (7) Mallard, W. G.; Linstrom, P. J., Eds. *NIST Chemistry Webbook, NIST Standard Reference Database*, Number 69, November 1998, National Institute of Standards and Technology: Gaithersburg, MD 20899 (<http://webbook.nist.gov>).
- (8) Curtiss, L. A.; Raghavachari, K.; Trucks, G. W.; Pople, J. A. *J. Chem. Phys.* **1991**, *94*, 7221–7230.
- (9) Curtiss, L. A.; Raghavachari, K.; Pople, J. A. *J. Chem. Phys.* **1993**, *98*, 1293–1298.
- (10) Roux, M. V.; Dávalos, J. Z.; Jiménez, P.; Flores, H.; Juaristi, E.; Saiz, J. L.; Abboud, J.-L. M. *J. Chem. Thermodyn.*, in press.
- (11) Jiménez, P.; Menéndez, V.; Roux, M. V.; Turrión, C. *J. Chem. Thermodyn.* **1995**, *27*, 679–683, and references therein.
- (12) Roux, M. V.; Jiménez, P.; Dávalos, J. Z.; Abboud, J.-L.; Molina, M. T. *J. Chem. Thermodyn.* **1996**, *28*, 1029–1035.
- (13) Taken from ref 7. Whenever different values of comparable quality are reported for the same species, we use the average of these values.
- (14) Sharma, R. B.; Sen Sharma, D. K.; Hiraoka, K.; Kebarle, P. *J. Am. Chem. Soc.* **1985**, *107*, 3747–3757, and references therein.
- (15) This result is obtained by combining the value of $\Delta G^\circ(2, X = Br)$, -5.3 ± 0.4 kcal mol⁻¹, obtained (in two steps) by equilibrium FT ICR,^{2b} with $\Delta G^\circ(3)$, the Gibbs energy change for the isodesmic reaction $3: 1Cl(g) + 2Br(g) \rightarrow 1Br(g) + 2Cl(g)$. The latter was obtained computationally at the HF/6-31G(d) level¹⁶ and amounts to -0.20 kcal mol⁻¹.
- (16) Abboud, J.-L. M.; Herreros, M.; Notario, R.; Lomas, J. S.; Mareda, J.; Müller, P.; Rossier, J.-C. *J. Org. Chem.*, in press.
- (17) Curtiss, L. A.; Raghavachari, K.; Redfern, P. C.; Pople, J. A. *J. Chem. Phys.* **1997**, *106*, 1063–1079.
- (18) Kruppa, G. H.; Beauchamp, J. L. *J. Am. Chem. Soc.* **1986**, *108*, 2162–2169.
- (19) See ref 1, and references therein.
- (20) At the UG2(MP2) level. Gomperts, R.; Castaño, O.; Notario, R.; Abboud, J.-L. M. To be published.
- (21) (a) HPMS value from Szulejko, J. E.; McMahon, T. B. *J. Am. Chem. Soc.* **1993**, *115*, 4448–4456. (b) This value is in excellent agreement with G2-level data from Smith, B. J.; Radom, L. *J. Am. Chem. Soc.* **1993**, *115*, 4448–4456, and with PEPICO studies, Keister, J. W.; Riley, J. S.; Baer, T. *J. Am. Chem. Soc.* **1993**, *115*, 12613–12614.
- (22) Houle, F. A.; Beauchamp, J. L. *J. Am. Chem. Soc.* **1979**, *101*, 4067–4074.
- (23) Dyke, J.; Jonathan, N.; Lee, E.; Morris, A.; Winter, M. *Phys. Scr.* **1977**, *16*, 197, reported in refs 7 and 22.