

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/7393693>

# Antimycobacterial Scalarane-Based Sesterterpenes from the Red Sea Sponge *Hyrtios erecta*

ARTICLE in JOURNAL OF NATURAL PRODUCTS · JANUARY 2006

Impact Factor: 3.8 · DOI: 10.1021/np0502645 · Source: PubMed

CITATIONS

22

READS

71

## 3 AUTHORS:



**Diaa Youssef**

King Abdulaziz University

111 PUBLICATIONS 1,350 CITATIONS

SEE PROFILE



**Lamiaa A Shaala**

King Abdulaziz University

42 PUBLICATIONS 322 CITATIONS

SEE PROFILE



**Samy Emara**

Misr International University

65 PUBLICATIONS 528 CITATIONS

SEE PROFILE

# Antimycobacterial Scalarane-Based Sesterterpenes from the Red Sea Sponge *Hyrtios erecta*

Diaa T. A. Youssef,<sup>\*,†</sup> Lamiaa A. Shaala,<sup>†</sup> and Samy Emara<sup>‡</sup>

Department of Pharmacognosy and Department of Pharmaceutical Analytical Chemistry, Faculty of Pharmacy, Suez Canal University, Ismailia 41522, Egypt


Received July 23, 2005

A new scalarane-type pentacyclic sesterterpene, sesterstatin 7 (**1**), was isolated from the Red Sea sponge *Hyrtios erecta*, together with 16-*epi*-scalarolbutenolide (**2**), 25-dehydroxy-12-*epi*-deacetylscalarin (**3**), 3-acetylsesterstatin 1 (**4**), and 21-acetoxydeoxyscalarin. The structure of **1** was elucidated by spectroscopic data interpretation. Sesterstatin 7 (**1**) showed 63% inhibition of *Mycobacterium tuberculosis* (H<sub>37</sub>Rv) at a concentration of 6.25  $\mu$ g/mL. Compound **2** displayed moderate inhibitory activity, while **3** and **4** were weakly active against the same biological target.

Marine organisms have attracted considerable attention as a source of novel natural products with intriguing structures and useful biological activities.<sup>1</sup> Many scalarane-type sesterterpenoids have been isolated from marine sponges belonging to the order Dictyoceratida.<sup>1</sup> Scalarane-type sesterterpenes display a variety of biological activities such as cytotoxic,<sup>2–4a,b</sup> antimicrobial,<sup>5,6</sup> antifeedant,<sup>7</sup> ichthyotoxic,<sup>8</sup> anti-inflammatory,<sup>9,10</sup> and platelet-aggregation inhibitory effects,<sup>11</sup> as well as nerve growth factor synthesis-stimulating action.<sup>12</sup> The genus *Hyrtios* has proven to be a rich source of structurally diverse substances with potentially useful biological activities including terpenoids,<sup>2–4</sup> macrolides,<sup>3a,13</sup> and tryptamine-derived alkaloids.<sup>14</sup>

In the course of our ongoing efforts to locate drug leads from Red Sea marine invertebrates, we have studied an extract of the Red Sea sponge *Hyrtios erecta*. We report herein the identification and antimycobacterial activity of several scalarane-type sesterterpenes including the new compound sesterstatin 7 (**1**), together with the previously reported compounds, 16-*epi*-scalarolbutenolide (**2**),<sup>15</sup> 25-dehydroxy-12-*epi*-deacetylscalarin (**3**),<sup>16</sup> 3-acetylsesterstatin 1 (**4**),<sup>2</sup> and 21-acetoxydeoxyscalarin.<sup>17</sup> Sesterstatins 1–6 were reported from the sponge *H. erecta* collected in the Republic of Maldives.<sup>3</sup> In addition,  $\beta$ -acetylsesterstatin 1 and 21-acetylsesterstatin 3 were isolated from a Red Sea collection of the sponge *H. erecta*.<sup>2</sup> The previously known sesterstatins exhibit cytotoxic activity against several cancer cell lines.<sup>2,3</sup>

Sesterstatin 7 (**1**) was isolated as a white solid with a molecular formula of C<sub>27</sub>H<sub>40</sub>O<sub>5</sub>, as established from its HRFABMS and <sup>13</sup>C NMR data. Its <sup>1</sup>H NMR spectrum (Table 1) displayed resonances for 40 protons including six singlets assigned to six methyl groups ( $\delta$  0.83, 0.80, 0.89, 0.83, 1.19, and 2.12), eight methylenes, five aliphatic methines, and an exchangeable singlet at  $\delta$  5.75 for an OH moiety. The <sup>13</sup>C NMR spectrum revealed signals for 27 carbons including six methyls, eight methylenes, five methines, and eight quaternary carbons. Analysis of the <sup>1</sup>H,<sup>1</sup>H-COSY and the HMQC NMR experiments led to the assembly of the following structural fragments: C-1 to C-3; C-5 to C-7; C-9 to C-12 with a hydroxyl group at C-12; and C-14 to C-16 with an acetyl moiety at C-16. The proton



signals at  $\delta$  4.76 (H-24a) and 4.71 (H-24b) ( $J_{24a/b}$  = 18.3 Hz) together with the carbon resonances at  $\delta$  70.5 (CH<sub>2</sub>, C-24), 158.5 (qC, C-17), 139.2 (qC, C-19), and 174.9 (qC, C-25) were representative of an  $\alpha,\beta$ -unsaturated butenolide moiety in **1**.<sup>2,3a</sup> This was supported by interpretation of the HMBC spectrum (Table 1), which indicated that C-16 and C-24 were connected to the sp<sup>2</sup> carbon C-17, with these situated, along with C-18 and C-25, in the  $\alpha,\beta$ -unsaturated lactone ring.<sup>2,3a</sup> Connectivities of the five ring systems of **1**, as well as the assignments of all quaternary carbons, were supported unequivocally by HMBC data (Table 1). In addition, the placements of the OH and acetyl moieties at C-12 and C-16, respectively, were secured from HMBC correlations of H-11/C-12, H-16/C-17, and H-16/C-18. Moreover, COSY cross-peaks of H-12/H-11a, H-12/H-11b, H-16/H-15a, and H-16/H-15b supported these assignments. The equatorial geometry of OH at C-12 was established from the large coupling constant of H-12 ( $J$  = 11.5 Hz).<sup>2,3a</sup> Similarly, the  $\beta$ -configuration of the acetyl moiety at C-16 was established from the coupling constant ( $J$  = 10.3 Hz).<sup>15</sup> From the above discussion, compound **1** was assigned as 16 $\beta$ -acetoxydeoxyscalarolide.

Compounds **2–4** and 21-acetoxydeoxyscalarin were identified by interpretation of their spectroscopic data, including HRMS and high-field NMR, and by comparing these data with literature values.<sup>2,15–17</sup>

Compounds **1–4** were evaluated for their in vitro anti-tuberculosis activity at a concentration of 6.25  $\mu$ g/mL,

\* To whom correspondence should be addressed. Tel: +20-64-3342486. Fax: +20-64-3355741. E-mail: youssefdiaa@hotmail.com.

<sup>†</sup> Department of Pharmacognosy.

<sup>‡</sup> Department of Pharmaceutical Analytical Chemistry.

**Table 1.**  $^1\text{H}$  and  $^{13}\text{C}$  NMR Data of **1** ( $\text{CDCl}_3$ )

position	$\delta_{\text{C}}$ (mult.)	$\delta_{\text{H}}$ [mult., $J$ (Hz)]	HMBC <sup>a</sup>
1	39.6, $\text{CH}_2$	1.73, m, 0.79, m	C-10
2	18.5, $\text{CH}_2$	1.56, m, 1.42, m	C-4, C-10
3	42.0, $\text{CH}_2$	1.79, m 1.11, ddd (13.5, 13.5, 4.0)	C-4
4	33.2, qC		
5	56.6, CH	0.79, m	C-4
6	18.1, $\text{CH}_2$	1.57, m, 1.41, m	
7	41.6, $\text{CH}_2$	1.41, m, 0.94, m	C-8
8	37.1, qC		
9	58.0, CH	0.88, m	C-10, C-12
10	37.4, qC		
11	25.7, $\text{CH}_2$	1.88, ddd (11.5, 4.2, 1.8) 1.50, m	C-10, C-12
12	75.2, CH	3.66, dd (11.5, 4.4)	C-13
OH		5.75, s	
13	42.8, qC		
14	54.1, CH	1.21, m	C-8, C-13, C-16
15	23.8, $\text{CH}_2$	2.25, dd (11.7, 7.1) 1.74, m	C-16
16	69.0, CH	5.50, brdd (10.3, 7.1)	C-17, C-18
17	158.5, qC		
18	139.2, qC		
19	33.2, $\text{CH}_3$	0.83, s	C-4
20	20.8, $\text{CH}_3$	0.80, s	C-4
21	17.2, $\text{CH}_3$	0.89, s	C-8
22	15.9, $\text{CH}_3$	0.83, s	C-1, C-10
23	16.5, $\text{CH}_3$	1.19, s	C-12, C-13, C-18
24	70.5, $\text{CH}_2$	4.76, dd (18.3, 1.3) 4.71, d (18.3)	C-17, C-18, C-25
25	174.9, qC		
26	170.5, qC		
27	21.3, $\text{CH}_3$	2.12, s	C-26

<sup>a</sup> Protons that correlated with carbons.

against *Mycobacterium tuberculosis* ( $\text{H}_37\text{Rv}$ ) (ATCC 27294), in BACTEC 12B medium using the Microplate Alamar Blue Assay (MABA).<sup>18</sup> Compound **1** displayed 63% inhibition at this concentration. Compounds **2**–**4** showed inhibitory activities of 40%, 16%, and 13%, respectively. A minimum inhibition of 90% against *M. tuberculosis* ( $\text{H}_37\text{Rv}$ ) is needed to progress compounds to an in vivo study.<sup>19</sup>

## Experimental Section

**General Experimental Procedures.** Optical rotations were measured on a Jasco-DIP-700 digital polarimeter using  $\text{CH}_2\text{Cl}_2$  at 20 °C at the sodium D line (589 nm). NMR spectra were determined on a Varian Unity 500 instrument (500 MHz for  $^1\text{H}$  and 125 MHz for  $^{13}\text{C}$  NMR). Homonuclear  $^1\text{H}$  connectivities were determined by using the 2D double-quantum filtered COSY technique. One-bond heteronuclear  $^1\text{H}$ – $^{13}\text{C}$  connectivities were determined by a 2D proton-detected HMQC experiment; two- and three-bond  $^1\text{H}$ – $^{13}\text{C}$  connectivities were determined by a 2D proton-detected HMBC experiment. HRFABMS were determined on a Finnigan MAT-312 spectrometer using 3-NBA/NaCl as matrix. Reversed-phase HPLC was conducted on a semipreparative  $\text{C}_{18}$  Ultracarb column (5  $\mu\text{m}$  ODS 30, 250  $\times$  10 mm, Phenomenex).

**Biological Material.** The sponge was collected by hand using scuba at a depth of 15 m off Safaga at the Egyptian Red Sea coast in December 2002. The sponge material was frozen immediately and kept frozen at –20 °C until workup. The sponge was identified as *Hyrtios erecta* (Keller) (family Thorectidae, order Dictyoceratida).<sup>2</sup> A voucher specimen has been deposited in the Natural History Museum, London, UK, and in the Red Sea invertebrates collection at the Faculty of Pharmacy, Suez Canal University, under the registration number DY-29.

**Extraction and Isolation.** Freshly collected specimens (800 g, wet wt) of the sponge were frozen immediately. The sponge was extracted with MeOH (3  $\times$  1.5 L) at room temperature. The combined extracts were concentrated under reduced pressure, dissolved in 500 mL of MeOH/ $\text{H}_2\text{O}$  (9:1), and extracted with hexane (4  $\times$  250 mL) to give 585 mg of hexane

residue. The remaining methanolic layer was diluted with  $\text{H}_2\text{O}$  to (3:2) MeOH– $\text{H}_2\text{O}$  and then extracted with  $\text{CH}_2\text{Cl}_2$  (4  $\times$  250 mL) to afford 431 mg of a  $\text{CH}_2\text{Cl}_2$  extract. The hexane residue was subjected to silica flash column chromatography using hexane/ $\text{CH}_2\text{Cl}_2$ /acetone. Fractions of 100 mL were collected and monitored by TLC. Similar fractions were pooled to give 10 further fractions. Of these, fraction 5 (110 mg) was purified on a semipreparative  $\text{C}_{18}$  reversed-phase HPLC column using 80% MeCN (2.0 mL/min at 220 nm) to give **1** (2.3 mg) and **2** (0.8 mg). In turn, fraction 6 (72 mg) was further purified on a semipreparative  $\text{C}_{18}$  reversed-phase HPLC column using 90% MeCN (2.0 mL/min at 220 nm) to afford **3** (1.9 mg), **4** (2.1 mg), and 21-acetoxydeoxyxylarinin  $\{[\alpha]_{\text{D}} +61.1$  (c 1.0,  $\text{CH}_2\text{Cl}_2$ ); HRFABMS  $m/z$  511.5041 (calcd for  $\text{C}_{29}\text{H}_{44}\text{O}_6\text{Na}$ ,  $[\text{M} + \text{Na}]^+$ , 511.5036) $\}$  (1.5 mg).

**Sesterstatin 7 (1):** amorphous solid;  $[\alpha]_{\text{D}} -19.5$  (c 1.15,  $\text{CH}_2\text{Cl}_2$ ); NMR data, see Table 1; HRFABMS  $m/z$  467.2769 (calcd for  $\text{C}_{27}\text{H}_{40}\text{O}_5\text{Na}$   $[\text{M} + \text{Na}]^+$ , 467.2773).

**16-*epi*-Sclarolbutenolide (2):** amorphous solid;  $[\alpha]_{\text{D}} -7.5$  (c 1.2,  $\text{CH}_2\text{Cl}_2$ ); HRFABMS  $m/z$  467.2769 (calcd for  $\text{C}_{27}\text{H}_{40}\text{O}_5\text{Na}$   $[\text{M} + \text{Na}]^+$ , 467.2773).

**25-Dehydroxy-12-*epi*-deacetylscalarin (3):** amorphous solid;  $[\alpha]_{\text{D}} -22.5$  (c 1.0,  $\text{CH}_2\text{Cl}_2$ ); HREIMS  $m/z$  409.2711 (calcd for  $\text{C}_{25}\text{H}_{38}\text{O}_3\text{Na}$ ,  $[\text{M} + \text{Na}]^+$ , 409.2719).

**3-Acetylsesterstatin 1 (4):** amorphous solid;  $[\alpha]_{\text{D}} +82.5$  (c 0.6,  $\text{CH}_2\text{Cl}_2$ ); HRFABMS  $m/z$  467.2769 (calcd for  $\text{C}_{27}\text{H}_{40}\text{O}_5\text{Na}$   $[\text{M} + \text{Na}]^+$ , 467.2773).

**Acknowledgment.** This research was supported by the U.S.-Egypt Science and Technology Joint Fund in cooperation with the USDA and Ministry of Scientific Research in Egypt under Project #BIO5-002-021. Our thanks go to the members of the Egyptian Environmental Affairs Agency (EEAA) and the Red Sea Protectorate of Egypt for collection permission. We acknowledge V. Brecht for the NMR and MS data. Antimicrobial testing was conducted by the Tuberculosis Antimicrobial Acquisition and Coordinating Facility (TAACF) through a research and development contract #1540 between D.T.A.Y. and the U.S. National Institute of Allergy and Infectious Diseases, NIH.

## References and Notes

- Blunt, J. W.; Copp, B. R.; Munro, M. H. G.; Northcote, P. T.; Prinsep, M. R. *Nat. Prod. Rep.* **2005**, *22*, 15–61, and previous reviews in this series.
- Youssef, D. T. A.; Yamaki, R. K.; Kelly, M.; Scheuer, P. J. *J. Nat. Prod.* **2002**, *65*, 2–6.
- (a) Pettit, G. R.; Cichacz, Z. A.; Tan, R.; Hoard, M. S.; Melody, N.; Pettit, R. K. *J. Nat. Prod.* **1998**, *61*, 13–16. (b) Pettit, G. R.; Cichacz, Z. A.; Tan, R.; Herald, D. L.; Melody, N.; Hoard, M. S.; Doubek, D. L.; Hooper, J. N. A. *Collect. Czech. Chem. Commun.* **1998**, *63*, 1671–1677. (c) Pettit, G. R.; Tan, R.; Melody, N.; Cichacz, Z. A.; Herald, D. L.; Hoard, M. S.; Pettit, R. K.; Chapuis, J.-C. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 2093–2098. (d) Pettit, G. R.; Tan, R.; Cichacz, Z. A. *J. Nat. Prod.* **2005**, *68*, 1253–1255.
- (a) Ryu, G.; Matsunaga, S.; Fusetani, N. *J. Nat. Prod.* **1996**, *59*, 515–517. (b) Nasu, S. S.; Yeung, B. K. S.; Hamann, M. T.; Scheuer, P. J.; Kelly-Borges, M.; Goins, K. J. *Org. Chem.* **1995**, *60*, 7290–7292. (c) Williams, D. E.; Tahir, A.; Andersen, R. J. *J. Nat. Prod.* **1999**, *62*, 653–654. (d) Salmoun, M.; Devijver, C.; Daloze, D.; Braekman, J. C.; Gomez, R.; de Kluijver, M.; van Soest, R. W. M. *J. Nat. Prod.* **2000**, *63*, 452–456.
- Hochlowski, J. E.; Faulkner, D. J.; Bass, L. S.; Clardy, J. *J. Org. Chem.* **1983**, *48*, 1738–1740.
- Berquist, P. R.; Cambie, R. C.; Kernan, M. R. *Biochem. Syst. Ecol.* **1990**, *18*, 349–357.
- Walker, R. P.; Thompson, J. E.; Faulkner, D. J. *J. Org. Chem.* **1980**, *45*, 4976–4979.
- Braekman, J. C.; Daloze, D.; Kaisin, M.; Moussiaux, B. *Tetrahedron* **1985**, *41*, 4603–4614.
- Kikuchi, H.; Tsukitani, Y.; Shimizu, I.; Kobayashi, M.; Kitagawa, I. *Chem. Pharm. Bull.* **1983**, *31*, 552–556.
- Crews, P.; Bescansa, P. *J. Nat. Prod.* **1986**, *49*, 1041–1052.
- Nakagawa, M.; Hamamoto, Y.; Ishihama, M.; Hamasaki, S.; Endo, M. *Tetrahedron Lett.* **1987**, *28*, 431–434.
- Doi, Y.; Shigemori, H.; Ishibashi, M.; Mizobe, F.; Kawashima, A.; Nakaike, S.; Kobayashi, J. *Chem. Pharm. Bull.* **1993**, *41*, 2190–2191.
- (a) Kobayashi, M.; Aoki, S.; Sakai, H.; Kawazoe, N.; Kihara, N.; Sasaki, T.; Kitagawa, I. *Tetrahedron Lett.* **1993**, *34*, 2795–2798. (b) Kobayashi, M.; Aoki, S.; Sakai, H.; Hihara, N.; Sasaki, T.; Kitagawa, I. *Chem. Pharm. Bull.* **1993**, *41*, 989–991. (c) Pettit, G. R.; Cichacz,

- Z. A.; Gao, F.; Herald, C. L.; Boyd, M. R.; Schmidt, J. M.; Hooper, J. N. A. *J. Org. Chem.* **1993**, *58*, 1302–1304.
- (14) (a) Youssef, D. T. A. *J. Nat. Prod.* **2005**, *68*, 1416–1419. (b) Kobayashi, J.; Murayama, T.; Ishibashi, M.; Kosuge, S.; Takamatsu, M.; Ohizumi, Y.; Kobayashi, H.; Ohta, T.; Nozoe, S.; Sasaki, T. *Tetrahedron* **1990**, *46*, 7699–7702. (c) Bourguet-Kondracki, M. L.; Martin, M. T.; Guyot, M. *Tetrahedron Lett.* **1996**, *37*, 3457–3460. (d) Salmoun, M.; Devijver, C.; Daloze, D.; Braekman, J.-C.; van Soest, R. W. M. *J. Nat. Prod.* **2002**, *65*, 1173–1176.
- (15) Rueda, A.; Zubia, E.; Ortega, M. J.; Carballo, J. L.; Salva, J. *J. Nat. Prod.* **1998**, *61*, 258–261.
- (16) Venkateswarlu, Y.; Bibani, M. A. F.; Rao, T. P. *Ind. J. Chem.* **1995**, *34B*, 563–564.
- (17) Miyamoto, T.; Sakamoto, K.; Amano, H.; Arakawa, Y.; Nagarekawa, Y.; Komori, T.; Higuchi, R.; Sasaki, T. *Tetrahedron* **1999**, *55*, 9133–9142.
- (18) Collins, L.; Franzblau, S. G. *Antimicrob. Agents Chemother.* **1997**, *41*, 1004–1009.
- (19) <http://www.taacf.org/Process-text.htm#nhdp-text>.

NP0502645