

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/6286354>

# Keeping an eye on I1:I1 Dynein as a model for flagellar dynein assembly and regulation

ARTICLE *in* CELL MOTILITY AND THE CYTOSKELETON · AUGUST 2007

Impact Factor: 4.19 · DOI: 10.1002/cm.20211 · Source: PubMed

---

CITATIONS

59

---

READS

29

3 AUTHORS, INCLUDING:



[Maureen Wirschell](#)

University of Mississippi Medical Center

22 PUBLICATIONS 462 CITATIONS

[SEE PROFILE](#)



[Triscia Hendrickson](#)

Morehouse College

5 PUBLICATIONS 175 CITATIONS

[SEE PROFILE](#)

## **View Article**

# Keeping an Eye on I1: I1 Dynein as a Model for Flagellar Dynein Assembly and Regulation

Maureen Wirschell,<sup>1\*</sup> Triscia Hendrickson,<sup>2</sup> and Winfield S. Sale<sup>1</sup>

<sup>1</sup>*Department of Cell Biology, Emory University School of Medicine, Atlanta, Georgia*

<sup>2</sup>*Biology Department, Morehouse College, Atlanta, Georgia*

Among the major challenges in understanding ciliary and flagellar motility is to determine how the dynein motors are assembled and localized and how dynein-driven outer doublet microtubule sliding is controlled. Diverse studies, particularly in *Chlamydomonas*, have determined that the inner arm dynein I1 is targeted to a unique structural position and is critical for regulating the microtubule sliding required for normal ciliary/flagellar bending. As described in this review, I1 dynein offers additional opportunities to determine the principles of assembly and targeting of dyneins to cellular locations and for studying the mechanisms that regulate dynein activity and control of motility by phosphorylation. *Cell Motil. Cytoskeleton* 64:569–579, 2007. © 2007 Wiley-Liss, Inc.

**Key words:** I1 dynein; f-dynein; cilia; flagella; axoneme

## **INTRODUCTION AND OVERVIEW**

Genetic, proteomic, and computational analyses have recently shed much light on the importance of cilia and flagella in human health [El Zein et al., 2003; Avidor-Reiss et al., 2004; Li et al., 2004; Pazour, 2004; Snell et al., 2004; Pazour et al., 2005]. Failure to assemble these structures or failure in their motility can lead to severe consequences in development or adult functions in many organs [Snell et al., 2004; Davenport and Yoder, 2005; Pan et al., 2005; Quarmby and Parker, 2005; Davis et al., 2006; Christensen et al., 2007]. Therefore, understanding how cilia and flagella assemble and the processes that drive motility—particularly the dynein motors—are of great interest. Here, we focus on a conserved dynein motor, inner arm dynein I1, and how studies, primarily in *Chlamydomonas*, have revealed new features of dynein assembly and regulation of axonemal motility by phosphorylation.

*Chlamydomonas* is a unicellular, biflagellate, green alga that is an excellent model system for the discovery of conserved genes required for flagellar assembly and

function [Silflow and Lefebvre, 2001] and in particular for study of the flagellar dyneins—the microtubule motors that drive flagellar motility [DiBella and King, 2001; Kamiya, 2002; Oiwa and Sakakibara, 2005]. Structural and biochemical studies of wild-type and mutant *Chlamydomonas* have revealed the axoneme bears at least seven different inner arm dyneins that differ in composi-

Contract grant sponsor: NIH; Contract grant numbers: GM075446, GM051173, GM067078; Contract grant sponsor: March of Dimes; Contract grant number: FY04-115.

\*Correspondence to: Maureen Wirschell, Department of Cell Biology, Emory University School of Medicine, Whitehead Biomedical Research Building, Suite 465, 615 Michael St., Atlanta, GA 30322, USA. E-mail: wirsch@cellbio.emory.edu

Received 28 March 2007; Accepted 17 April 2007

Published online 4 June 2007 in Wiley InterScience (www.interscience.wiley.com).  
DOI: 10.1002/cm.20211

tion and structural location [DiBella and King, 2001]. For example, in the axoneme, I1-dynein is a trilobed structure originally identified as the “triad” [Goodenough and Heuser, 1985a]. Mutations affecting certain I1 subunits often result in a complete failure in I1-dynein assembly [*ida1*, *ida2*, *ida7*—See Table I for nomenclature, Kamiya et al., 1991] leaving a gap in the I1 position that repeats every 96-nm and does not affect the assembly of the other flagellar dynein isoforms [Piperno et al., 1990; Kamiya et al., 1991; Porter et al., 1992; Smith and Sale, 1992b; Perrone et al., 1998]. Thus I1-dynein must be targeted to a unique position in the axoneme by specialized proteins or domains that repeat every 96-nm.

Phenotypic analysis has revealed that the inner arm dyneins are responsible for determining the size and shape of the flagellar bend, parameters that define waveform [Brokaw and Kamiya, 1987; Kamiya et al., 1991; Kamiya, 1995]. In some cases, mutations that affect I1-dynein assembly suppress paralysis in a central pair mutant [Porter et al., 1992]. Also, I1-dynein assembly mutants exhibit defects in phototaxis—a behavioral response to a light stimulus that requires precise control of flagellar bending indicating I1 is important, although not absolutely required, in this process [King and Dutcher, 1997; Okita et al., 2005]. These phenotypic data demonstrate that I1 dynein is a critical component of a complex system, involving multiple axonemal dyneins, which controls axonemal bending. Extensive in vitro functional and pharmacological data also indicate I1 dynein is required for control of axonemal sliding [Habermacher and Sale, 1997; King and Dutcher, 1997; Yang and Sale, 2000; Hendrickson et al., 2004], and that the mechanism of regulation involves reversible phosphorylation of the I1 subunit IC138 in a process that also may be controlled by calcium [Hennessey et al., 2002; Smith, 2002a; Dymek and Smith, 2006].

Phylogenetic analyses have revealed that I1-dynein proteins are highly conserved, thus the study of I1 in *Chlamydomonas* is relevant to a wide range of organisms. For example, not only is I1-dynein activity regulated by phosphorylation in *Chlamydomonas*, recent data suggests that I1 activity plays a role in regulating motility in *Tetrahymena thermophila* [Hennessey et al., 2002; Deckman and Pennock, 2004].

In this commentary, we discuss mechanisms for assembly and transport of I1, anchoring I1-dynein to the doublet microtubule, and localizing I1-dynein within the 96-nm repeat. This review also incorporates very recent and exciting unpublished work that has been presented in Abstract form at the 2006 American Society for Cell Biology annual meeting. We describe a model for how localized control of I1-dynein phosphorylation operates to locally control microtubule sliding and adjust axonemal waveform. In particular, we summarize a model in

which I1 dynein locally inhibits microtubule sliding driven by other dynein isoforms.

## ORGANIZATION OF THE INNER ARMS: THE 96-NM REPEAT AND SUBUNIT COMPOSITION OF I1 DYNEIN

Compared to the outer dynein arms, the inner arm dyneins are heterogeneous with respect to their organization [Porter and Sale, 2000; DiBella and King, 2001; Kamiya, 2002]. Analysis of mutant phenotypes, biochemical identification of the inner arm dynein subtypes, and eloquent ultrastructural studies have helped to define the composition and arrangement of the complex inner dynein arm system, particularly the I1-dynein. The complexity of composition and structure has led to different nomenclatures, but on the basis of heavy chain (HC) number, there appear to be two types of inner arm dyneins. I1-dynein, also called f-dynein, contains two HCs, whereas the inner arm dyneins containing one HC are designated a, b, c, d, e, and g, and were initially thought to be organized in structures referred to as I2 and I3 [Piperno et al., 1990; Kagami and Kamiya, 1992, 1995].

The inner arm dyneins are organized around a regular repeating pattern with a 96-nm repeat length (Fig. 1). This pattern has been elucidated in exquisite detail by rapid freeze-deep etch microscopy [Goodenough and Heuser, 1985b], analysis of inner arm mutants [Piperno et al., 1990], image analysis of thin sections [Mastromarino et al., 1992; Porter et al., 1992; Perrone et al., 1998], and most recently by cryoEM tomography of axonemes [Nicastro et al., 2005, 2006]. The I1 dynein is located in the proximal position of the 96-nm repeat, in close proximity to radial spoke S1, and is found along the entire length of the axoneme (Fig. 1). Although it is presumed that I1 dynein is found on all nine outer doublets, this has not yet been determined by direct localization at the ultrastructural level. Based on analysis in a number of organisms, the organization of the 96-nm repeat is a highly conserved feature of the axoneme [Goodenough and Heuser, 1985b; Woolley, 1997; Nicastro et al., 2005, 2006]. Additional features of the 96-nm repeat module are the single headed inner arm dyneins, two radial spokes (S1 and S2), and the dynein regulatory complex (DRC)-structures that are critical for regulation of dynein function [Gardner et al., 1994; Piperno, 1995; Smith and Yang, 2004], as well as novel structures connecting I1 dynein and the DRC to specific outer dynein arms within the 96-nm module [Nicastro et al., 2006]. The novel structures linking I1-dynein to the outer arm may be composed of the Oda7 protein [Freshour et al., 2006; Nicastro et al., 2006].

TABLE I. Inner Dynein Arm Components and Mutations that Affect II-Dynein Assembly or Function<sup>a</sup>

| Subunit   | <i>M<sub>r</sub></i> | Mutant <sup>a,b</sup>                    | Mutant phenotype  | Predicted function  | References  |
|-----------|----------------------|--|---|---|---|
| 1αHC      | 523                  | <i>ida1</i> ( <i>pf9</i> , <i>pf30</i> ) | Lacks II, slow-swimming   | Motor activity<br>Altered waveform, phototaxis  | Kamiya et al., 1991; Porter et al., 1992; Okita et al., 2005; Piperno et al., 1990; Brokaw and Kamiya, 1987 |
| 1βHC      | 511                  | <i>G41a</i><br><i>ida2</i>               | Truncated motor domain<br>Lacks II, slow-swimming                                 | Required for II assembly<br>Motor activity  | Myster et al., 1999<br>Kamiya et al., 1991  |
| IC140     | 140                  | <i>D11</i><br><i>ida7</i>                | Altered waveform, phototaxis<br>Truncated motor domain<br>Lacks II, slow-swimming | Required for II assembly<br>Required for II assembly  | Perrone et al., 2000<br>Yang and Sale, 1998; Perrone et al., 1998   |
| IC138     | 138                  | 5A<br><i>bop5-1</i>                      | Altered waveform, phototaxis<br>Truncated IC140<br>Truncated IC138, II lacks LC7b | Not required for II assembly<br>Not required for II assembly  | Perrone et al., 1998<br>Hendrickson et al., 2004; Dutcher et al., 1988                                      |
| ?<br>IC97 | ?<br>97 <sup>c</sup> | <i>mia1</i> , <i>mia2</i><br>–           | Slow-swimming<br>Hyperphosphorylated IC138  | Not required for II assembly<br>Not required for II assembly  | King and Dutcher, 1997<br>Wirschell et al., 2006  |
| TcTex1    | 13                   | –  | –   | Cargo binding   | Harrison et al., 1998   |
| TcTex2b   | 13.7                 | <i>pf16-D2::PF16</i>                     | Slow-swimming   | Not required for II assembly<br>May modulate motor function   | DiBella et al., 2004b   |
| LC8       | 10                   | <i>fla14</i>                             | Short immotile flagella,<br>Altered waveform, phototaxis                          | Stability of II<br>Stable assembly of II found in outer arms, inner arms, radial spokes, and the retrograde IFT motor | Pazour et al., 1998   |
| LC7a      | 14                   | <i>oda15</i>                             | Partial outer arm assembly defect   | Stable association of LC7b in II  | DiBella et al., 2004a   |
| LC7b      | 11                   | –  | Missing in <i>bop5-1</i> mutant   | Not required for II assembly<br>Interacts with IC138  | Hendrickson et al., 2004<br>Hendrickson et al., 2004; DiBella et al., 2004a                                 |
| ?         | ?                    | <i>ida3</i>                              | Lacks II, slow-swimming<br>Altered waveform, phototaxis                           | May modulate II function  | Kamiya et al., 1991   |

<sup>a</sup>Mutations affecting the inner dynein arm system generally are referred to as *ida* mutants (Kamiya et al., 1991). Exceptions to this system of nomenclature include mutants designated as *pf* (paralyzed flagella), *bop* (bypass of paralysis), or *mia* (modifiers of inner arms); named either prior to the establishment of the *ida* nomenclature or prior to the recognition that the mutation affected inner arm dyneins.

<sup>b</sup>Individual alleles at each locus are not detailed here.

<sup>c</sup>In some reports using different SDS-PAGE markers, IC97 is reported to migrate at 110 kDa and is cited as IC110 (Dibella and King, 2001; King and Dutcher, 1997; Porter et al., 1992).




Fig. 1. Schematic of the 96-nm axonemal dynein repeat. The arrangement of the inner dynein arms, outer dynein arms, radial spokes, and DRC on the A-tubule is shown [adapted from Porter and Sale, 2000; Nicastro et al., 2006]. These structures repeat every 96-nm along the length of the axoneme (the outer dynein arms repeat every 24-nm). The proximal end (closest to the cell body) faces left. I1-dynein is the trilobed inner arm just proximal to radial spoke S1 in each repeat (green); it is followed by a series of single headed inner arms (labeled 2–6). PKA is thought to be localized at the base of the radial spokes [Gaillard et al., 2006]; CK1 and PP2A are predicted to

be anchored near I1 dynein; a fraction of PP1 located on the outer doublets. The inset shows a diagram of the I1-dynein and its individual subunits [Harrison et al., 1998; Perrone et al., 1998; Yang and Sale, 1998; Porter and Sale, 2000; DiBella et al., 2004a,b; Hendrickson et al., 2004]. I1-dynein is composed of two motor subunits (1α- and 1β-HC), three intermediate chains (IC140, IC138, and IC97), and five LC subunits (Tctex1, Tctex2b, LC8, LC7a, and LC7b). IC138 is a phosphoprotein in I1 and is one of the key regulatory subunits critical for I1-dynein activity.

I1-deficient axonemes reveal that I1 is localized to a unique position in the 96-nm repeat and that failure in I1 assembly does not affect assembly of other axonemal structures [Piperno et al., 1990; Kamiya et al., 1991; Porter et al., 1992; Smith and Sale, 1992b; Nicastro et al., 2006]. This result is important in several ways. First, the other axonemal dyneins assemble normally indicating mutations in I1 assembly do not disrupt assembly of other dynein isoforms. Second, the results also imply that I1-dynein is localized in the axoneme by a docking mechanism that resides near the S1 radial spoke, at the proximal end of the 96-nm repeat (Fig. 1). Consistent with this prediction, *in vitro* reconstitution of I1-deficient axonemes results in a restoration of functional I1 structures to the original position in the 96-nm repeat [Smith and Sale, 1992b; Yamamoto et al., 2006]. However, to date, no docking complex has been identified.

I1 is the only inner dynein arm with two HCs, designated 1α and 1β, [Piperno et al., 1990]. I1 dynein forms a two headed structure as revealed by electron microscopy of isolated I1 particles [Goodenough et al., 1987; Smith and Sale, 1991; Sakakibara et al., 2006].

Along with the HCs, are three intermediate chains (IC140, IC138, and IC97), and five known light chains (LC) (Tctex1, Tctex2b, LC8, LC7b, and LC7a) [Porter et al., 1992; Harrison et al., 1998; Perrone et al., 1998; Yang and Sale, 1998; DiBella et al., 2001, 2004a,b; Hendrickson et al., 2004; Wirschell et al., 2006] (Fig. 1 inset and Table I). The HCs form stem domains which connect to an IC/LC complex that forms the cargo-binding domain of the dynein motor as well as globular head domains that possess the motor activity. For dyneins in general, the intermediate and LCs play important roles in motor assembly and regulation [Perrone et al., 1998; Yang and Sale, 1998; DiBella et al., 2001, 2004a,b]. Several of the LCs are not unique to I1, notably LC8, LC7a, LC7b, and Tctex1. Rather, they are also LCs in other dyneins and additional enzymes and structures [Pazour et al., 1998; DiBella et al., 2004a,b]. These LCs may have common functions in stable assembly of dynein structures (LC8), cargo binding (Tctex1-rhodopsin), or regulation (LC7b) [King et al., 1996; Harrison et al., 1998; Pazour et al., 1998; Tai et al., 1999; Hendrickson et al., 2004].

## I1-DYNEIN MUTANTS REVEAL NEW FEATURES OF ASSEMBLY AND DOCKING

Mutations that disrupt specific I1 subunits or specific domains of those subunits often result in assembly of incomplete, or partial, I1 dynein complexes (Table I). For example, mutant strains containing truncated HC genes, lacking the motor domains, still assemble an I1 complex [Myster et al., 1999; Perrone et al., 2000]. These results are consistent with previous studies indicating that N-terminal fragments between 140- and 160-kDa are capable of complex assembly [Sakakibara et al., 1993; Koonce and Knecht, 1998; Iyadurai et al., 1999]. Importantly, electron microscopy of these HC truncations revealed the location of the globular motor domains of the  $1\alpha$  and  $1\beta$ -HCs in the axoneme [Myster et al., 1999; Perrone et al., 2000]. They also indirectly reveal the location of the IC/LC complex at the base of I1 dynein where it is anchored to the A-tubule (Fig. 1). The assembly of I1 dynein complexes that lack one of the HC motor domains provides an opportunity to dissect the individual roles of the  $1\alpha$ - and  $1\beta$ -HCs [Fox et al., 2006; Toba et al., 2006].

The ICs typically contain WD repeats and are integral structural components of dynein motors. This indeed seems to be the case for the IC140 subunit (see below). Like the ICs of the outer arm, IC140 is required for assembly of the I1 dynein complex and thus is implicated in anchoring of I1 to the axoneme [Perrone et al., 1998; Yang and Sale, 1998]. In support of this, the C-terminus of IC140 interacts with the axoneme directly [Yang and Sale, 1998] and thus may interact with components that localize I1 in the 96-nm axonemal repeat structure. However, while IC140 is required for I1 assembly in the axoneme, the N-terminal region is not critical for its assembly function (Table I-strain 5A, [Perrone et al., 1998]). Alternatively, in the absence of IC140, I1-dynein components in the cytoplasm may be unstable and degraded, thus contributing to the failure of I1-dynein to assemble in the axoneme.

The *bop5-1* mutant expresses a truncated IC138 protein that lacks the C-terminal WD repeat domain (Table I). All I1 subunits assemble in this mutant, except for LC7b indicating that LC7b and IC138 interact [DiBella et al., 2004a; Hendrickson et al., 2004]. A second IC138 mutation indicates that, unlike IC140 and the outer arm ICs, IC138 is not required for assembly of the entire I1-dynein complex, but may be required for assembly of the IC97 and LC7b subunits [Hendrickson et al., 2004; Bower et al., 2006]. These studies indicate that the primary role of IC138 is not in assembly and is consistent with the known regulatory role of IC138 in control of microtubule sliding (see below). The results also reveal that IC97 is not required for assembly of I1, but they do

not reveal the exact function of IC97 within I1. This mutation further indicates that IC140 along with the N-terminal HC domains are principally responsible for targeting and anchoring I1 to its specific location in the 96-nm repeat.

Based on crosslinking studies, IC138 and IC140 interact directly with axonemal proteins [Yang and Sale, 1998; Hendrickson et al., 2004] and at least some of these interactions may reflect interactions with tubulin (Wirschell, Hendrickson, and Sale, unpublished observations). IC97 interacts directly with both  $\alpha$ - and  $\beta$ -tubulin subunits [Wirschell et al., 2006]. Given that all three ICs may interact directly with tubulin, anchoring of I1 dynein to the doublet microtubule involves multiple interactions. However, IC-tubulin interactions cannot explain the exact placement of I1 within the 96-nm repeat. One idea is that I1 interacts with an intrinsic component of the outer doublet that establishes the 96-nm repeat structure that governs placement of the inner dynein arms, radial spokes, and DRC. Proteins that might be part of such a complex could include the tektins [Norlander et al., 1996; Yanagisawa and Kamiya, 2004; Setter et al., 2006]. Consistent with this idea, tektins may act as a molecular ruler that defines the axonemal 96-nm repeat [Setter et al., 2006] and in *Chlamydomonas*, defects in tektin assembly are associated with defective inner dynein arm assembly [Yanagisawa and Kamiya, 2004].

Structural analyses of axonemes from dynein mutants have demonstrated that each dynein isoform is targeted to its own unique position along the length and around the circumference of the A-microtubule of most outer doublets. Anchoring of the outer dynein arm is mediated in part through additional complexes called outer dynein arm-docking complexes (ODA-DC and the Oda5 complex) [Takada and Kamiya, 1994; Wakabayashi et al., 2001; Casey et al., 2003; Wirschell et al., 2004]. Moreover, additional mutations have revealed the outer dynein arm is transported into the axoneme specifically through an interaction with the intraflagellar transport component IFT46 and Oda16p—a novel non-dynein transport factor required for outer dynein arm assembly [Ahmed and Mitchell, 2005; Hou et al., 2007]. Evidence suggests that single-headed inner arms are also transported into the flagellum by IFT [Piperno et al., 1996]. Given that other dyneins are transported by IFT, it is likely that I1 is as well. However, to date, similar IFT transport mechanisms have not been defined for I1 dynein. Predictably, recovery of additional mutants that fail to assemble I1 will reveal new genes required for assembly, transport, targeting, or docking of I1 dynein. The *ida3* mutant may be an example. The *ida3* mutation causes a complete I1 assembly defect, but the *IDA3* gene product remains unknown (Table I).



# FUNCTIONAL ROLE OF I1 IN REGULATION OF FLAGELLAR BENDING: A MODEL FOR LOCAL REGULATION OF MICROTUBULE SLIDING BY PHOSPHORYLATION

Among the major challenges in understanding the mechanism of ciliary motility is to determine how dynein-driven microtubule sliding is regulated. Based on models, oscillatory bending and bend propagation require mechanical feedback mechanisms that control the timing and location of active sliding [Brokaw, 1985, 1994; Lindemann, 2002]. However, the experimental evidence for the identity and nature of such a mechanical feedback mechanism is lacking.

In contrast, direct experimental evidence, through genetic and in vitro functional studies using *Chlamydomonas* axonemes, has revealed I1-dynein is important for control of microtubule sliding and is likely designed for modulating the size and shape of the axonemal bend [Brokaw and Kamiya, 1987]. We summarize data indicating I1-dynein is important for control of flagellar waveform and that the mechanism involves regulation of phosphorylation of the I1-dynein subunit IC138 and control of microtubule sliding [Habermacher and Sale, 1997; King and Dutcher, 1997; Yang and Sale, 2000; Hendrickson et al., 2004]. We also discuss a model in which localized inhibition of microtubule sliding [Smith, 2002b] is mediated by I1-dynein and IC138 phosphorylation and is required for normal control of bending behavior.

Mutations in I1-dynein subunits or mutations that alter I1 phosphorylation result in failure of control of normal axonemal waveform and phototaxis in *Chlamydomonas* [Brokaw and Kamiya, 1987; King and Dutcher, 1997; Okita et al., 2005]. For example, the *mia* mutants were isolated in a screen for phototaxis mutations and the mutant axonemes exhibit hyperphosphorylated IC138 [King and Dutcher, 1997]. Based on microtubule sliding assays and pharmacological analysis, dynein-driven sliding is inhibited in *mia* mutant axonemes. However, addition of kinase inhibitors rescues the microtubule sliding defects. The data revealed a correlation between IC138 phosphorylation and regulation of dynein; phosphorylated IC138 inhibited microtubule sliding and dephosphorylation rescued microtubule sliding (Fig. 2). Okita et al. have further examined the role of I1 in phototaxis and suggest that I1 may be involved in one of many pathways required for generating phototactic behavior [Okita et al., 2005]. Additionally, mutations in I1 subunits can suppress paralysis in a central pair mutant, indicating a functional regulatory interaction between I1-dynein activity and the central pair-radial spoke structures for control of microtubule sliding and axonemal motility [Porter et al., 1992]. Thus, I1




Fig. 2. Regulation of IC138 phosphorylation. IC138 is a critical component of I1 involved in the radial spoke-central pair mechanism for regulating I1 dynein activity [adapted from Habermacher and Sale, 1997; Porter and Sale, 2000]. This regulatory process involves changes in IC138 phosphorylation such that in the phosphorylated state, I1 is inactive; when IC138 is dephosphorylated, I1-dynein activity is restored. The critical axonemal kinases and phosphatases involved in this pathway are indicated [Porter and Sale, 2000; Smith and Yang, 2004].

dynein appears to regulate axonemal motility through a mechanism that controls microtubule sliding.

Consistent with the idea that I1-dynein plays a role in control, in vitro functional assays, using isolated axonemes and microtubule sliding, have also revealed that I1-dynein plays a central role in regulation of microtubule sliding and that the mechanism depends upon changes in phosphorylation of IC138. Figure 2 illustrates the role of CK1 and PKA in phosphorylation of IC138. Interestingly, a mating induced elevation in intraflagellar cAMP [Hasegawa et al., 1987; Pasquale and Goodenough, 1987] levels is presumably responsible for quiescence of flagella during mating (observations by Ursula Goodenough, personal communication). An interesting and testable idea is that the increased intraflagellar cAMP levels affect IC138 phosphorylation and thus globally inhibits microtubule sliding leading to cessation or slowing of flagellar motility during the mating response. Moreover, dynein-driven microtubule sliding in paralyzed axonemes, lacking either the radial spokes or the central pair, is inhibited presumably throughout the axoneme [Smith and Sale, 1992a]. The inhibition of microtubule sliding in these paralyzed mutant axonemes can be rescued by addition of kinase inhibitors indicating abnormal phosphorylation in the axoneme inhibits microtubule sliding [Howard et al., 1994; Habermacher and Sale, 1995, 1997; Yang and Sale, 2000; Smith, 2002b]. The results are consistent with the model shown in Fig. 2 indicating that inhibition of dynein in the paralyzed mutants is due, in part, to phosphorylation of IC138 and that dephosphorylation rescues microtubule sliding [Habermacher and Sale, 1997; King and Dutcher, 1997; Yang and Sale, 2000; Smith, 2002b; Hendrickson et al., 2004]. However, the in vitro experiments were performed in a paralyzed flagella



Fig. 3. A model for localized regulation of I1 and axonemal bending. The axoneme is shown in cross section illustrating the 9 + 2 arrangement of the axoneme and the asymmetric central pair apparatus, the radial spokes, and the dynein arms [Smith, 2007]. The black line indicates the plane of bending [Mitchell, 2003b; Mitchell and Yokoyama, 2003]. The current model for regulation of I1-dynein is that chemical

and/or mechanical signals from the central pair are transmitted through the radial spokes to affect IC138 phosphorylation on specific outer doublets (e.g., outer doublets  $N$ ,  $N + 1$ ,  $N - 1$ ). Thus, phosphorylation of IC138 in an asymmetrical manner is predicted to result in local inhibition of sliding. This in turn would alter the form of the bend.

mutants, lacking either the central pair or radial spokes, in which axonemal kinase activity is misregulated, revealing a predicted global change in dynein activity [Gaillard et al., 2001, 2006; Gokhale et al., 2006]. The same in vitro measurements of microtubule sliding in wild-type axonemes have not been informative, most likely because the axonemal kinases are tightly and locally controlled. Thus, a major challenge is to determine how I1-dynein operates in wild-type axonemes for control of movement.

One idea, illustrated in Fig. 3 is that in wild-type axonemes, I1-dynein is locally and asymmetrically regulated in response to chemical or mechanical signals that impinge on the central pair apparatus. In this model, signals originating from the central pair, an inherently asymmetric structure, are directed to specific doublet microtubules through the radial spokes [Smith and Yang, 2004; Smith, 2007]. The idea is that an asymmetric signal, for example from one of the central pair microtubules, is directed at doublet “ $N$ ” (Fig. 3) and locally induces phosphorylation of IC138 on doublet “ $N$ ”. In contrast, signals from the other central pair microtubule would have a different affect on the phosphorylation of IC138 on other outer doublet microtubules. Thus, following local changes in phosphorylation, sliding between doublets “ $N$ ” and “ $N + 1$ ” will be different relative to sliding between other doublets, therefore altering bending. As discussed before [Yang et al., 2004], the input for the asymmetric central pair-radial spoke signal could be a second messenger, such as calcium impinging upon the central pair [Wargo et al., 2005; Dymek and Smith, 2006], and/or a mechanical interaction with the radial spoke [Warner and Satir, 1974; Smith and Yang,

2004; Yang et al., 2004]. The output signal, presumably mediated by a radial spoke on one doublet microtubule, would be the activation of axonemal kinases located near I1-dynein [Yang et al., 2004; Gaillard et al., 2006].

Consistent with these ideas, the central pair apparatus is asymmetric in structure and composition [Mitchell and Sale, 1999; Mitchell and Yokoyama, 2003; Mitchell, 2003a; Wargo and Smith, 2003; Smith and Yang, 2004; Lehtreck and Witman, 2007] and the position of the central pair correlates with sites of active sliding [Wargo and Smith, 2003; Wargo et al., 2004] and the position of doublets relative to the bend [Mitchell, 2003a; Smith and Yang, 2004]. The results have also revealed a functional role for kinases (PKA, CK1) and phosphatases (PP1, PP2A) that are physically localized in the axoneme [Habermacher and Sale, 1996, 1997; King and Dutcher, 1997; Porter and Sale, 2000; Yang et al., 2000; Gaillard et al., 2001, 2006; Gokhale et al., 2006]. Much effort is underway to identify molecular mechanisms for targeting otherwise ubiquitous signaling molecules to precise locations in the axoneme [Gaillard et al., 2001, 2006]. Further tests of these ideas will require new assays linking radial spoke function to the direct control of axonemal kinases located on the outer doublet microtubule.

Related challenges include determining the mechanism for how IC138 can alter microtubule sliding. The implication of the model is the I1-dynein locally operates to inhibit microtubule sliding [Smith, 2002b]. Consistent with this idea, isolated I1-dyneins generate relatively slow microtubule translocation indicating that I1 dynein is unlike the outer arm dyneins or other inner arm dyneins [Smith and Sale, 1991; Kagami and Kamiya, 1992;



Sakakibara and Nakayama, 1998; Sakakibara et al., 1999; Yagi et al., 2005]. Although it is possible that the I1-dynein is not well suited to such in vitro assays that involve absorption to inert surfaces, recent advances in the K. Oiwa lab (Kobe Advanced ICT Research Center, Kobe, Japan) have confirmed that I1-dynein generates only slow microtubule translocation compared to other dyneins [Sakakibara et al., 2006; Toba et al., 2006].

When IC138 is phosphorylated, microtubule sliding is inhibited in the axoneme [Habermacher and Sale, 1997; King and Dutcher, 1997; Yang and Sale, 2000; Hendrickson et al., 2004]. One hypothesis is that phosphorylation of IC138 inhibits I1-dynein motor activity. However, using purified I1-dynein and in vitro motility assays, it has been determined that motor activity of isolated I1 dynein is not altered by IC138 phosphorylation, yet in the axoneme microtubule sliding is inhibited by the same changes in IC138 [Sakakibara et al., 2006]. Thus further progress will require measurements of single I1 motor characteristics including assessment of processivity and duty cycle as has been done for other inner arm dyneins and further structural analyses of I1 [Sakakibara et al., 2006, 1999]. Additionally, our collaborative studies with M. Porter (University of Minnesota), Toba, Oiwa, Sakakibara, and colleagues (Kobe Advanced ICT Research Center, Kobe, Japan) are addressing models of I1-dynein mechanism using single motor analysis and dyneins isolated from wild-type and I1-mutant axonemes [Myster et al., 1999; Perrone et al., 2000; Fox et al., 2006; Toba et al., 2006]. The goal is to further test the hypothesis that I1 dynein acts as a “brake” to locally inhibit microtubule sliding driven by other dynein isoforms and thus control axonemal waveform.

## ACKNOWLEDGMENTS

We are grateful to Dr. Elizabeth Smith (Dartmouth College) and Dr. Steven King (University of Connecticut Health Science Center) for helpful discussions of I1 dynein and this manuscript.

## REFERENCES

- Ahmed NT, Mitchell DR. 2005. ODA16p, a *Chlamydomonas* flagellar protein needed for dynein assembly. *Mol Biol Cell* 16(10):5004–5012.
- Avidor-Reiss T, Maer AM, Koundakjian E, Polyanovsky A, Keil T, Subramaniam S, Zuker CS. 2004. Decoding cilia function: Defining specialized genes required for compartmentalized cilia biogenesis. *Cell* 117(4):527–539.
- Bower R, Perrone CA, O'Toole E, Fox L, Wirschell M, Sale WS, Porter ME. 2006. IC138 is required for regulation but not assembly of the I1 inner arm dynein. San Diego, CA: Abstracts from the 2006 Annual Meeting of the American Society for Cell Biology and published at [www.ASCB.org](http://www.ASCB.org).
- Brokaw CJ. 1985. Computer simulation of flagellar movement. VI. Simple curvature-controlled models are incompletely specified. *Biophys J* 48(4):633–642.
- Brokaw CJ. 1994. Control of flagellar bending: A new agenda based on dynein diversity. *Cell Motil Cytoskeleton* 28(3):199–204.
- Brokaw CJ, Kamiya R. 1987. Bending patterns of *Chlamydomonas* flagella: IV. Mutants with defects in inner and outer dynein arms indicate differences in dynein arm function. *Cell Motil Cytoskeleton* 8(1):68–75.
- Casey DM, Inaba K, Pazour GJ, Takada S, Wakabayashi K, Wilkerson CG, Kamiya R, Witman GB. 2003. DC3, the 21-kDa subunit of the outer dynein arm-docking complex (ODA-DC), is a novel EF-hand protein important for assembly of both the outer arm and the ODA-DC. *Mol Biol Cell* 14(9):3650–3663.
- Christensen ST, Pedersen LB, Schneider L, Satir P. 2007. Sensory cilia and integration of signal transduction in human health and disease. *Traffic* 8(2):97–109.
- Davenport JR, Yoder BK. 2005. An incredible decade for the primary cilium: A look at a once-forgotten organelle. *Am J Physiol Renal Physiol* 289(6):F1159–1169.
- Davis EE, Brueckner M, Katsanis N. 2006. The emerging complexity of the vertebrate cilium: New functional roles for an ancient organelle. *Dev Cell* 11(1):9–19.
- Deckman CM, Pennock DG. 2004. Dephosphorylation of inner arm 1 is associated with ciliary reversals in *Tetrahymena thermophila*. *Cell Motil Cytoskeleton* 57(2):73–83.
- DiBella LM, Benashski SE, Tedford HW, Harrison A, Patel-King RS, King SM. 2001. The Tctex1/Tctex2 class of dynein light chains. Dimerization, differential expression, and interaction with the LC8 protein family. *J Biol Chem* 276(17):14366–14373.
- DiBella LM, King SM. 2001. Dynein motors of the *Chlamydomonas* flagellum. *Int Rev Cytol* 210:227–268.
- DiBella LM, Sakato M, Patel-King RS, Pazour GJ, King SM. 2004a. The LC7 light chains of *Chlamydomonas* flagellar dyneins interact with components required for both motor assembly and regulation. *Mol Biol Cell* 15(10):4633–4646.
- DiBella LM, Smith EF, Patel-King RS, Wakabayashi K, King SM. 2004b. A novel Tctex2-related light chain is required for stability of inner dynein arm I1 and motor function in the *Chlamydomonas* flagellum. *J Biol Chem* 279(20):21666–21676.
- Dymek EE, Smith EF. 2006. Regulation of axonemal dynein activity by a calmodulin and radial spoke associated complex (CRC). San Diego, CA: Abstracts from the 2006 Annual Meeting of the American Society for Cell Biology and published at [www.ASCB.org](http://www.ASCB.org).
- El Zein L, Omran H, Bouvagnet P. 2003. Lateralization defects and ciliary dyskinesia: Lessons from algae. *Trends Genet* 19(3):162–167.
- Fox LA, Tritschler D, Porter ME, Sale WS. 2006. Regulation of flagellar inner arm dynein I1: The I1 $\beta$  heavy chain motor domain is necessary for control of microtubule sliding by phosphorylation. San Diego, CA: Abstracts from the 2006 Annual Meeting of the American Society for Cell Biology and published at [www.ASCB.org](http://www.ASCB.org).
- Freshour J, Yokoyama R, Mitchell DR. 2006. *Chlamydomonas* flagellar outer row dynein assembly protein Oda7 interacts with both outer row and I1 inner row dyneins. *J Biol Chem* 282(8):5404–5412.
- Gaillard AR, Diener DR, Rosenbaum JL, Sale WS. 2001. Flagellar radial spoke protein 3 is an A-kinase anchoring protein (AKAP). *J Cell Biol* 153(2):443–448.
- Gaillard AR, Fox LA, Rhea JM, Craig B, Sale WS. 2006. Disruption of the A-kinase anchoring domain in flagellar radial spoke protein 3 results in unregulated axonemal cAMP-dependent protein kinase activity and abnormal flagellar motility. *Mol Biol Cell* 17(6):2626–2635.

- Gardner LC, O'Toole E, Perrone CA, Giddings T, Porter ME. 1994. Components of a "dynein regulatory complex" are located at the junction between the radial spokes and the dynein arms in *Chlamydomonas* flagella. *J Cell Biol* 127(5):1311–1325.
- Gokhale A, Wirschell M, Sale WS. 2006. Characterization of the protein kinase CK1 that regulates dynein-driven microtubule sliding and is anchored in *Chlamydomonas* flagellar axonemes. San Diego, CA: Abstracts from the 2006 Annual Meeting of the American Society for Cell Biology and published at www.ASCB.org.
- Goodenough UW, Gebhart B, Mermall V, Mitchell DR, Heuser JE. 1987. High-pressure liquid chromatography fractionation of *Chlamydomonas* dynein extracts and characterization of inner-arm dynein subunits. *J Mol Biol* 194(3):481–494.
- Goodenough UW, Heuser JE. 1985a. Outer and inner dynein arms of cilia and flagella. *Cell* 41(2):341–342.
- Goodenough UW, Heuser JE. 1985b. Substructure of inner dynein arms, radial spokes, and the central pair/projection complex of cilia and flagella. *J Cell Biol* 100(6):2008–2018.
- Habermacher G, Sale WS. 1995. Regulation of dynein-driven microtubule sliding by an axonemal kinase and phosphatase in *Chlamydomonas* flagella. *Cell Motil Cytoskeleton* 32(2):106–109.
- Habermacher G, Sale WS. 1996. Regulation of flagellar dynein by an axonemal type-1 phosphatase in *Chlamydomonas*. *J Cell Sci* 109(Part 7):1899–1907.
- Habermacher G, Sale WS. 1997. Regulation of flagellar dynein by phosphorylation of a 138-kD inner arm dynein intermediate chain. *J Cell Biol* 136(1):167–176.
- Harrison A, Olds-Clarke P, King SM. 1998. Identification of the t complex-encoded cytoplasmic dynein light chain tctex1 in inner arm II supports the involvement of flagellar dyneins in meiotic drive. *J Cell Biol* 140(5):1137–1147.
- Hasegawa E, Hayashi H, Asakura S, Kamiya R. 1987. Stimulation of in vitro motility of *Chlamydomonas* axonemes by inhibition of cAMP-dependent phosphorylation. *Cell Motil Cytoskeleton* 8(4):302–311.
- Hendrickson TW, Perrone CA, Griffin P, Wuichet K, Mueller J, Yang P, Porter ME, Sale WS. 2004. IC138 is a WD-repeat dynein intermediate chain required for light chain assembly and regulation of flagellar bending. *Mol Biol Cell* 15(12):5431–5442.
- Hennessey TM, Kim DY, Oberski DJ, Hard R, Rankin SA, Pennock DG. 2002. Inner arm dynein 1 is essential for Ca<sup>++</sup>-dependent ciliary reversals in *Tetrahymena thermophila*. *Cell Motil Cytoskeleton* 53(4):281–288.
- Hou Y, Qin H, Folliot JA, Pazour GJ, Rosenbaum JL, Witman GB. 2007. Functional analysis of an individual IFT protein: IFT46 is required for transport of outer dynein arms into flagella. *J Cell Biol* 176(5):653–665.
- Howard DR, Habermacher G, Glass DB, Smith EF, Sale WS. 1994. Regulation of *Chlamydomonas* flagellar dynein by an axonemal protein kinase. *J Cell Biol* 127(6, Part 1):1683–1692.
- Iyadurai SJ, Li MG, Gilbert SP, Hays TS. 1999. Evidence for cooperative interactions between the two motor domains of cytoplasmic dynein. *Curr Biol* 9(14):771–774.
- Kagami O, Kamiya R. 1992. Translocation and rotation of microtubules caused by multiple species of *Chlamydomonas* inner-arm dynein. *J Cell Sci* 103(3):653–664.
- Kagami O, Kamiya R. 1995. Separation of dynein species by high-pressure liquid chromatography. *Methods Cell Biol* 47:487–489.
- Kamiya R. 1995. Exploring the function of inner and outer dynein arms with *Chlamydomonas* mutants. *Cell Motil Cytoskeleton* 32(2):98–102.
- Kamiya R. 2002. Functional diversity of axonemal dyneins as studied in *Chlamydomonas* mutants. *Int Rev Cytol* 219:115–155.
- Kamiya R, Kurimoto E, Muto E. 1991. Two types of *Chlamydomonas* flagellar mutants missing different components of inner-arm dynein. *J Cell Biol* 112(3):441–447.
- King SJ, Dutcher SK. 1997. Phosphoregulation of an inner dynein arm complex in *Chlamydomonas reinhardtii* is altered in phototactic mutant strains. *J Cell Biol* 136(1):177–191.
- King SM, Barbarese E, Dillman JF III, Patel-King RS, Carson JH, Pfister KK. 1996. Brain cytoplasmic and flagellar outer arm dyneins share a highly conserved Mr 8,000 light chain. *J Biol Chem* 271(32):19358–19366.
- Koonce MP, Knecht DA. 1998. Cytoplasmic dynein heavy chain is an essential gene product in *Dictyostelium*. *Cell Motil Cytoskeleton* 39(1):63–72.
- Lechtreck K-F, Witman GB. 2007. *Chlamydomonas reinhardtii* hidin is a central pair protein required for flagellar motility. *J Cell Biol* 176(4):473–482.
- Li JB, Gerdes JM, Haycraft CJ, Fan Y, Teslovich TM, May-Simera H, Li H, Blacque OE, Li L, Leitch CC, Lewis RA, Green JS, Parfrey PS, Leroux MR, Davidson WS, Beales PL, Guay-Woodford LM, Yoder BK, Stormo GD, Katsanis N, Dutcher SK. 2004. Comparative genomics identifies a flagellar and basal body proteome that includes the BBS5 human disease gene. *Cell* 117(4):541–552.
- Lindemann CB. 2002. Geometric Clutch model version 3: The role of the inner and outer arm dyneins in the ciliary beat. *Cell Motil Cytoskeleton* 52(4):242–254.
- Mastronarde DN, O'Toole ET, McDonald KL, McIntosh JR, Porter ME. 1992. Arrangement of inner dynein arms in wild-type and mutant flagella of *Chlamydomonas*. *J Cell Biol* 118(5):1145–1162.
- Mitchell DR. 2003a. Orientation of the central pair complex during flagellar bend formation in *Chlamydomonas*. *Cell Motil Cytoskeleton* 56(2):120–129.
- Mitchell DR. 2003b. Reconstruction of the projection periodicity and surface architecture of the flagellar central pair complex. *Cell Motil Cytoskeleton* 55(3):188–199.
- Mitchell DR, Sale WS. 1999. Characterization of a *Chlamydomonas* insertional mutant that disrupts flagellar central pair microtubule-associated structures. *J Cell Biol* 144(2):293–304.
- Mitchell DR, Yokoyama R. 2003. Structural analysis of central pair function in *Chlamydomonas* flagella. *Mol Biol Cell* 14:436a.
- Myster SH, Knott JA, Wysocki KM, O'Toole E, Porter ME. 1999. Domains in the 1alpha dynein heavy chain required for inner arm assembly and flagellar motility in *Chlamydomonas*. *J Cell Biol* 146(4):801–818.
- Nicastro D, McIntosh JR, Baumeister W. 2005. 3D structure of eukaryotic flagella in a quiescent state revealed by cryo-electron tomography. *Proc Natl Acad Sci USA* 102(44):15889–15894.
- Nicastro D, Schwartz C, Pierson J, Gaudette R, Porter M, McIntosh J. 2006. The molecular architecture of axonemes revealed by cryo-electron tomography. *Science* 313(5789):944–948.
- Norrander JM, Perrone CA, Amos LA, Linck RW. 1996. Structural comparison of tektins and evidence for their determination of complex spacings in flagellar microtubules. *J Mol Biol* 257(2):385–397.
- Oiwa K, Sakakibara H. 2005. Recent progress in dynein structure and mechanism. *Curr Opin Cell Biol* 17(1):98–103.
- Okita N, Isogai N, Hirono M, Kamiya R, Yoshimura K. 2005. Phototactic activity in *Chlamydomonas* 'non-phototactic' mutants deficient in Ca<sup>2+</sup>-dependent control of flagellar dominance or in inner-arm dynein. *J Cell Sci* 118(Part 3):529–537.
- Pan J, Wang Q, Snell WJ. 2005. Cilium-generated signaling and cilia-related disorders. *Lab Invest* 85(4):452–463.
- Pasquale SM, Goodenough UW. 1987. Cyclic AMP functions as a primary sexual signal in gametes of *Chlamydomonas reinhardtii*. *J Cell Biol* 105(5):2279–2292.

- Pazour GJ. 2004. Intraflagellar transport and cilia-dependent renal disease: The ciliary hypothesis of polycystic kidney disease. *J Am Soc Nephrol* 15(10):2528–2536.
- Pazour GJ, Agrin N, Leszyk J, Witman GB. 2005. Proteomic analysis of a eukaryotic cilium. *J Cell Biol* 170(1):103–113.
- Pazour GJ, Wilkerson CG, Witman GB. 1998. A dynein light chain is essential for the retrograde particle movement of intraflagellar transport (IFT). *J Cell Biol* 141(4):979–992.
- Perrone CA, Myser SH, Bower R, O'Toole ET, Porter ME. 2000. Insights into the structural organization of the I1 inner arm dynein from a domain analysis of the I $\beta$  dynein heavy chain. *Mol Biol Cell* 11(7):2297–2313.
- Perrone CA, Yang P, O'Toole E, Sale WS, Porter ME. 1998. The *Chlamydomonas* IDA7 locus encodes a 140-kDa dynein intermediate chain required to assemble the I1 inner arm complex. *Mol Biol Cell* 9(12):3351–3365.
- Piperno G. 1995. Regulation of dynein activity within *Chlamydomonas* flagella. *Cell Motil Cytoskeleton* 32(2):103–105.
- Piperno G, Mead K, Henderson S. 1996. Inner dynein arms but not outer dynein arms require the activity of kinesin homologue protein KHP1(FLA10) to reach the distal part of flagella in *Chlamydomonas*. *J Cell Biol* 133(2):371–379.
- Piperno G, Ramanis Z, Smith EF, Sale WS. 1990. Three distinct inner dynein arms in *Chlamydomonas* flagella: Molecular composition and location in the axoneme. *J Cell Biol* 110(2):379–389.
- Porter ME, Power J, Dutcher SK. 1992. Extragenic suppressors of paralyzed flagellar mutations in *Chlamydomonas reinhardtii* identify loci that alter the inner dynein arms. *J Cell Biol* 118(5):1163–1176.
- Porter ME, Sale WS. 2000. The 9 + 2 axoneme anchors multiple inner arm dyneins and a network of kinases and phosphatases that control motility. *J Cell Biol* 151(5):F37–F42.
- Quarby LM, Parker JD. 2005. Cilia and the cell cycle? *J Cell Biol* 169(5):707–710.
- Sakakibara H, Burgess S, Sakai Y. 2006. Conformational changes of inner-arm dynein-f from *Chlamydomonas* coupled with phosphorylation of its intermediate chain IC138. San Diego, CA: Abstracts from the 2006 Annual Meeting of the American Society for Cell Biology and published at www.ASCB.org.
- Sakakibara H, Kojima H, Sakai Y, Katayama E, Oiwa K. 1999. Inner-arm dynein c of *Chlamydomonas* flagella is a single-headed processive motor. *Nature* 400(6744):586–590.
- Sakakibara H, Nakayama H. 1998. Translocation of microtubules caused by the alpha, beta and gamma outer arm dynein subparticles of *Chlamydomonas*. *J Cell Sci* 111(Part 9):1155–1164.
- Sakakibara H, Takada S, King SM, Witman GB, Kamiya R. 1993. A *Chlamydomonas* outer arm dynein mutant with a truncated beta heavy chain. *J Cell Biol* 122(3):653–661.
- Setter PW, Malvey-Dorn E, Steffen W, Stephens RE, Linck RW. 2006. Tektin interactions and a model for molecular functions. *Exp Cell Res* 312(15):2880–2896.
- Silflow CD, Lefebvre PA. 2001. Assembly and motility of eukaryotic cilia and flagella. Lessons from *Chlamydomonas reinhardtii*. *Plant Physiol* 127(4):1500–1507.
- Smith EF. 2002a. Regulation of flagellar dynein by calcium and a role for an axonemal calmodulin and calmodulin-dependent kinase. *Mol Biol Cell* 13(9):3303–3313.
- Smith EF. 2002b. Regulation of flagellar dynein by the axonemal central apparatus. *Cell Motil Cytoskeleton* 52(1):33–42.
- Smith EF. 2007. Hydin seek: Finding a function in ciliary motility. *J Cell Biol* 176(4):403–404.
- Smith EF, Sale WS. 1991. Microtubule binding and translocation by inner dynein arm subtype I1. *Cell Motil Cytoskeleton* 18(4):258–268.
- Smith EF, Sale WS. 1992a. Regulation of dynein-driven microtubule sliding by the radial spokes in flagella. *Science* 257(5076):1557–1559.
- Smith EF, Sale WS. 1992b. Structural and functional reconstitution of inner dynein arms in *Chlamydomonas* flagellar axonemes. *J Cell Biol* 117(3):573–581.
- Smith EF, Yang P. 2004. The radial spokes and central apparatus: Mechano-chemical transducers that regulate flagellar motility. *Cell Motil Cytoskeleton* 57(1):8–17.
- Snell WJ, Pan J, Wang Q. 2004. Cilia and flagella revealed: From flagellar assembly in *Chlamydomonas* to human obesity disorders. *Cell* 117(6):693–697.
- Tai AW, Chuang JZ, Bode C, Wolfrum U, Sung CH. 1999. Rhodopsin's carboxy-terminal cytoplasmic tail acts as a membrane receptor for cytoplasmic dynein by binding to the dynein light chain Tctex-1. *Cell* 97(7):877–887.
- Takada S, Kamiya R. 1994. Functional reconstitution of *Chlamydomonas* outer dynein arms from alpha-beta and gamma subunits: Requirement of a third factor. *J Cell Biol* 126(3):737–745.
- Toba S, Sakakibara H, Porter ME, Sale WS, Oiwa K. 2006. Motility of the I $\alpha$  and I $\beta$  heavy chains of the inner arm dynein I1 of *Chlamydomonas* flagella. San Diego, CA: Abstracts from the 2006 Annual Meeting of the American Society for Cell Biology and published at www.ASCB.org.
- Wakabayashi K, Takada S, Witman GB, Kamiya R. 2001. Transport and arrangement of the outer-dynein-arm docking complex in the flagella of *Chlamydomonas* mutants that lack outer dynein arms. *Cell Motil Cytoskeleton* 48(4):277–286.
- Wargo MJ, Dymek EE, Smith EF. 2005. Calmodulin and PF6 are components of a complex that localizes to the C1 microtubule of the flagellar central apparatus. *J Cell Sci* 118(Part 20):4655–4665.
- Wargo MJ, McPeck MA, Smith EF. 2004. Analysis of microtubule sliding patterns in *Chlamydomonas* flagellar axonemes reveals dynein activity on specific doublet microtubules. *J Cell Sci* 117(Part 12):2533–2544.
- Wargo MJ, Smith EF. 2003. Asymmetry of the central apparatus defines the location of active microtubule sliding in *Chlamydomonas* flagella. *Proc Natl Acad Sci USA* 100(1):137–142.
- Warner FD, Satir P. 1974. The structural basis of ciliary bend formation. Radial spoke positional changes accompanying microtubule sliding. *J Cell Biol* 63(1):35–63.
- Wirschell M, Pazour G, Yoda A, Hirono M, Kamiya R, Witman GB. 2004. Oda5p, a novel axonemal protein required for assembly of the outer dynein arm and an associated adenylate kinase 10.1091/mbc.E03-11-0820. *Mol Biol Cell* 15(6):2729–2741.
- Wirschell M, Yanagisawa HA, Kamiya R, Witman GB, Porter ME, Sale WS. 2006. The role of the IC97 in I1-dynein assembly and axonemal anchoring. San Diego, CA: Abstracts from the 2006 Annual Meeting of the American Society for Cell Biology and published at www.ASCB.org.
- Woolley DM. 1997. Studies on the eel sperm flagellum. I. The structure of the inner dynein arm complex. *J Cell Sci* 110(Part 1):85–94.
- Yagi T, Minoura I, Fujiwara A, Saito R, Yasunaga T, Hirono M, Kamiya R. 2005. An axonemal dynein particularly important for flagellar movement at high viscosity. Implications from a new *Chlamydomonas* mutant deficient in the dynein heavy chain gene *DHC9*. *J Biol Chem* 280(50):41412–41420.

- Yamamoto R, Yagi T, Kamiya R. 2006. Functional binding of inner-arm dyneins with demembranated flagella of *Chlamydomonas* mutants. *Cell Motil Cytoskeleton* 63(5):258–265.
- Yanagisawa H-A, Kamiya R. 2004. A tektin homologue is decreased in *Chlamydomonas* mutants lacking an axonemal inner-arm dynein. *Mol Biol Cell* 15(5):2105–2115.
- Yang P, Fox L, Colbran RJ, Sale WS. 2000. Protein phosphatases PP1 and PP2A are located in distinct positions in the *Chlamydomonas* flagellar axoneme. *J Cell Sci* 113(Part 1):91–102.
- Yang P, Sale WS. 1998. The Mr 140,000 intermediate chain of *Chlamydomonas* flagellar inner arm dynein is a wd-repeat protein implicated in dynein arm anchoring. *Mol Biol Cell* 9(12):3335–3349.
- Yang P, Sale WS. 2000. Casein kinase I is anchored on axonemal doublet microtubules and regulates flagellar dynein phosphorylation and activity. *J Biol Chem* 275(25):18905–18912.
- Yang P, Yang C, Sale WS. 2004. Flagellar radial spoke protein 2 is a calmodulin binding protein required for motility in *Chlamydomonas reinhardtii*. *Eukaryot Cell* 3(1):72–81.