

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/10646842>

The clinical and pathogenetic significance of estrogen receptor- α expression in chronic liver diseases and liver carcinoma

ARTICLE in *CANCER* · AUGUST 2003

Impact Factor: 4.89 · DOI: 10.1002/cncr.11528 · Source: PubMed

CITATIONS

29

READS

10

7 AUTHORS, INCLUDING:



P. Lampertico

University of Milan

240 PUBLICATIONS 4,271 CITATIONS

SEE PROFILE



Guido Ronchi

Fondazione IRCCS Ca' Granda - Ospedale M...

26 PUBLICATIONS 1,798 CITATIONS

SEE PROFILE

The Clinical and Pathogenetic Significance of Estrogen Receptor- β Expression in Chronic Liver Diseases and Liver Carcinoma

Massimo Iavarone, M.D., Ph.D.
 Pietro Lampertico, M.D., Ph.D.
 Chiara Seletti, M.D., Ph.D.
 Maria Francesca Donato, M.D.
 Guido Ronchi, M.D.
 Ersilio Del Ninno, M.D.
 Massimo Colombo, M.D.

A.M. and A. Migliavacca Center, Division of Hepatology and FIRC Liver Cancer Unit, IRCCS Maggiore Policlinico Hospital, University of Milan, Milan, Italy.

Supported by grants from the Italian Ministry of University and Technological Research ("40%-CO-FIN 2000"), Consorzio Interuniversitario Trapianti d'Organo and Fondazione Italiana Ricerca Cancro (FIRC).

The authors thank Professor A. Morabito for statistical support, and Professor M. Levrero for his critical advice.

Address for reprints: Massimo Colombo, M.D., Ph.D., Division of Hepatology, IRCCS Maggiore Hospital, Via Pace 9, 20122 Milan, Italy; Fax: (011) 39-0250320700; E-mail: massimo.colombo@unimi.it

Received January 14, 2003; revision received April 11, 2003; accepted April 17, 2003.

BACKGROUND. Estrogen receptor- α (ER α) is variably expressed in hepatocellular carcinoma (HCC) and is believed to be correlated with prognosis and survival. Recently, another estrogen receptor (ER β) has been identified, but its relevance in liver diseases is unknown.

METHODS. The expression of ER β in the liver of 42 patients with HCC (10 with paired extratumoral tissues) and 26 with chronic liver disease without HCC was studied by a reverse transcriptase-polymerase chain reaction method, and correlated with the expression of ER α and severity of the liver disease.

RESULTS. Both ER β and wild-type ER α were found to be expressed more often in patients with chronic liver disease compared with those with HCC (69% vs. 45% [$P = 0.046$] and 46% vs. 10% [$P = 0.0008$], respectively). ERs were similarly expressed in HCC and in the paired extratumoral tissue. Wild-type receptors, either alone or together with the deleted mutants ER $\delta 5$, were more often coexpressed in chronic liver disease (58%) than in HCC (29%); in 13 tumors (31%), either ER $\delta 5$ or no receptors at all were detected ($P = 0.006$). Hepatitis B virus (HBV)-related tumors either did not appear to express ERs or expressed ER $\delta 5$ more often than hepatitis C virus (HCV)-related tumors (67% vs. 15%; $P = 0.007$). The same was true for multinodular compared with single nodular tumors (50% vs. 19%; $P = 0.04$).

CONCLUSIONS. Both receptors were expressed in chronic liver disease and neoplastic livers demonstrating different patterns in relation to the etiology and clinical presentation of the tumor. These differences might underscore different pathogenetic mechanisms in HBV-related and HCV-related HCC and a different evolutionary course for the tumor. *Cancer* 2003;98:529-34.

© 2003 American Cancer Society.

KEYWORDS: estrogen receptor (ER)- α , estrogen receptor- β , hepatocellular carcinoma (HCC), chronic hepatitis, cirrhosis, hepatitis B virus (HBV), hepatitis C virus (HCV).

Several lines of evidence suggest that sex hormones and their receptors may play a role in liver carcinogenesis.¹ Estrogen receptor- α (ER α) is expressed in the liver of both healthy individuals and patients with chronic hepatitis and hepatocellular carcinoma (HCC), apparently with no differences in the pattern of expression.^{2,3} Conversely, the mutant form with the entire exon 5 deleted (ER $\alpha\delta 5$) is preferentially expressed in patients with HCC compared with patients with normal livers or chronic hepatitis.^{4,5} Because ER $\alpha\delta 5$ lacks the hormone-binding domain, HCC patients expressing mutated receptors might form a subset of patients who respond less favorably to antiestrogen drugs.⁵

With the identification of ER β ,^{6,7} the cellular network of ERs

TABLE 1
Epidemiologic and Clinical Features of the 68 Patients Enrolled into the Study

	Chronic liver disease without HCC	HCC
No. of patients	26	42
Males, no.	18	35
Age (yrs) ^a	55 ± 12	65 ± 9
HBsAg	2	9
Anti-HCV	23	26
Alcohol consumption	1	2
Mixed	0	2
Cryptogenic	0	3

HCC: hepatocellular carcinoma; HBsAg: hepatitis B surface antigen; HCV: hepatitis C virus.

^a Mean ± the standard deviation.

appears to be more complex than previously thought, thus partly explaining the different patterns of clinical response to anticancer hormones reported to date.⁶ Despite the high degree of sequence homology between ER α and ER β ,⁸ animal studies have revealed distinct patterns of expression for each receptor, with ER α being expressed preferentially in female reproductive organs and ER β being predominant in nonreproductive organs and in the male reproductive tract.⁶⁻⁹ The ER α :ER β expression ratio apparently changes during carcinogenesis and is believed to play a role in tumor development.¹⁰ In tissues of the breast, prostate, and colon, ER β appears to be underexpressed during malignant transformation and tumor progression.¹⁰⁻¹⁴

Coexpression of ER α and ER β may further contribute to the range of clinical responses to hormone therapy because it increases the chances of producing heterodimer receptors, whose transcriptional profiles are different from those given by homodimer receptors.⁸

To assess whether the pattern of ERs plays a role in human liver carcinogenesis, we studied the expression of ER subtypes in the liver in a well defined group of patients with chronic hepatitis and HCC and correlated the results with disease etiology and severity.

MATERIALS AND METHODS

Patients

The current study included 68 patients who were attending the Liver Center in the Division of Hepatology at Maggiore Hospital in Milan (Table 1). There were 26 patients with chronic liver disease without HCC (9 with chronic viral hepatitis with a mean necroinflammatory activity score of 9 ± 2 as assessed by the Ishak score and 17 with compensated cirrhosis). Forty-two patients had a histologic diagnosis of HCC (40 patients

had underlying cirrhosis and 26 had a single-nodular tumor measuring 38 mm, on average). The grade of tumor differentiation was assessed according to the Edmondson-Steiner score and was high (Grade 1) in 2 patients (5%), moderate (Grade 2) in 16 patients (38%), poor (Grades 3-4) in 19 patients (45%), and not evaluable in 5 patients (12%). Ten patients with HCC also had paired tissues from extratumoral liver. None of the patients had received antiviral or antitumoral treatment at the time of the study. All liver biopsies were performed with a 16-gauge Tru-Cut[®] needle (TSK, Tokyo, Japan) under ultrasound guidance. Approximately two-thirds of each tissue sample was formalin fixed and embedded in paraffin for routine histopathology, whereas the remainder of the tissue sample was snap-frozen at -80°C until RNA extraction. Overall, 78 liver tissue samples were analyzed.

ER Assay

Total RNA was extracted from frozen tissue by RNeasy B (TEL-TEXT, Inc., Friendswood, TX), and the acid guanidium-thiocyanate-phenol-chlorophorm method. Primers were designed from published gene sequences, with particular attention paid to the high homology between ER α and ER β . In particular, the primer couples for ER α were located in exons 4 and 6 (ER1: 5'-GGAGACATGAGAGCTGCCAAC-3', ER2: 5'-CCAGCAGCATGTCGAAGATC-3', ER3: 5'-CACAAAGCGCCAGAGAGATGA-3', and ER4: 5'-CATCATGCGGAACCGAGATG-3'), according to Fuqua et al.,¹⁵ to allow the amplification of a possible exon 5 deletion. With regard to primer couples for ER β , these were located in exons 2 and 7 (ER- β 1: 5'-TCACTTCTGCGCTGTCTGCAGCG-3', ER- β 2: 5'-CCTGGGTCGCTGTGACCAGA-3', ER- β 3: 5'-GGCCAAGAGAAGTGGCGGCCACG-3', and ER- β 4: 5'-AAACCTTGAAGTAGTTGCCAGGAGC-3'), according to the method of Vladusic et al.¹⁶

Total RNA was reverse transcribed to cDNA. A total of 10 μL of RNA were added to a volume of 20 μL containing 4 μL of polymerase chain reaction (PCR) buffer 5 \times , 2.5 μL of dNTPs (2.5 mM), 5 μL of specific outer antisense primer (ER4 for ER α and ER β 2 for ER β), 15 units (U) of reverse transcriptase-avian myeloblastosis virus (RT-AMV) (Thermoscript[™] [RNase H⁻ Reverse Transcriptase], Life Technologies, Gaithersburg, MD), 40 U of RNase inhibitor (RNase OUT; Invitrogen, Groningen, the Netherlands), and 1 μL of DTT (0.1 mM, Life Technologies). Complementary DNA (cDNA) was synthesized by incubation at 42 $^{\circ}\text{C}$ for 60 minutes, cooled rapidly at 4 $^{\circ}\text{C}$, and stored at -20°C until first PCR.

Approximately 10 μL of cDNA were added to a final volume of 50 μL containing 5 μL of PCR buffer 10 \times , 3 μL of dNTPs (2.5 mM), 2 μL each primer

FIGURE 1. Example of a deleted sequence of estrogen receptor- β (ER β) in a patient with cirrhosis (CIR14- β). Reference sequences (ER α -ref and ER β -ref) were found in GenBank (National Institutes of Health, Bethesda, MD), access numbers NM_000125 and NM_001437, respectively.

		exon IV	exon V
ER α -ref	GAGGGTGCCAG	GCTTTGTGGATTGACCTCCATGATCAGGTCCACCTTCTAGAATGTG	
ER β -ref	GAAGATTCCCG	GCTTTGTGGAGCTCAGCCTGTTCGACCAAGTGGCGCTCTTGAGAGCT	
CIR14- β	GAAGATTCCCG	-----	-----
		811	
ER α -ref	CCTGGCTAGAGATCCTGATGATTGGTCTCGTCTGGCGCTCCATGGAGCACCCAGTGAAGC		
ER β -ref	GTGGATGGAGGTGTTAATGATGGGGCTGATGTGGCGCTCAATTGACCACCCGGCAAGC		
CIR14- β	-----		
			exon VI
ER α -ref	TACTGTTTGCTCCTAACTTGCTCTTGACAG	GAACCAGGGAAAAATGTGTAGAGGGCATG	
ER β -ref	TCATCTTTGCTCCAGATCTTGTCTTGACAG	GGATGAGGGGAAATGCGTAGAAGGAATT	
CIR14- β	-----	GGATGAGGGGAAATGCGTAGAAGGAATT	
			951

couples (ER3/ER4 primers for ER α and ER β 1/ER β 2 for ER β), and 2.5 U of Taq Gold DNA polymerase (AmpliTaq Gold[®]; Applied Biosystems, Foster City, CA). The reaction mixture was heated at 95 °C for 4 minutes and then was subjected to a total of 40 cycles (1 minute at 95 °C, 1 minute at 58 °C for ER α and 62 °C for ER β , and 1 minute at 72 °C). The reaction cycle was terminated by heating at 72 °C for 10 minutes and cooling at 4 °C. Second PCR steps were performed with 5 μ L of the product of the first step of PCR, using the inner primer couples (ER1/ER2 primers for ER α and ER β 3/ER β 4 for ER β), with the same profile as the first PCR. A total of 5 μ L of the final PCR product was analyzed by 3% agarose gel electrophoresis, stained with ethidium bromide, and detected by ultraviolet lights.

The expected size was 438 base pairs (bp) for ER α and 430 bp for ER β . Deletion mutants were identified by the different electrophoretic mobility and confirmed by direct sequencing, after purification (QIAquick PCR Purification Kit; Qiagen, Milan, Italy).

All samples that tested negative for either ER α or ER β were retested and β -actin was amplified as a control of RNA quality. As a positive control, total RNA from MCF-7 breast carcinoma cells was reverse-transcribed and amplified in parallel with samples under investigation. As a negative control, reaction mixtures without RNA were processed.

Histopathologic Studies

Liver tissue was fixed in 10% formalin, embedded in paraffin, and routinely stained with hematoxylin and eosin. The histologic diagnoses of chronic hepatitis, cirrhosis, and HCC were based on internationally accepted criteria. Each liver biopsy was reviewed by a single pathologist (M. F. D.) and scored for histologic activity¹⁷ or grade of tumor cell differentiation.¹⁸

Serum Assays

Hepatitis B surface antigen (HBsAg) and antibody to HBsAg were detected by a microparticle enzyme immunoassay (AXSYM; Abbott Laboratories, North Chi-

cago, IL). Antibody to hepatitis C virus (HCV) was measured by a second-generation test system (Ortho DS, Raritan, NJ). Serum α -fetoprotein (AFP) was measured using a standard assay (immunoradiometric assay [IRMA]; Abbott Laboratories).

Statistical Analysis

The Fisher exact test, chi-square test, and Student *t* test were used to test the differences between groups.

RESULTS

All 78 tissue samples reacted for β -actin and therefore were analyzed for ER mRNA.

Chronic Liver Disease without HCC

ER β mRNA was detected in 18 patients with chronic liver disease (69%). Fifteen patients had wild-type ER β alone, 1 patient coexpressed wild-type ER β and a deletion variant, and 2 patients were found to have the deletion variants alone. Sequence analysis of the ER β deletion variant PCR product revealed the absence of 139 bp within the hormone-binding domain (i.e., the presence of the mutant ER β 85) (Fig. 1). ER α mRNA was detected in all tissues from patients with chronic liver disease. Twelve patients had wild-type ER α , 13 coexpressed wild-type ER α and ER α 85, and 1 patient had ER α 85. Combining the ER α and ER β results, we obtained four different patterns. Fifteen tissue samples (58%) coexpressed ER α and ER β (either the wild-type alone or together with the deleted mutants), 1 tissue sample (4%) expressed ER α 85 together with wild-type ER β , and 10 tissue samples (38%) expressed wild-type ER α alone or in association with ER β 85, together with ER β 85 or without ER β expression. All tissue samples demonstrated some form of ER expression (Table 2).

Hepatocellular Carcinoma

ER β mRNA was detected in 19 HCC tumor samples (45%). Fifteen samples expressed wild-type ER β alone, 1 sample coexpressed wild-type ER β and ER β 85, and 3

TABLE 2
Expression of ER α and ER β mRNA in 26 Chronic Liver Disease Tissues and 42 HCC Specimens

Liver tissue	Patterns of ER mRNA expression			
	$\alpha+\beta+$	$\alpha-\beta+$	$\alpha+\beta-$	$\alpha-\beta-$
Chronic liver disease without HCC	15 (58%)	1 (4%)	10 (38%)	0
HCC	12 (29%)	4 (9%)	13 (31%)	13 (31%)

$P = 0.006$, chi-square test.

ER: estrogen receptor; HCC: hepatocellular carcinoma; $\alpha+$: either ER α wt alone or both ER α wt and ER $\alpha\delta 5$; $\alpha-$: ER $\alpha\delta 5$ or no expression of ER α at all; $\beta+$: either ER β wt or both ER β wt and ER $\beta\delta 5$; $\beta-$: ER $\beta\delta 5$ or no expression of ER β at all.

samples expressed ER $\beta\delta 5$ only. ER α mRNA was detected in 41 HCC samples (98%). Four patients had wild-type ER α , 21 samples coexpressed wild-type ER α and ER $\alpha\delta 5$, and 16 samples were found to express ER $\alpha\delta 5$. Combining the results of ER α and ER β expression provided four different patterns. Twelve samples (29%) co-expressed ER α and ER β , either wild-type alone or associated with deleted forms; 4 samples (9%) expressed ER $\alpha\delta 5$ together with wild-type ER β ; 13 samples (31%) expressed wild-type ER α alone or associated with ER $\alpha\delta 5$ together, with ER $\beta\delta 5$ or without ER β expression; and 13 samples (31%) expressed either deleted forms or no ER mRNA. The four patterns of ER expression did not appear to be influenced by age, gender, alcohol consumption, portal invasion by the tumor, serum level of AFP, or tumor grading (Table 3). However, more hepatitis B virus (HBV)-related tumors either did not express ERs or expressed only deleted forms compared with HCV tumors (67% vs. 15%; $P = 0.007$). Either negative or deleted ER was more common among multinodular tumors compared with single-nodular tumors (50% vs. 19%; $P = 0.04$).

ER β was found to be expressed less frequently in HCC compared with in chronic hepatitis or cirrhotic tissues (45% vs. 69%; $P = 0.046$), whereas ER α was similarly expressed in both. However, wild-type ER α was expressed significantly more often in chronic hepatitis cases (5 of 9 cases) compared with cirrhosis cases (7 of 17 cases) and HCC cases (4 of 42 cases) (chi-square test, $P = 0.0017$). The four patterns of expression of ER mRNA significantly differed for chronic liver disease and HCC tissues (Table 2). Benign tissues co-expressed wild-type forms of both ER α and ER β (58%) alone or together with the deleted forms more often than malignant tissues (29%), whereas 31% of tumors were found to have either deleted forms of both receptors or no receptors at all, a pattern that was not present in benign liver disease cases (overall, $P = 0.006$).

Among the 10 extratumoral cirrhotic livers, 7 cases

TABLE 3
Clinical and Histologic Characteristics of 42 HCC Specimens According to ER status

	Patterns of ER mRNA expression			
	$\alpha+\beta+$	$\alpha-\beta+$	$\alpha+\beta-$	$\alpha-\beta-$
Patients	12	4	13	13
Males	9	3	11	12
Age (yrs) ^a	64 \pm 6	65 \pm 9	64 \pm 6	65 \pm 10
HBsAg	1	1	1	6
Anti-HCV	8	3	11	4
Alcohol consumption	0	0	1	1
Multifocal HCC ^b	4	1	3	8
Portal invasion ^b	2	0	1	1
Serum AFP, 7 IU/mL	9	3	7	10
Tumor grade ^c 1-2	4	2	7	5
3-4	6	2	3	8

HCC: hepatocellular carcinoma; ER: estrogen receptor; $\alpha+$: either ER α wt alone or both ER α wt and ER $\alpha\delta 5$; $\alpha-$: ER $\alpha\delta 5$ or no expression of ER α at all; $\beta+$: either ER β wt or both ER β wt and ER $\beta\delta 5$; $\beta-$: ER $\beta\delta 5$ or no expression of ER β at all; HBsAg: hepatitis B surface antigen; HCV: hepatitis C virus; AFP: α -fetoprotein.

^a Mean \pm the standard deviation.

^b Detected by abdominal ultrasound.

^c According to Edmondson-Steiner score.

had no or deleted ER β . Seven extratumoral samples expressed ER $\alpha\delta 5$ alone. The analysis of paired biopsies demonstrated no differences in ER expression between tumoral and extratumoral samples.

DISCUSSION

Carcinogenesis is a complex mechanism that involves constitutive cell proliferation, with the expression of ER believed to play an important role in this process.¹ Estrogens, in fact, are main regulators of those cell functions associated with the growth and development of well defined tumors such as those of the uterus and breast.¹⁹ The fact that the effects of estrogen and its antagonist on various target tissues are markedly different and can be largely influenced by alterations in the structure of ER itself¹⁹ makes the study of ER expression of potential interest in liver carcinogenesis as well. A potential role for ER in liver carcinogenesis has been suggested by the striking prevalence of HCC observed in males, the increased expression of the variant transcript of ER α (ER $\alpha\delta 5$) in the peritumoral cirrhotic tissue of patients with HCC, and the correlation between ER α and increased proliferation rates in HCC cases.¹ Recently, Villa et al.⁵ demonstrated high rates of ER $\alpha\delta 5$ expression in men at high-risk for HCC development.

To our knowledge, this is the first study to report the expression of ER β and its variants in the liver of patients with chronic liver diseases and liver carci-

noma. Previously, ER β was found to be expressed in breast, prostate, and gastrointestinal tissues of healthy individuals or patients with a variety of clinical conditions including cancer.¹⁰⁻¹⁴ We found that ER β was underexpressed compared with ER α in patients with chronic hepatitis or cirrhosis and those with HCC, with a tendency toward loss of ER β expression in HCC patients compared with those with chronic inflammatory liver diseases. Because a similar behavior of ER β already has been demonstrated during neoplastic transformation in breast, prostate, and gastrointestinal tissues,¹⁰⁻¹⁴ one could hypothesize that the neoplastic transformation of several epithelial lineages, including the liver, is associated with a reduction in expression of ER β .

In the current study, all but one tissue sample expressed ER α . In greater than half the cases, ER α was expressed as the deletion mutant ER $\alpha\delta 5$, with the tissue expression of this receptor being higher in patients with HCC than in patients with chronic hepatitis or cirrhosis, as already demonstrated by Villa et al.⁵ The higher prevalence of ER $\alpha\delta 5$ in neoplastic livers and paired extratumoral samples in the current study is in keeping with the hypothesis that sex hormones may play a role in HCC.⁵ The presence of mutated ER α may be biologically significant because the growth rate of HCC in patients with this variant was found to be significantly higher than in tumors expressing wild-type ER α only.²⁰ Expression of ER $\alpha\delta 5$ also may be a useful clinical predictor because in one study²¹ patients with this variant were found to have shorter survivals than those expressing wild-type ER α . Moreover, ER $\alpha\delta 5$ -positive tumors are more resistant to palliative treatment with antiestrogen therapy, whereas they respond better to progestinic drugs in terms of patient survival at 1 year.²²

To our knowledge, the current study was the first to assess the clinical significance of ER α and ER β coexpression in liver tissue, and demonstrated higher coexpression of ER α and ER β in patients with chronic liver disease compared with those with HCC (58% vs. 29%) and excess ER $\alpha\delta 5$ associated with wild-type ER β among patients with severe liver disease and/or HCC. These other observations suggest that a lack of coexpression could have pathogenetic implications in HCC patients. A previous work reported less expression of wild-type ER α among patients with cirrhosis and HCC with respect to patients with mild liver disease. In that study, diminution of ER α paralleled the increased expression of ER $\alpha\delta 5$. An *in vitro* study²³ demonstrated that ER $\alpha\delta 5$ could inhibit wild-type ER β , thus confirming the interaction between ERs as well as the potential importance of ER coexpression in a patient's response to hormone therapy. Our finding that the

deletion mutant of ER β mRNA lacked a domain that exactly corresponds to exon 5 of ER α , parallels the observations in MCF-7 breast carcinoma cells and in the estrogen-unresponsive breast carcinoma cell line MDA-MB-231¹⁶ and stresses further the importance of assessing ER expression in the liver prior to the initiation of hormone therapy in patients with HCC.

Isolated expression of wild-type ER α was demonstrated in six patients with chronic hepatitis or cirrhosis and in two patients with a well differentiated single-nodular HCC. Expression of ER α only in patients with liver disease is unprecedented because previous studies have lacked concurrent analyses for ER β .

It is interesting to note that we demonstrated lower ER β expression associated with ER $\alpha\delta 5$ in HCC and in paired extratumoral cirrhotic tissue. *In vitro* studies have demonstrated a correlation between ER α and ER β , often in forms of heterodimer-activated receptors, a condition in which the transcriptional profiles are different from those generated by homodimer-activated receptors.⁸ Coexpression of ER α and ER β may have a significant clinical impact because epithelial tumors that coexpress ER α and ER β are reported to be less invasive and dedifferentiated than tumors expressing a single ER.¹⁰⁻¹⁴ Studies of chemical carcinogenesis suggested that ER may modulate HCC risk by inhibiting the malignant transformation of preneoplastic liver cells.^{24,25} It also is interesting to note that we found that wild-type forms of ER were more often coexpressed in the less severe, single-nodular tumors compared with multinodular HCCs.

In the current study, ER α and ER β were found to be underexpressed in HBV-related tumors in terms of either an excess of deletion mutants for both receptors or no ER expression. These data are in keeping with recent studies based on polymorphic microsatellite markers that suggested the existence of two distinct chromosome patterns underlying distinct pathways of hepatocarcinogenesis, viral etiology, and clinical severity.²⁶ HBV-related tumors appear to have greater chromosome instability, poorer differentiation, and poorer prognosis compared with HBV-negative tumors. The number and size of HCC nodules, that are clinically significant surrogates for liver cell dedifferentiation and vascular invasiveness, also were found to be the two stronger predictors of survival after radical treatment of HCC.²⁷

The results of the current study demonstrated expression of ER β in both chronic liver disease and liver carcinoma, and a change in the coexpression pattern of ER α and ER β in patients with HCC. The differences in ER expression may point to different pathogenetic mechanisms in HBV-related and HCV-related HCC and a different evolutionary tumor course.

REFERENCES

- De Maria N, Manno M, Villa E. Sex hormones and liver cancer. *Mol Cell Endocrinol*. 2002;193:59–63.
- Porter LE, Elm MS, Van Thiel DH, Dugas MC, Eagon PK. Characterization and quantitation of human hepatic estrogen receptor. *Gastroenterology*. 1983;84:704–712.
- Rossini GP, Baldini GM, Villa E, Manenti F. Characterization of estrogen receptor from human liver. *Gastroenterology*. 1989;96:1102–1109.
- Villa E, Camellini L, Dugani A, et al. Variant estrogen receptor messenger RNA species detected in human primary hepatocellular carcinoma. *Cancer Res*. 1995;55:498–500.
- Villa E, Dugani A, Moles A, et al. Variant liver estrogen receptor transcripts already occur at early stage of chronic liver disease. *Hepatology*. 1998;27:983–988.
- Mosselman S, Polman J, Dijkema R. ER β : identification and characterization of a novel estrogen receptor. *FEBS Lett*. 1996;392:49–53.
- Kuiper GGJM, Enmark E, Peltö-Huikko M, Nilsson S, Gustafsson JA. Cloning a novel estrogen receptor expressed in rat prostate and ovary. *Proc Natl Acad Sci U S A*. 1996;93:5925–5930.
- Couse JF, Korach KS. Estrogen receptors null mice: what we have learned and where will they lead us? *Endocr Rev*. 1999;20(3):358–417.
- Couse JF, Lindzey J, Grandien K, Gustafsson JA, Korach KS. Tissue distribution and quantitative analysis of estrogen receptor- α (ER α) and estrogen receptor- β (ER β) messenger ribonucleic acid in the wild-type and ER α -knock-out mouse. *Endocrinology*. 1997;138:4613–4621.
- Leygue E, Dotzlaw H, Watson PH, Murphy LC. Altered estrogen receptor α and β messenger RNA expression during human breast tumorigenesis. *Cancer Res*. 1998;58:3197–3201.
- Horvath LG, Henshall SM, Lee CS, et al. Frequent loss of estrogen receptor- β expression in prostate cancer. *Cancer Res*. 2001;61:5331–5335.
- Foley EF, Jazaeri AA, Shupnik MA, Jazaeri O, Rice LW. Selective loss of estrogen receptor β in malignant human colon. *Cancer Res*. 2000;60:245–248.
- Campbell-Thompson M, Lynch JJ, Bhardwaj B. Expression of estrogen receptor (ER) subtypes and ER β isoforms in colon cancer. *Cancer Res*. 2001;61:632–640.
- Roger P, Sahla ME, Makela S, Gustafsson JA, Baldet P, Rochefort H. Decreased expression of estrogen receptor β protein in proliferative preinvasive mammary tumors. *Cancer Res*. 2001;61:2537–2541.
- Fuqua SAW, Fitzgerald SD, Chamness GC, et al. Variant human breast estrogen receptor with constitutive transcriptional activity. *Cancer Res*. 1991;51:105–109.
- Vladusic EA, Hornby AE, Guerra-Vladusic FK, Lupu R. Expression of estrogen receptor β messenger RNA variant in breast cancer. *Cancer Res*. 1998;58:210–214.
- Ishak K, Baptista A, Bianchi L, et al. Histopathological grading and staging of chronic hepatitis. *J Hepatol*. 1995;22:696–699.
- Edmondson HA, Steiner PE. Primary carcinoma of the liver: a study of 100 cases among 48,900 necroscopies. *Cancer*. 1954;7:462–503.
- Murphy LC, Dotzlaw H, Leygue E, Douglas D, Coutts A, Watson PH. Estrogen receptors variants and mutations. *J Steroid Biochem Mol Biol*. 1997;62:363–372.
- Villa E, Camellini L, Fantoni E, et al. Type of estrogen receptors determines response to antiestrogen therapy in hepatocellular carcinoma. *Cancer Res*. 1996;56:3883–3885.
- Villa E, Ferretti I, Buttafuoco P, Grottola A, Ferrari A, Manenti F. Natural history of inoperable hepatocellular carcinoma: estrogen receptor status in the tumor is the strongest prognostic factor for survival. *Hepatology*. 2000;32:233–238.
- Villa E, Ferretti I, Grottola A, et al. Hormonal therapy with megestrol in inoperable hepatocellular carcinoma characterized by variant estrogen receptors: a prospective, randomized, controlled trial. *Br J Cancer*. 2001;84:881–885.
- Inoue S, Ogawa S, Horie K, et al. An estrogen receptor β isoform that lacks exon 5 has dominant negative activity on both ER α and ER β . *Biochem Biophys Res Commun*. 2000;279:814–819.
- Mishkin SY, Farber E, Ho RK, Mulay S, Mishkin S. Tamoxifen alone or in combination with estradiol-17 β inhibits the growth and malignant transformation of hepatic hyperplastic nodules. *Eur J Cancer Clin Oncol*. 1985;21:615–623.
- Shimizu I, Yasuda M, Mizobuchi Y, et al. Suppressive effect of oestradiol on chemical hepatocarcinogenesis in rats. *Gut*. 1998;42:112–119.
- Laurent-Puig P, Legoix P, Bluteau O, et al. Genetic alterations associated with hepatocellular carcinomas define distinct pathways of hepatocarcinogenesis. *Gastroenterology*. 2001;120:1763–1773.
- McPeake JR, O'Grady JC, Zaman S, et al. Liver transplantation for primary hepatocellular carcinoma: tumor size and number determine outcome. *J Hepatol*. 1993;18:226–234.