

Chiral Recognition of Cyclic α -Hydroxyketones by CD-Sensitive Zinc Tetraphenylporphyrin Tweezer

HIDEKI ISHII,¹ YIHUI CHEN,¹ ROSS A. MILLER,² SANDOR KARADY,²

KOJI NAKANISHI,^{1*} AND NINA BEROVA^{1*}

¹Department of Chemistry, Columbia University, 3000 Broadway, New York, New York 10027

²Department of Process Research, Merck and Co., P.O. Box 2000, Rahway, New Jersey 07765

Dedicated to Professor Piero Salvadori, Pisa, on the occasion of his 70th birthday.

ABSTRACT A combined chemical/chiroptical microscale protocol for the determination of absolute configurations of cyclic α -hydroxyketones is described. The hydroxyl group in cyclic α -hydroxyketones is converted into (3-aminopropylamino)acetate ($\text{NH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{NHCH}_2\text{COOR}$), or more generally, according to a newly developed protocol, into (3-hydroxypropylamino)acetate group ($\text{HOCH}_2\text{CH}_2\text{CH}_2\text{NHCH}_2\text{COOR}$). The resultant conjugated compound forms a 1:1 host–guest complex with a dimeric zinc porphyrin tweezer, which exhibits exciton-coupled bisignate CD spectrum centered around the 420-nm porphyrin Soret band due to induced helicity between the two porphyrins in the complex. The absolute configurations of the α -stereogenic center is then determined by comparison of the sign of the observed CD exciton couplet of the complex with that of the preferred porphyrin twist predicted by the Merck Molecular Force Field (MMFFs) method. *Chirality* 17:305–315, 2005. © 2005 Wiley-Liss, Inc.

KEY WORDS: α -hydroxyketones; circular dichroism; exciton chirality; porphyrin tweezer; host–guest complexation

Chiral cyclic α -hydroxyketones are frequently found in bioactive compounds and are also important building blocks for the syntheses of complex chiral compounds including natural products and pharmaceutically important compounds,^{1–8} e.g., HIV protease inhibitors, such as indinavir (Crivian), which has been used in controlling the progression of AIDS in HIV-infected patients since 1996.^{9–13} However, due to the fast emergence of resistant viral strains of HIV,^{14,15} second-generation analogues with improved potency against resistant strains and improved pharmacokinetic profiles are needed. The Merck Research Laboratories synthesized compound **I**, which showed superior properties versus indinavir and was chosen as a developmental candidate (Chart 1).^{16–18} The synthesis of non-racemic amino chromanol fragment **II** has presented a synthetic challenge and several synthetic routes to **II** have been identified via the chiral non-racemic hydroxyketone **III**.¹⁹

Many synthetic methods have been developed for the preparation of chiral α -hydroxyketones,^{19–22} and their absolute configurations have been determined by stereoselective reactions,^{23,24} by spectroscopic methods such as Mosher methods,¹⁹ or by comparison of optical rotation/CD with known compounds.^{25,26} Although these methods have been used in various specific cases, they are not general enough to be applicable to all α -hydroxyketones, such as closely related hydroxyketones that Merck synthesized as part of this program. For example, in at least one case an assignment was tentatively made based on similar retention times by chiral HPLC. In addition, configurational analysis has been hampered by various

difficulties.^{27,28} In the application of the dibenzoate chirality method, reduction of α -hydroxyketones to 1,2-diols is required. However, this reduction frequently produces cis and trans diol mixtures, thus necessitating further stereochemical assignments.^{22,29}

In the following we describe studies aimed at developing a combined chemical/chiroptical protocol for configurational assignments of cyclic α -hydroxyketones by application of the exciton-coupled CD method based on chiral recognition by dimeric zinc porphyrin host **1**. As typical ketones, we chose five cyclic α -hydroxyketones, (S)- α -hydroxychromanone **2**, (R)-hydroxytetralone **3**, (R)- α -hydroxybenzosuberone **4**, (R)- α -hydroxycyclohexanone **5**, and (R)- α -hydroxycycloheptanone **6**, which were synthesized by Sharpless asymmetric dihydroxylation of the corresponding ketones (Chart 2).¹⁹

In the conventional application of the exciton method, the substrate should contain at least two covalently attached chromophores. More recently, a supramolecular approach based on chiral exciton coupling between porphyrin chromophores has been developed. It is based

This article includes Supplementary Material available via the Internet at <http://www.interscience.wiley.com/jpages/0899-0042/suppmat>
Contract grant sponsor: NIH; Contract grant number: GM 34509 (K.N., N.B.)

*Correspondence to: K. Nakanishi, E-mail: kn5@columbia.edu and N. Berova, E-mail: ndb1@columbia.edu

Received for publication 8 February 2005; Accepted 22 February 2005

DOI: 10.1002/chir.20166

Published online in Wiley InterScience (www.interscience.wiley.com).




Chart 1.

on a 1:1 complex formation between a chiral substrate linked to an achiral bidentate carrier (guest) and an achiral zinc tetraphenylporphyrin molecular tweezer **1** (host). Because the complexation proceeds in a stereocontrolled manner, it favors 1:1 host–guest complexes that adopt a preferred porphyrin–porphyrin twist. The latter leads to a strong induced CD exciton couplet in the porphyrin Soret band region which is diagnostic for the absolute configuration assignment of the chiral substrate.^{30–37} In the following an improved method is described that allows the stereochemical determination of cyclic α -hydroxyketones **2–6** by usage of an improved carrier molecule (Scheme 1).

MATERIALS AND METHODS

General

High-resolution FABMS was performed on a JMS-110/110 tandem mass spectrometer (JMS, Tokyo, Japan) using a *m*-nitrobenzyl alcohol (*m*-NBA) matrix. ¹H NMR spectra were recorded on a Bruker DPX-300 (300 MHz; Bruker, Billerica, MA) spectrometer in CDCl₃ or CD₃OD. Chemical shifts are reported in ppm relative to Me₄Si (0.0 ppm) or CD₂HOD (3.3 ppm). Data are reported as follows: (s) singlet, (d) doublet, (t) triplet, and (m) multiplet; coupling constants are in Hz; integration. The CD spectra, recorded on a JASCO-810 spectrophotometer, were converted into $\Delta\epsilon_{\text{max}}$ [l mol^{−1} cm^{−1}]/ λ [nm] units. The zinc tetraarylporphyrin tweezer (**1**) is commercially available from TCI (Portland, OR, catalogue no. P1364).

Computational Section

Molecular modeling calculations were performed as described earlier^{33,34,36,37} with the following modification: (i) sampling of all the structures was adopted within 15 kJ/mol instead of 10 kJ/mol of the lowest-energy conformer over 1,000 fully optimized steps, (ii) the distance checks for primary alcohol O–Zn were set to 2.4 ± 0.5 Å,³⁸ and (iii) the dihedral angles around the stereogenic center

[H α –C α –O–C(=O)] of the guest, α -hydroxyketone derivatives, were locked in order to find the lowest-energy conformations within a limited calculation time.^{39,40} After all the calculations (3,000 times), the most stable conformers within a 15-kJ/mol energy window were analyzed using the Boltzmann distribution based on energy difference.

Synthesis of (3-Aminopropylamino)acetate Derivatives

Preparation of di-BOC-protected diaminopropionic acid and methods **A** and **B** are described in ref. 31.

Synthesis of 2-(*N*-*tert*-Butoxycarbonyl-3-(*tert*-butyldimethylsilyloxy)propylamino)acetic acid is reported in the Supplemental Material.

Synthesis of (3-Hydroxypropylamino)acetate Derivatives

Typical procedure of method C. To a solution of cyclic α -hydroxyketone **2** (5.3 mg, 33 μ mol) in CH₂Cl₂ (2 ml) were added bromoacetyl bromide (20 μ l, 230 μ mol) and triethylamine (100 μ l, 1.0 mmol). The reaction mixture was stirred for 30 min. After the solvent had been removed, the residue was purified by preparative TLC (SiO₂) developed with *n*-hexane/ethyl acetate (2:1) to give bromoacetate derivative **8** (6.2 mg, 66%). To a solution of bromoacetate **8** (6.2 mg, 22 μ mol) in THF (1.0 ml) was added sodium carbonate (3.0 mg, 28 μ mol) followed by 3-aminopropanol (3.6 mg, 48 μ mol). The reaction mixture was stirred for 30 min and purified by silica gel column chromatography (ethyl acetate/methanol, 5:1) to give conjugate **2b** (5.0 mg, 81%).

Typical procedure of method D. To a solution of cyclic α -hydroxyketone **2** (4.0 mg, 24 μ mol) in CH₂Cl₂ (1.0 ml) were added {*tert*-BOC-[3-(*tert*-butyldimethylsilyloxy)propyl]amino}acetic acid (17 mg, 49 μ mol), EDC (11 mg, 58 μ mol), and DMAP (5.0 mg, 41 μ mol). The reaction mixture was stirred for 12 h at room temperature





Chart 2.



Scheme 1.

and then concentrated in vacuo. The crude product was purified by preparative TLC (SiO₂) with *n*-hexane/ethyl acetate (1:1) to give protected conjugate **11** (8.3 mg, 70%). To a solution of **11** (8.3 mg, 17 μ mol) in CH₂Cl₂ (1.0 ml) was added TFA (100 μ l) and stirred for 3 h at room temperature. The solvent and excess TFA were then removed under reduced pressure to give TFA salt **12** (6.6 mg). The salt was neutralized with sodium carbonate (10 mg) in methanol (1.0 ml). After removal of methanol in vacuo, conjugate **2b** was used in the complexation with the zinc porphyrin tweezer without any further purification.

Hydroxyconjugate 2b. ¹H NMR (300 MHz, CDCl₃) δ 7.87 (dd, J = 7.9, 1.7 Hz, 1 H), 7.50 (ddd, J = 7.7, 7.3, 1.7 Hz, 1 H), 7.06 (ddd, J = 7.9, 7.3, 0.9 Hz, 1 H), 6.99 (dd, J = 8.2, 0.5 Hz, 1 H), 5.71 (dd, J = 11.8, 5.6 Hz, 1 H), 4.58 (dd, J = 11.2, 5.6 Hz, 1 H), 4.41 (dd, J = 11.8, 11.2 Hz, 1 H), 3.80 (t, J = 5.3 Hz, 2 H), 3.59 (s, 1 H), 3.58 (s, 1 H), 2.92 (t, J = 5.8 Hz, 2 H), 1.73 (tt, J = 5.8, 5.3 Hz, 2 H); HRFABMS m/z 280.1163 [M + H]⁺, calcd. for C₁₄H₁₈O₅N, 280.1185.

Hydroxyconjugate 3b. ¹H NMR (300 MHz, CDCl₃) δ 8.00 (dd, J = 7.8, 1.1 Hz, 1 H), 7.51 (ddd, J = 7.5, 7.4, 1.1 Hz, 1 H), 7.33 (dd, J = 7.8, 7.5 Hz, 1 H), 7.26 (d, J = 7.8 Hz, 1 H), 5.60 (dd, J = 12.8, 5.4 Hz, 1 H), 3.85 (t, J = 5.3 Hz, 1 H), 3.68 (s, 1 H), 3.67 (s, 1 H), 3.3–3.0 (m, 2 H), 3.03 (t, J = 5.8 Hz, 2 H), 2.48–2.24 (m, 2 H), 1.82 (tt, J = 5.8, 5.3 Hz, 2 H); HRFABMS m/z 278.1396 [M + H]⁺, calcd. for C₁₅H₂₀O₄N, 278.1392.


Hydroxyconjugate 4b. ¹H NMR (300 MHz, CD₃OD) δ 7.65 (d, J = 7.7 Hz, 1 H), 7.48 (dd, J = 7.7, 7.3 Hz, 1 H),

7.34–7.28 (m, 2 H), 5.61 (dd, J = 11.2, 5.8 Hz, 1 H), 4.17 (s, 2 H), 3.71 (t, J = 5.7 Hz, 2 H), 3.23 (t, J = 5.5 Hz, 2 H), 3.18–2.97 (m, 2 H), 2.33 (m, 1 H), 2.22 (m, 1 H), 2.08–1.74 (m, 4 H); HRFABMS m/z 292.1554 [M + H]⁺, calcd. for C₁₆H₂₂O₄N, 292.1549.


Hydroxyconjugate 5b. ¹H NMR (300 MHz, CD₃OD) δ 5.38 (dd, J = 11.8, 6.6 Hz, 1 H), 4.11 (s, 2 H), 3.71 (t, J = 5.0 Hz, 2 H), 3.23 (t, J = 5.5 Hz, 2 H), 2.54–2.34 (m, 2 H), 2.22–2.09 (m, 2 H), 1.96–1.70 (m, 6 H); HRFABMS m/z 230.1400 [M + H]⁺, calcd. for C₁₁H₂₀O₄N, 230.1392.

Hydroxyconjugate 6b. ¹H NMR (300 MHz, CD₃OD) δ 5.47 (dd, J = 9.1, 5.5 Hz, 1 H), 4.07 (s, 2 H), 3.70 (t, J = 5.8 Hz, 2 H), 3.22 (t, J = 7.2 Hz, 2 H), 2.68 (ddd, J = 16.4, 5.4, 5.3 Hz, 1 H), 2.34 (ddd, J = 16.4, 5.4, 5.3 Hz, 1 H), 2.23–2.10 (m, 2 H), 1.95–1.70 (m, 8 H); HRFABMS m/z 244.1550 [M + H]⁺, calcd. for C₁₂H₂₂O₄N, 244.1549.

Aminoconjugate 3a. ¹H NMR (300 MHz, CD₃OD) δ 7.94 (dd, J = 8.3, 1.4 Hz, 1 H), 7.58 (ddd, J = 7.5, 7.4, 1.3 Hz,



Scheme 2.

Fig. 1. CD and UV spectra of **2** and **7** in MeCN (6.8×10^{-5} M).




Fig. 2. (a) Electric transition dipole moment (\leftrightarrow) of 3-methoxychroman-4-one calculated by ZINDO/S-CI. (b) Dihedral angle between electric transitions dipole moments in the lowest conformation of compound **7**.

1 H), 7.38–7.34 (m, 2 H), 5.75 (dd, $J = 14.6, 5.3 \text{ Hz}$, 1 H), 4.21 (s, 2 H), 3.44–3.14 (m, 4 H), 3.08 (dd, $J = 7.8, 7.7 \text{ Hz}$, 2 H), 2.53–2.45 (m, 1 H), 2.33 (ddd, $J = 12.6, 12.6, 4.9 \text{ Hz}$, 1 H), 2.13 (m, 2 H); HRFABMS m/z 277.1569 $[\text{M} + \text{H}]^+$, calcd. for $\text{C}_{15}\text{H}_{21}\text{O}_3\text{N}_2$, 277.1552.

Aminoconjugate 4b. ^1H NMR (300 MHz, CD_3OD) δ 7.66 (dd, $J = 8.0, 1.6 \text{ Hz}$, 1 H), 7.49 (ddd, $J = 7.5, 7.3, 1.5 \text{ Hz}$, 1 H), 7.35–7.31 (m, 2 H), 5.62 (dd, $J = 11.2, 5.9 \text{ Hz}$, 1 H), 3.21–3.16 (m, 2 H), 3.11–3.03 (m, 4 H), 2.38–2.15 (m, 4 H), 2.14–1.97 (m, 3 H), 1.92–1.75 (m, 1 H); HRFABMS m/z 291.1714 $[\text{M} + \text{H}]^+$, calcd. for $\text{C}_{16}\text{H}_{23}\text{O}_3\text{N}_2$, 291.1709.


Preparation of Host/Guest Complex for CD Measurement

Porphyrin tweezer **1** solution, 0.1 mM in anhydrous CH_2Cl_2 (10 μl), was added to methylcyclohexane (MCH) (1 ml). The exact concentration of the diluted tweezer **1** solution was determined by UV from the known ϵ value of the Soret band in methylcyclohexane ($\epsilon = 640,000 \text{ l mol}^{-1}$


cm^{-1}). To this solution was added 3 mM **2b** in anhydrous CH_2Cl_2 (10 μl). This solution was shaken, and UV and CD spectra were then recorded at room temperature and corrected by background subtraction. The CD spectra were measured in millidegrees and normalized into $[1 \text{ mol}^{-1} \text{ cm}^{-1}]/\lambda$ [nm] units.

RESULTS AND DISCUSSION

The aim of this paper is first to test the applicability of the conventional exciton chirality method to α -hydroxyketones and, second, to develop a more general protocol based on the porphyrin tweezer approach. For this purpose, we first attempted to apply the conventional approach of the exciton chirality method, namely, chromophoric derivatization of a functional group close to the stereogenic center, i.e., the 3-OH. Compound **2** was converted into its *p*-bromobenzoate **7** ($\lambda = 244 \text{ nm}$, $\epsilon = 19,500$ in EtOH) (Scheme 2). However, as shown in



Scheme 3.




Scheme 4.

Figure 1, benzoate **7** gave only a weak and ill-defined CD couplet at 258 nm ($\Delta\epsilon = -2.0$) and 240 nm ($\Delta\epsilon = +8.2$). A comparison with the CD of the original **2** (Fig. 1, dotted line) shows that it is impossible to interpret this very weak couplet between the chromanone ($\lambda = 250$ nm, $\epsilon = 4,500$ in MeCN) and *p*-bromobenzoate chromophores. In order to analyze this weak CD couplet we performed ZINDO/S-CI calculation of 3-methoxychroman-4-one, which showed the electric transition dipole around 262 nm to be oriented as depicted in Figure 2a. Furthermore, the dihedral angle between this dipole and that 4-bromobenzoate CT band in the optimized conformation of **7** found by MMFFs conformational search in chloroform (Fig. 2b) was calculated to be circa -151° . Thus, although the weak couplet between 258 nm ($\Delta\epsilon = -2.0$) and 240 nm ($\Delta\epsilon = +8.2$) originates from a negative exciton couplet between the two electric transition dipoles, its low intensity precludes application in configurational analysis.

In order to overcome the restriction due to such unsuitable CD coupling orientation between interacting dipoles, we chose the host-guest chiral recognition approach using dimeric zinc porphyrin tweezer **1** ($\lambda = 417$ nm, $\epsilon = 640,000$, methylcyclohexane) (Scheme 1).^{30–34} As shown in Scheme 3, it was attempted to convert α -

hydroxyketone **2** into its bifunctional ester conjugate **2a** according to methods **A** and **B**.³¹ In method **A**, alcohols are converted into bromoacetyl derivatives followed by reaction with diaminopropane to give aminoconjugate **2a**. Advantages of method **A** are that all reagents are commercially available and that the reactions are fast. The disadvantage, however, is that it is not applicable to compounds that are sensitive under basic conditions, due to the strong basicity of diaminopropane. On the other hand, in method **B**, di-BOC-protected diaminopropane ester is reacted with the hydroxyketone, followed by deprotection and neutralization to give the aminoconjugate. Although di-BOC-protected diaminopropionic acid has to be prepared, the aminoconjugate **10** is obtained under milder conditions. However, attempts to neutralize the TFA salt resulted in aminolysis, thereby regenerating the starting α -hydroxyketone **2** instead of the desired conjugate **2a** ("aminoconjugate"). With method **A**, the intermediate α -bromomethyl acetate **8** survived treatment with sodium carbonate in THF, but addition of



Scheme 5.












Fig. 3. CD and UV spectra of complex **1-2b** ($c = 1.1 \times 10^{-6}$ M in MCH).

TABLE 1. Predicted CD exciton chirality and observed CD amplitude (A_{CD}) of the complexes between zinc porphyrin tweezer **1** and conjugate compounds (**13a–16a** and **13b–16b**)

Parent alcohol	Predicted CD exciton chirality (MMFFs)	Conjugate compound	R = NH ₂ Preparation method A A_{CD}	R = OH Preparation method C A_{CD}
 (S)-13	Negative		13a –17	13b –17
 (S)-14	Positive		14a +40	14b +44
 (1R,2S,5R)-15	Negative		15a –494	15b –317
 (R)-16	Negative		16a –202	16b –198

1,3-diaminopropane resulted in its immediate decomposition to α -hydroxyketone **2**. With method **B**, TFA salt **9** was obtained as a stable compound, but neutralization with sodium carbonate in methanol also gave the α -hydroxyketone **2**. In both cases, the ester bonds in **8** and **9** are readily cleaved by inter- or intramolecular aminolysis which is well-documented as general base-catalysis arising from the 1,3-diaminopropane moiety.^{41,42} Therefore we

examined the employment of milder conjugates that are capable of coordinating with the zinc of the tweezer molecule, to prevent undesirable aminolysis.

Although oxygen nucleophilicity is weaker than that of nitrogen, it is known that coordination occurs between the substrate that contains a methyl ether oxygen and the zinc ion in the porphyrin.^{38,43} Thus the (3-hydroxypropylamino)acetyl group $\text{HOCH}_2\text{CH}_2\text{CH}_2\text{NHCH}_2\text{CO-}$ was




Fig. 4. (a) CD and UV spectra in MCH of the complex between the zinc tweezer and (a) conjugate **16a** ($c = 1.3 \times 10^{-6}$ M) and (b) conjugate **16b** ($c = 1.3 \times 10^{-6}$ M).














Fig. 5. Job plot for the complexation of Zn tweezer **1** with **16b**. The total concentration of two components was 10 μ M in MCH, with the molar fraction varying from 0 to 1.

tested as a new carrier moiety to produce “hydroxyconjugate” **2b** (Scheme 4). First, we applied method C, where 3-aminopropane is used instead of diaminopropane (see method A). An advantage of method C is that no racemization occurs because of the rapid formation of bromoacetate **8**. Addition of 3-aminopropanol to bromoacetate **8** in THF gave the desired products **2b** without ester bond hydrolysis. Method D was also tested. In this case, the carrier hydroxypropylamino moiety was introduced to give **11**, which was deprotected to TFA salt **12** and subsequently neutralized to yield the desired compound **2b**. Thus, both methods yielded the desired compound **2b**. The conjugate **2b** was then subjected to complexation with zinc porphyrin tweezer host **1**, giving rise to complex **1–2b** (Scheme 5). As shown in Figure 3, complex **1–2b** exhibited a clear-cut positive exciton couplet at 431 nm ($\Delta\epsilon = +10.5$) and 419 nm ($\Delta\epsilon = -8.7$), amplitude $A_{CD} = +19.2$. Furthermore, formation of the 1:1 complex between zinc porphyrin tweezer **1** and bidentate conjugate guest **2b** is supported by the shift of the broad UV/vis peak from 416 nm to the sharper 418 nm absorption (Fig. 3).³⁰

Because complex **1–2b** and those of **13b–16b** are a new type of host/guest complexes involving coordination between zinc and a primary alcohol function,

TABLE 2. Predicted CD exciton chirality and observed CD amplitude (A_{CD}) of the complexes between zinc porphyrin tweezer **1** and conjugate compounds (**2a–6a** and **2b–6b**)

Parent alcohol	Predicted CD exciton chirality (MMFFs)	Conjugate compound	R = NH ₂ Preparation method B A_{CD}	R = OH Preparation method D A_{CD}
 (S)- 2	Positive		2a —	2b +19.2
 (R)- 3	Negative		3a –283	3b –149
 (R)- 4	Negative		4a –180	4b –175
 (R)- 5	Positive		5a —	5b +137
 (R)- 6	Positive		6a —	6b +42.8




Fig. 6. Boltzmann distribution of θ values of the MC/MMFF calculated host-guest complex within 15 kJ/mol (3.6 kcal/mol) of the lowest-energy conformation for various conjugate (**13b**–**16b**).

their CD data were compared with those of conventional aminoconjugates **13a**–**16a** (Table 1). As shown, the sign of exciton-coupled CD and amplitudes (A_{CD}) of each set, aminoconjugates **13a**–**16a** and the corresponding hy-

droxyconjugates **13b**–**16b** were in good agreement. Note that in both cases of **1**–**13a** and **1**–**13b**, the zinc porphyrin tweezer recognizes even the subtle difference between methyl and ethyl groups ($A_{CD} = -17$). Although




Fig. 7. Front view of the most probable structure of complex **1**–**14b** calculated by MC/MMFFs. Complex **1**–**14b** was colored as follows: (pink) Zn-porphyrin (P-1) bound with primary alcohol; (yellow) Zn-porphyrin (P-2) bound with secondary amine; hydroxyconjugate **14b** is sky blue except for the primary alcohol (red) and secondary amine (dark blue).




Fig. 8. Boltzmann distribution of θ values of the MC/MMFFs calculated host–guest complex within 15 kJ/mol (3.6 kcal/mol) of the lowest-energy conformation for conjugate **2b**.

the UV/vis shift upon complexation in the hydroxyconjugate, from 416 to 419 nm, is smaller than that of the aminoconjugate, 416 and 422 nm, respectively, the

conformations of complexes between zinc porphyrin tweezer **1** and amino conjugates **13a–16a** are expected to be similar to those of hydroxyconjugates **13b–16b** (Fig. 4).

The Job plot performed for complex **1–16b**, where the relation between CD amplitude and molar fraction exhibited a maximum at molar fraction 0.5, suggests a 1:1 complexation between **1** and **16b** as previously reported for other tweezer complexes^{33,34} (Fig. 5).

After successful testing of hydroxyconjugates of model compounds **13b–16b**, the new conjugate system was applied to cyclic α -hydroxyketones (**3–6**), the conjugates being prepared according to methods shown in Table 2; in all cases, the hydroxyconjugates exhibited clear exciton CD. Interestingly, the negative exciton couplets of benzo-condensed cyclic α -hydroxyketones **3** and **4** opposite those of the corresponding cyclic α -hydroxyketones **5** and **6** (see below).

We also attempted to prepare the aminoconjugates of **3–6**. With aminoconjugates **3a** and **4a**, the A_{CD} was similar to those of the corresponding hydroxyconjugates **3b** and **4b**; however, the aminoconjugates of **5** and **6** could not be prepared.




Fig. 9. Boltzmann distribution of θ values of the MC/MMFF calculated host–guest complex within 15 kJ/mol (3.6 kcal/mol) of the lowest-energy conformation for various conjugates (**3b–6b**).

In previous studies, the assignment of absolute configurations from the observed CD couplet of the tweezer complex were mostly based on the relative steric sizes of the substituents at the stereogenic center, where the larger group protrudes from binding pocket while the medium group is clamped inside.^{30–32} However, in more complicated cases, this approach is not applicable. The Monte Carlo (MC) conformational search/Merck Molecular Force Field (MMFFs) molecular modeling⁴⁴ approaches developed more recently for predicting the porphyrin helicity in more sterically complicated substrates was found to be more favorable for α -hydroxyketone as well.^{33,35–37}

MMFFs calculations were thus performed in order to predict the porphyrin helicity in the complexes. As shown in Figure 6, the Boltzmann distribution of conformation with preferred porphyrin helicity (θ) of the complex was in good agreement with the observed sign of exciton-coupled CD. For example, the Boltzmann distribution of complex **1–13b** shows that 99% of the preferred conformations of complex **1–13b** adopt negatively arranged porphyrins (Fig. 6a), leading to a negative sign of the exciton-coupled CD, which is in agreement with the observed CD (Table 1). The lowest-energy conformation of complex **1–14b** is shown in Figure 7. Here the butyl group protrudes from the porphyrin sandwich to give a complex with positive helicity between the two porphyrin transition moments, 5–15 (P-1) and 5'–15' (P-2).⁴⁰

Similarly, the Boltzmann distribution of helicity (θ) between porphyrins in the complex **1–2b** was calculated using MMFFs. In agreement with experimental results shown in Figure 3, the calculation predicted a positive helicity (θ) between porphyrins in complex **1–2b** (Fig. 8). Furthermore, in complexes between hydroxyconjugate (**3b–6b**) and zinc porphyrin tweezer, the helicity based on MMFFs calculations also correctly predicted the observed results (Fig. 9).

Although the predicted porphyrin helicity by molecular modeling is in agreement with the observed CD exciton chirality, additional conformational studies aimed at clarification of the opposite interporphyrin helicity of **3a,b** vs. **5b** and of **4a,b** vs. **6b** are underway.

As mentioned above, in order to apply this method to α -hydroxyketones of unknown configurations, we prefer to make hydroxyconjugates according to **method D** and determine the configuration by comparison between the observed exciton coupled CD of the corresponding complex molecule and the results of the MMFFs calculations.

CONCLUSION

An efficient procedure for determination of absolute configurations of chiral cyclic α -hydroxyketones has been developed. The use of the (3-hydroxypropylamino)acetyl moiety as an improved carrier over the (3-aminopropylamino)acetyl moiety yields hydroxyconjugates that complex with the zinc porphyrin tweezer. This further extends the applicability of the tweezer method to cyclic α -hydroxycarbonyl systems for which there exists no general method. The absolute configurations of re-

sulting complexes can be determined from both the observed CD sign and from the predicted porphyrin helicity, as calculated by the Merck Molecular Force Field (MMFFs) method.

ACKNOWLEDGMENTS

This work was supported by the NIH (grants GM 34509 to K.N. and N.B.). We thank Dr. Katsunori Tanaka, Columbia University, for calculation of the electron transition of 3-methoxychroman-4-one using ZINDO/S-CI and Professor Gloria Proni, John Jay College, for helpful discussions and comments on the manuscript.

LITERATURE CITED

- Hanessian S. Organic chemistry series. Vol 3: Total synthesis of natural products: the "chiron" approach. New York: Pergamon Press; 1983. Chap 2, p 7–19.
- Oppolzer W. Asymmetric Diels–Alder and ene reactions in organic synthesis. *Angew Chem Int Ed Engl* 1984;23:876–889.
- Fujita M, Hiyama T. Highly diastereocontrolled reduction of ketones by means of hydrosilanes. Practical synthesis of optically active 1,2-diols and 2-amino alcohols of threo or erythro configuration. *J Am Chem Soc* 1984;106(16):4629–4630.
- Whitesell JK, Buchanan CM. Synthesis of (–)- and (+)-frontalin. *J Org Chem* 1986;51(26):5443–5445.
- Girijavallabhan VM, Ganguly AK, Pinto PA, Sarre OZ. Synthesis of the antifungal agent SCH 42427 (SM 9164). *Bioorg Med Chem Lett* 1991;1(7):349–352.
- Konosu T, Miyaoka T, Tajima Y, Oida S. Triazole antifungals. III. Stereocontrolled synthesis of an optically active triazolymethyl-oxirane precursor to antifungal oxazolidine derivatives. *Chem Pharm Bull* 1991;39(9):2241–2246.
- Chaudhary AG, Kingston DGI. Modified taxols. 11. Synthesis of 10-deacetoxytaxol and 10-deoxytaxotere. *Tetrahedron Lett* 1993;34(31):4921–4924.
- Gala D, DiBenedetto DJ, Clark JE, Murphy BL, Schumacher DP, Steinman M. Preparations of antifungal Sch 42427/MS 9164: preparative chromatographic resolution, and total asymmetric synthesis via enzymic preparation of chiral α -hydroxy arylketones. *Tetrahedron Lett* 1996;37(5):611–614.
- Vacca JP, Condra JH. Clinically effective HIV-1 protease inhibitors. *Drug Discov Today* 1997;2(7):261–272.
- Ghosh AK, Bilcer G, Schiltz G. Syntheses of FDA approved HIV protease inhibitors. *Synthesis* 2001(15):2203–2229.
- Askin D. The synthesis of indinavir and other clinically useful HIV-1 protease inhibitors. *Curr Opin Drug Discov Dev* 1998;1(3):338–348.
- Davies I, Taylor M, Marcoux J-F, Matty L, Wu J, Hughes D, Reider PJ. Stereoselective hydrogen bromide-promoted hydrogenation of an alpha-hydroxy oxime. *Tetrahedron Lett* 2000;41(42):8021–8025.
- Hansen KB, Rabbat P, Springfield SA, Devine PN, Grabowski EJJ, Reider PJ. Asymmetric synthesis of *cis*-aminochroman-ol. *Tetrahedron Lett* 2001;42(50):8743–8745.
- Lu Z, Raghavan S, Bohn J, Charest M, Stahlhut MW, Rutkowski CA, Simcoe AL, Olsen DB, Schleif WA, Carella A, et al. Design and synthesis of highly potent HIV protease inhibitors with activity against resistant virus. *Bioorg Med Chem Lett* 2003;13(10):1821–1824.
- Schock HB, Garsky VM, Kuo LC. Mutational anatomy of an HIV-1 protease variant conferring cross-resistance to protease inhibitors in clinical trials. Compensatory modulations of binding and activity. *J Biol Chem* 1996;271(50):31957–31963.
- Tata JR, Chapman KT, Duffy JL, Kevin NJ, Cheng Y, Rano TA, Zhang F, Huenig T, Kirk BA, et al., inventors; Merck & Co., Inc., USA, assignee. Preparation of γ -hydroxy-2-fluoroalkylaminocarbonyl-1-piperazinepentanamides as HIV protease inhibitors. *PCT Int Appl WO* 0138332, 2001.

17. Tata JR, Charest M, Lu Z, Raghavan S, Huening T, Rano T, inventors; Merck and Co., Inc., USA, assignee. Preparation of α -hydroxy- γ -[(carbocyclic- or heterocyclic-substituted)amino]carbonyl]alkanamides as HIV protease inhibitors. PCT Int Appl WO 0105230, 2001.
18. Tata JR, Raghavan S, Lu Z, Zhang F, Cheng Y, Chang J, Kim RM, Bohn JM, Rano T, Shen D-M, and others, inventors; Merck & Co., Inc., USA, and Huening T, assignees. Preparation of piperazine pentanamide HIV protease inhibitors and methods of treating AIDS. PCT Int Appl WO 0296359, 2002.
19. Marcune BF, Karady S, Reider PJ, Miller RA, Biba M, DiMichele L, Reamer RA. Asymmetric synthesis of cyclic hydroxy ketones derived from enol ethers via Sharpless asymmetric dihydroxylation. A study in the correlation of the enol ether chain length and enantioselectivity. *J Org Chem* 2003;68(21):8088–8091.
20. Hashiyama T, Morikawa K, Sharpless KB. α -Hydroxy ketones in high enantiomeric purity from asymmetric dihydroxylation of enol ethers. *J Org Chem* 1992;57(19):5067–5068.
21. Adam W, Diaz MT, Fell RT, Saha-Moeller CR. Kinetic resolution of racemic α -hydroxy ketones by lipase-catalyzed irreversible transesterification. *Tetrahedron: Asymmetry* 1996;7(8):2207–2210.
22. Borgevig A, Sunden H, Cordova A. Direct catalytic enantioselective α -aminooxylation of ketones: a stereoselective synthesis of α -hydroxy and α,α' -dihydroxy ketones. *Angew Chem Int Ed* 2004;43(9):1109–1112.
23. Lee LG, Whitesides GM. Preparation of optically active 1,2-diols and α -hydroxy ketones using glycerol dehydrogenase as catalyst: limits to enzyme-catalyzed synthesis due to noncompetitive and mixed inhibition by product. *J Org Chem* 1986;51:25–36.
24. Momiyama N, Yamamoto H. Catalytic enantioselective synthesis of α -aminooxy and α -hydroxy ketone using nitrosobenzene. *J Am Chem Soc* 2003;125:6038–6039.
25. Masui M, Ando A, Shiojiri T. New methods and reagents in organic synthesis. 75. Asymmetric synthesis of α -hydroxy ketones using chiral phase transfer catalysts. *Tetrahedron Lett* 1988;29:2835–2838.
26. Cain CM, Cousins RPC, Coumbarides G, Simpkins NS. Asymmetric deprotonation of prochiral ketones using chiral lithium amide bases. *Tetrahedron* 1990;46:523–544.
27. Specht KM, Nam J, Ho DM, Berova N, Kondru RK, Beratan DN, Wipf P, Pascal Jr RA, Kahne D. Determining absolute configuration in flexible molecules: a case study. *J Am Chem Soc* 2001;123:8961–8966.
28. Momiyama N, Yamamoto H. Catalytic enantioselective synthesis of α -aminooxy and α -hydroxy ketone using nitrosobenzene. *J Am Chem Soc* 2004;126:6498.
29. Hayashi Y, Yamaguchi J, Sumiya T, Hibino K, Shoji M. Direct proline-catalyzed asymmetric α -aminooxylation of aldehydes and ketones. *J Org Chem* 2004;69:5966–5973.
30. Huang X, Borhan B, Rickman BH, Nakanishi K, Berova N. Zinc porphyrin tweezer in host–guest complexation: determination of the absolute configurations of primary monoamines by circular dichroism. *Chem Eur J* 2000;6:216–224.
31. Kurtan T, Nesnas N, Li Y-Q, Huang X, Nakanishi K, Berova N. Chiral recognition by CD-sensitive dimeric zinc porphyrin host. 1. Chiroptical protocol for absolute configurational assignments of monoalcohols and primary monoamines. *J Am Chem Soc* 2001;123:5962–5973 and references therein.
32. Kurtan T, Nesnas N, Li Y-Q, Koehn FE, Nakanishi K, Berova N. Chiral recognition by CD-sensitive dimeric zinc porphyrin host. 2. Structural studies of host–guest complexes with chiral alcohol and monoamine conjugates. *J Am Chem Soc* 2001;123:5974–5982.
33. Huang X, Fujioka N, Pescitelli G, Koehn FE, Williamson RT, Nakanishi K, Berova N. Absolute configurational assignments of secondary amines by CD-sensitive dimeric zinc porphyrin host. *J Am Chem Soc* 2002;124:10320–10335.
34. Proni G, Pescitelli G, Huang X, Nakanishi K, Berova N. Magnesium tetraarylporphyrin tweezer: a CD-sensitive host for absolute configurational assignments of α -chiral carboxylic acids. *J Am Chem Soc* 2003;125(42):12914–12927.
35. Solladie-Cavallo A, Roje M, Giraud-Roux M, Chen Y, Berova N, Sunjic V. *trans*-Diaryl epoxides: asymmetric synthesis, ring-opening, and absolute configuration. *Chirality* 2004;16:196–203.
36. Ishii H, Krane S, Itagaki Y, Berova N, Nakanishi K, Weldon PJ. Absolute configuration of a hydroxyfuranoid acid from the pelage of the genus *Bos*, 18-(6S,9R,10R)-bovicidic acid. *J Nat Prod* 2004;67:1426–1430.
37. Van Klink JW, Baek S-H, Barlow AJ, Ishii H, Nakanishi K, Berova N, Perry NB, Weavers RT. Determination of the absolute configuration of anisotome irregular diterpenes: application of CD and NMR methods. *Chirality* 2004;16:549–558.
38. Teo T-L, Vetrichelvan M, Lai Y-H. Infinite three-dimensional polymeric metalloporphyrin network via six-coordinate Zn(II) and two axial oxygen donors. *Org Lett* 2003;5(22):4207–4210.
39. Jones GIL, Owen NL. Molecular structure and conformation of carboxylic esters. *J Mol Struct* 1973;18(1):1–32.
40. Pescitelli G, Gabriel S, Wang Y, Fleischhauer J, Woody RW, Berova N. Theoretical analysis of the porphyrin–porphyrin exciton interaction in circular dichroism spectra of dimeric tetraarylporphyrins. *J Am Chem Soc* 2003;125(25):7613–7628.
41. Bruice TC, Willis RG. The reaction of aliphatic diamines with phenyl acetate. *J Am Chem Soc* 1965;87(3):531–536.
42. Rajarathnam D, Jeyakumar T, Nadar PA. Structure-reactivity correlation in the aminolysis of 4-fluorophenyl acetate in aqueous medium. *Int J Chem Kinet* 2002;34(6):366–373.
43. Senge MO, Speck M, Wiehe A, Dieks H, Aguirre S, Kurreck H. Structure and conformation of photosynthetic pigments and related compounds. 12. A crystallographic analysis of porphyrin–quinones and their precursors. *Photochem Photobiol* 1999;70(2):206–216.
44. Halgren TA. Merck molecular force field. I. Basis, form, scope, parameterization, and performance of MMFF94. *J Comput Chem* 1996;17:490–519.