

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/6387840>

Regional multilineage differentiation potential of meniscal fibrochondrocytes: Implications for meniscus repair

ARTICLE *in* THE ANATOMICAL RECORD ADVANCES IN INTEGRATIVE ANATOMY AND EVOLUTIONARY BIOLOGY · JANUARY 2007

Impact Factor: 1.54 · DOI: 10.1002/ar.20419 · Source: PubMed

CITATIONS

43

READS

17

4 AUTHORS, INCLUDING:



Gabriel J Martinez-Diaz

University of Pittsburgh

16 PUBLICATIONS 145 CITATIONS

SEE PROFILE



Rocky S Tuan

University of Pittsburgh

505 PUBLICATIONS 23,092 CITATIONS

SEE PROFILE

Regional Multilineage Differentiation Potential of Meniscal Fibrochondrocytes: Implications for Meniscus Repair

ROBERT L. MAUCK, GABRIEL J. MARTINEZ-DIAZ, XIAONING YUAN,
AND ROCKY S. TUAN*

Cartilage Biology and Orthopaedics Branch, National Institute of Arthritis and
Musculoskeletal and Skin Diseases, National Institutes of Health, Department of Health
and Human Services, Bethesda, Maryland

ABSTRACT

The knee menisci are wedge-shaped semilunar fibrocartilaginous structures that reside between the femur and tibia and function to transmit and distribute load. These structures have characteristics of both fibrous and cartilaginous tissues. The cartilage-like inner region and the fibrous vascularized outer region each has a distinct extracellular matrix, and resident meniscal fibrochondrocytes (MFCs) with distinct morphologies dependent on their location. Damage to the meniscus is common, and disruption of tissue structure and function result in erosion of the underlying articular cartilage. It has been observed that damage in the vascular periphery undergoes spontaneous repair, whereas damage of the inner region does not heal. While vascularity of the peripheral region plays a role in healing, recent findings have also suggested that local cellular composition influences local healing capacity. This study examined the variation in multipotential characteristics of cell populations isolated from different regions of the bovine meniscus. MFCs were isolated from the outer (vascular), inner (avascular), and horn (mixed) regions and induced toward chondrogenic, adipogenic, and osteogenic lineages. The results of this study suggest that MFCs from all regions of the meniscus possess a multilineage differentiation capability, particularly toward chondrogenesis and adipogenesis. MFCs from the outer region were most plastic, differentiating along all three mesenchymal lineages. These findings may underlie the experimental observation of improved integration of meniscus grafts from the outer zone and may have implications for developing strategies of cell-based meniscus repair. *Anat Rec*, 290:48–58, 2007.

© 2006 Wiley-Liss, Inc.

Key words: meniscus; fibrochondrocyte; multipotential mesenchymal stem cells; adipogenesis; osteogenesis; chondrogenesis

Grant sponsor: Intramural Research Program, National Institute of Arthritis and Musculoskeletal and Skin Diseases, National Institutes of Health; Grant number: ZO1 AR41113.

Robert L. Mauck's current address is: McKay Orthopaedic Research Laboratory, Department of Orthopaedic Surgery, University of Pennsylvania, Philadelphia, Pennsylvania.

*Correspondence to: Rocky S. Tuan, Cartilage Biology and Orthopaedics Branch, National Institute of Arthritis and Muscu-

loskeletal and Skin Diseases, National Institutes of Health, 50 South Drive, MSC 8022, Building 50, Room 1503, Bethesda, MD 20892. Fax: 301-480-4315. E-mail: tuanr@mail.nih.gov

Received 31 May 2006; Accepted 8 October 2006

DOI 10.1002/ar.a.20419

Published online in Wiley InterScience
(www.interscience.wiley.com).

The knee menisci are wedge-shaped semilunar fibrocartilaginous structures that reside between the femur and tibia (Rath and Richmond, 2000). The main function of the meniscus is to transmit and distribute load, increase joint congruency, and stabilize the joint (Ghosh and Taylor, 1987; Arnoczky et al., 1988a). As a fibrocartilaginous structure, the meniscus has characteristics of both fibrous and cartilaginous tissues, with the inner region more cartilage-like (O'Connor, 1976) and the outer region more fibrous (Petersen and Tillmann, 1998). The meniscus is predominantly composed of collagen (85–95% of the dry weight), of which > 90% is collagen type I, with the remaining 1–2% consisting of collagen types II, III, V, and VI (Eyre and Wu, 1983; Cheung, 1987; McDevitt and Webber, 1990; Kambic and McDevitt, 2005). Proteoglycans make up less than 2–3% of the dry weight (Fithian et al., 1990; McDevitt and Webber, 1990; Adams and Hukins, 1992; Buma et al., 2004) and are concentrated in the inner cartilaginous zones. In the outer fibrous region of the meniscus, collagen fibers are circumferentially oriented (Bullough et al., 1970; Arnoczky et al., 1988a; Mow et al., 1992; Petersen and Tillmann, 1998), resulting in highly anisotropic material properties that are greatest in the fiber direction (Bullough et al., 1970; Setton et al., 1999). These properties are essential for the mechanical function of the tissue.

The cellular component of the meniscus reflects its fibrocartilaginous nature. The main cell type in the structure is the meniscal fibrochondrocyte (MFC), which sparsely populates the tissue (McDevitt and Webber, 1990; Adams and Hukins, 1992; Setton et al., 1999; Benjamin and Ralphs, 2004). The appearance and phenotypic characteristics of these MFCs vary according to their position in the meniscus (Hellio Le Graverand et al., 2001a). For example, MFCs located in the inner avascular region resemble articular chondrocytes; they are rounded and well separated from one another by their surrounding extracellular matrix (ECM). Conversely, MFCs in the outer fibrous region are spindle-shaped, with numerous projections that form gap junctions with neighboring cells (Hellio Le Graverand et al., 2001a). Collectively, MFCs function to maintain and remodel the meniscus ECM (McDevitt and Webber, 1990; Adams and Hukins, 1992) and alter their biosynthetic activities in response to a changing mechanical environment (Shin et al., 2003; Upton et al., 2003) and/or onset of joint instability (Hellio Le Graverand et al., 2001b, 2001c). MFCs can be isolated directly from the meniscus via digestion of the ECM, or coaxed to migrate out of meniscus pieces in tissue culture (Webber et al., 1985, 1989; Webber, 1990).

It has long been appreciated that damage to the meniscus is a common occurrence (MacAusland, 1931; Cravener and MacElroy, 1941), with indications of scarring (indicating past insult) in most patients over the age of 45. Damage can be of many forms, including flap and horn tears, longitudinal tears, or transverse or oblique tears (MacAusland, 1931; Cravener and MacElroy, 1941). Not all meniscus tears are considered treatable, and partial resection is more common than repair (Rath and Richmond, 2000). Mechanical compromise of the meniscus decreases tibiofemoral contact area and thereby increases stress concentrations on the underlying cartilage (Krause et al., 1976; Baratz et al., 1986; Ahmed, 1992; Rath and Richmond, 2000) and changes joint motion (Levy et al., 1982; Lanzer and Komenda, 1990). In several animal models (Hoch et al., 1983; Elliott et al.,

1999; LeRoux et al., 2000), meniscectomy led to a decrease in the mechanical properties of the articular cartilage, with eventual degenerative changes, such as osteophyte formation and joint space narrowing (Petrosini and Sherman, 1996; Rath and Richmond, 2000).

One of the interesting features of meniscus healing arises from the finding that the tissue is differentially vascularized, with a vascular periphery (outer one-third) and an avascular (cartilaginous) inner region (Arnoczky and Warren, 1983). King (1936a, 1936b) was the first to suggest that tears connected to the vascular periphery undergo spontaneous repair, whereas those limited to the inner region do not (Petrosini and Sherman, 1996). This observation has led to the belief that vascularity is essential for healing, and consequently, surgical techniques have been designed to improve vascular access to sites of meniscus damage (Arnoczky and Warren, 1983). These methods include the introduction of vascular channels or the implantation of fibrin clots (Arnoczky et al., 1985, 1988b; Arnoczky, 1992). More recently, it has been observed that different zones of the meniscus have differing healing potentials, even in the absence of an active blood supply (Kobayashi et al., 2004). In these studies of whole meniscus explants, portions of the vascular region were transplanted to freshly prepared defects in the avascular region and found to integrate better than grafts derived from the avascular region. These findings suggest that the endogenous cellular composition of the meniscus may play a role in the local healing response.

It is generally believed that endogenous tissue repair is driven by progenitor cells located within the tissue that migrate to injury sites, proliferate, and deposit new matrix (Barry, 2003). Indeed, an increasing number of musculoskeletal tissues have been found to harbor cells that retain some multipotential characteristics, including adipose tissue (Zuk et al., 2001; Erickson et al., 2002), articular cartilage (Barbero et al., 2003; Tallheden et al., 2003; Alsalameh et al., 2004), trabecular bone (Osyczka et al., 2002; Tuli et al., 2003), synovium (Nishimura et al., 1999; De Bari et al., 2003), heart and skeletal muscle (Warejcka et al., 1996; Williams et al., 1999; Young et al., 2001), and periodontal ligaments (Lekic and McCulloch, 1996; Seo et al., 2005). In this study, we examined the ability of MFCs to undergo multipotential lineage differentiation (chondrogenesis, adipogenesis, and osteogenesis). These differentiation lineages were assessed in comparison to bone marrow-derived MSCs, a cell source known to exhibit such multipotential characteristics (Caplan, 1991; Prockop, 1997; Pittenger et al., 1999). Furthermore, to determine if there exists a regional variation in this cellular component of endogenous meniscus repair, this study examined the potentiality of MFCs isolated from different regions: the outer (vascular), inner (avascular), and horn (mixed) regions. We hypothesized that MFCs derived from the outer vascular region would possess a multilineage differentiation potential, while those from the avascular inner zones would not, with inherent cell plasticity reflecting the overall healing capacity of each region.

MATERIALS AND METHODS

Meniscal Fibrochondrocyte Isolation and Expansion

MFCs were isolated from the medial and lateral menisci of 3- to 6-month-old bovine calves within 36 hr of slaughter (Fresh Farms Beef, Rutland, VT). For each of

four replicate experiments, the menisci of 3–5 donors were isolated and combined. After removal of any extraneous soft capsular tissue, menisci were sectioned into an inner (I; inner one-third of central region), outer (O; outer one-third of central region), and horn (H; anterior and posterior full thickness) regions. These sections were manually diced into 1–2 mm³ pieces and plated in 10 cm tissue culture dishes with 40 mL of a basal medium consisting of high-glucose (hg) DMEM containing 1 × penicillin/streptomycin/fungizone (PSF) and 10% fetal bovine serum (FBS). MFCs were observed to emerge from meniscus pieces over a 1- to 2-week period, after which they were trypsinized, replated, and expanded through passage 2 (p2). Human bone marrow-derived mesenchymal stem cells (MSCs; p3 and p4) were isolated and expanded as described previously (Song and Tuan, 2004) according to an institutional review board-approved procedure (George Washington University) and used as a positive control for differentiation assays. For all studies, media were changed twice weekly over the 21-day culture period.

Multilineage Differentiation Culture Conditions and Analysis

Chondrogenesis. To assess the ability of meniscus-derived cells to undergo chondrogenesis, isolated MFCs and human MSCs were formed into pellets containing 250,000 cells in 96-well polypropylene plates as described previously (Mauck et al., 2006). In a preliminary study, cells from the entirety of the meniscus were isolated, while in subsequent studies cells from distinct regions were utilized. For long-term culture, pellets were maintained in either basal medium (BM) supplemented with 50 µg/mL ascorbate 2-phosphate or a chemically defined chondrogenic medium consisting of hgDMEM supplemented with 1 × PSF, 0.1 µM dexamethasone, 50 µg/mL ascorbate 2-phosphate, 40 µg/mL L-proline, 100 µg/mL sodium pyruvate, 1 × insulin/transferrin/selenium (ITS)⁺ Premix (Becton Dickinson, Franklin Lakes, NJ) with (CM⁺) or without (CM[−]) 10 ng/mL transforming growth factor β3 (TGF-β3) (R&D Systems, Minneapolis, MN). At weekly intervals through day 21, several pellets from each group were removed from culture and either suspended in TRIzol reagent (Invitrogen, Carlsbad, CA) and stored at −80°C for RNA isolation or embedded in 2% low-melting-temperature agarose and fixed overnight at 4°C in 4% phosphate-buffered paraformaldehyde. These pellet-laden agarose blocks were paraffin-embedded and sectioned to 8 µm thickness. Sections were stained with hematoxylin and eosin (H&E), Picrosirius red, or Alcian blue (pH 1.0; Rowley Biochemicals, Danvers, MA) to visualize cells, bulk collagen, and sulfated proteoglycan, respectively. Additionally, sulfated glycosaminoglycan (s-GAG) content was determined using the DMMB dye-binding assay (Farndale et al., 1986) on a per-pellet basis for 6–8 pellets/replicate on papain-digested samples (as in Mauck et al., 2003).

Adipogenesis and osteogenesis. To test for adipogenesis and osteogenesis, p2 MFCs derived from each region as well as human MSCs were plated at 20,000 cells/cm² in tissue culture-treated six-well plates and maintained in lineage-specific differentiation media (3 mL/well). Freshly prepared differentiation medium was changed twice weekly. Osteogenic medium (OS) consisted of DMEM

supplemented with 10% FBS, 1 × PSF, 10 nM dexamethasone, 10 mM β-glycerophosphate, 50 µg/mL ascorbate 2-phosphate, and 10 nM 1,25-dihydroxyvitamin D3. Adipogenic medium (AS) consisted of DMEM supplemented with 10% FBS, 1 × PSF, 1 µM dexamethasone, 1 µg/mL insulin, and 0.5 mM 3-isobutyl-1-methylxanthine (IBMX). Control cultures were maintained in basal medium.

On days 1 and 21, relative cell numbers in OS, AS, and control cultures were determined using the MTT Cell Proliferation Assay (Molecular Probes, Eugene, OR). A further sampling for each condition was collected in 1 mL of TRIzol reagent (Invitrogen) and stored at −80°C for RNA extraction. On day 21, OS and control samples were fixed with 60% isopropanol and stained with Alizarin red (2%; Rowley Biochemicals) for mineralized matrix. Day 21 AS and control samples were fixed with 4% phosphate-buffered paraformaldehyde (FD Neurotechnologies, Catonsville, MD) and stained with Oil Red O (Sigma Chemicals, St. Louis, MO; dissolved at 0.18% in 60% isopropanol) to visualize lipid droplets. Oil Red O content was further quantified by elution with isopropanol and absorbance measurements at OD 540 nm (Sen et al., 2001). For each replicate study, 1–2 wells from Oil Red O-stained cultures were counterstained with hematoxylin (Sigma Chemicals; to visual cell nuclei) and photographed using an inverted microscope.

One-Step Reverse Transcriptase and Real-Time PCR

For undifferentiated MFC monolayers cultures, as well as those maintained in adipogenic or osteogenic culture, total RNA was extracted via mechanical disruption in TRIzol reagent followed by phenol-chloroform phase separation. For chondrogenic pellets, total RNA was extracted from pellets by mechanical dissociation with a motorized pestle (Kimble/Kontes, Vineland, NJ) in 1 mL of TRIzol reagent. RNA purity and quantity was assessed spectrophotometrically on the basis of A_{260/280}. In preliminary chondrogenesis studies using cells from the entirety of the meniscus, eight pellets were harvested on days 7 and 21. These samples were analyzed for transcript levels of *aggrecan* (AGN) and *cartilage oligomeric matrix protein* (COMP) using the One-Step Reverse Transcriptase PCR kit (Invitrogen) with primers and amplification conditions as defined previously (Li et al., 2003). For assessment of regional variation in chondrogenesis, 12–16 pellets were harvested per replicate study on day 21. Reverse transcription was performed using the First Strand cDNA Synthesis kit (Invitrogen) with 1 µg of total RNA using random hexamers according to the manufacturer's instructions. cDNA amplification was carried out using an iCycler real-time PCR system (Bio-Rad, Hercules, CA) using SYBR Green PCR Reaction Mix (Bio-Rad). Starting quantities of target gene transcripts in unknowns were derived from corresponding standard curves for each primer pair, with overall expression normalized to *GAPDH* in each sample. For AS and OS cultures and controls, expression of *fatty acid binding protein* (FABP) and *alkaline phosphatase* (ALP) were assessed (Baksh et al., 2004), respectively. For chondrogenic conditions, the expression of the ECM genes, *AGN*, *collagen types I and II*, were assessed (Fitzgerald et al., 2004).

Statistical Analysis

Analysis of variance (ANOVA) was performed using Statistica (Statsoft, Tulsa, OK), with Fisher's LSD post-




Fig. 1. Chondrogenic differentiation of day 21 pellet cultures of MFCs isolated from the entirety of the bovine meniscus. Medium composition: BM, basal medium; CM⁺, chondrogenic medium with TGF- β 3; CM⁻, chondrogenic medium without TGF- β 3. **A:** Histology: staining with H&E (left), Alcian blue (middle), and Picrosirius red (right). CM⁺ cultures showed a dense collagenous matrix (right) that is also rich in proteoglycans (middle). In the absence of TGF- β 3 (CM⁻), collagenous ma-

trix (right) is deposited with little proteoglycan (middle). In BM, small pellets formed with little ECM deposition. Dashed line highlights pellet periphery in CM⁻ and BM conditions. Scale bar = 50 μ m. **B:** RT-PCR analysis of cartilaginous matrix gene expression. An enhanced chondrogenic phenotype is seen, with higher *aggrecan* and *COMP* expression, in pellet cultures maintained in CM⁺ compared to either CM⁻ or BM on days 7 and 21.

hoc testing between groups. Independent variables were meniscus zone (inner, outer, or horns) and culture condition (CM⁻/CM⁺, AS, OS, and control), with dependent variables including wet weight, s-GAG content, cell activity (MTT), and Oil Red O content. All data are reported as the mean and standard deviation of 4–15 samples derived from a minimum of three replicate studies, depending on the assay, with significance set at $P < 0.05$.

RESULTS

Chondrogenesis

In preliminary studies, culture of MFCs derived from the entirety of the meniscus in pellet format resulted in the accumulation of a proteoglycan-and-collagen-rich

ECM in CM⁺ medium. In CM⁻, a dense collagenous matrix was formed that was generally devoid of Alcian blue staining, indicating a lack of cartilage matrix accumulation. When maintained in BM for 21 days, small cellular pellets formed with little ECM (Fig. 1A). Gene expression corroborated these observations showing that CM⁺ medium sustained and/or enhanced expression of *aggrecan* and *COMP*, two markers of the chondrocyte phenotype, while BM and CM⁻ conditions did not (Fig. 1B).

We next examined MFCs derived from distinct regions of the meniscus. The initial morphologies of these cells were different; after 5 days of outgrowth from meniscus pieces, inner zone cells remained slightly rounded while those from the outer zone took on a spindle shape (Fig. 2). After monolayer expansion, culture of MFCs in pellets in CM⁺ led to pronounced cartilaginous ECM production in




Fig. 2. Isolation of fibrochondrocytes from three distinct regions of the meniscus: the inner avascular body, the outer vascular body, and the mixed vascular horns. Cells were isolated via outgrowth from small pieces of each region (top right). Primary meniscal fibrochondrocytes in monolayer culture (day 5) showed distinct morphologies (bottom). Scale bar = 100 μ m.




Fig. 3. Chondrogenic differentiation in day 21 pellet cultures of MFCs isolated from different regions of bovine meniscus compared to that in human MSCs maintained under identical CM^+ conditions. All cultures showed pronounced chondrogenic differentiation, with human

MSCs showing more homogeneous distribution of cartilaginous ECM. H&E (top row), Picrosirius red (middle row), and Alcian blue (bottom row). Scale bar = 50 μ m.

cells from all regions, particularly in the pellet periphery (Fig. 3). Real-time RT-PCR analysis of gene expression demonstrated a marked induction of *aggrecan* (2- to 20-fold) and *collagen type II* (8- to 180-fold) in all MFC populations compared to monolayers, with no distinct differences in cell response observed between groups (Fig. 4). *Collagen type I* gene expression was high for all populations, did not change in pellet cultures compared to monolayers with culture in CM^+ , and remained much

higher (1–2 orders of magnitude) than collagen type II expression levels for these same pellets. Over a 21-day period, pellet wet weights increased significantly in CM^+ compared to those in CM^- for cells from each zone of the meniscus ($P < 0.01$; $n = 4$). On day 21, s-GAG content increased with CM^+ for each group as well ($P < 0.001$; $n = 4$), with no significant differences observed between groups (I: 54 ± 8 ; O: 66 ± 5 ; H: 57 ± 7 ; MSC: 61 ± 3 μ g s-GAG/pellet).




Fig. 4. Expression of cartilage matrix genes, and wet weight and s-GAG content of day 21 pellet cultures MFCs isolated from different regions of bovine meniscus. **A–C:** Real-time RT-PCR analysis showing fold change in expression in *collagen type I*, *aggrecan*, and *collagen type II* (compared to monolayer) for pellets maintained in chondrogenic (CM⁺) conditions (data from three replicate experiments). Cartilage

matrix genes, *collagen type II* and *aggrecan*, are both upregulated, whereas *collagen type I* levels remain unchanged. Wet weight (**D**) and s-GAG content (**E**; per pellet) of pellets from four replicate experiments cultured in either CM⁺ (black circles) or CM⁻ (gray circles). All chondrogenic cultures showed increased weight and s-GAG content. Asterisk indicates $P < 0.05$ for CM⁺ pellets vs. CM⁻ pellets.




Fig. 5. Adipogenic differentiation of day 21 monolayer cultures of MFCs isolated from each region of the meniscus maintained in adipogenic medium and examined by histological staining. AS, adipogenic medium; CT, control in basal medium. AS cultures of all MFCs showed

intracellular deposition of lipid droplets as shown by Oil Red O staining with hematoxylin counterstain. Bone marrow-derived human MSCs were also similarly cultured and stained as a positive control. Scale bar = 50 μm.

Adipogenesis

Monolayer cultures of MFCs derived from each region of the meniscus were treated with adipogenic supplements (AS) for 21 days. Compared to controls, MFCs in AS conditions increased their expression of *FABP* by 10- to 100-fold (data not shown), with no clear differences observed among cells from the different meniscus regions. Staining with

Oil Red O showed positive lipid droplet accumulation in cells derived from all regions of the meniscus (Fig. 5). Quantification of bound dye demonstrated significant increases in AS cultures of cells from every region, with significantly higher levels in outer zone and horn MFCs maintained in AS cultures compared to inner MFCs similarly maintained ($P < 0.05$; $n = 15$; Fig. 6). In these cultures, cell number increased by four- to sevenfold in both




Fig. 6. Adipogenic differentiation of day 21 monolayer cultures of MFCs isolated from each region of the meniscus maintained in adipogenic medium. AS, adipogenic medium; CT, control in basal medium. Top: Quantification of Oil Red O-stained lipid droplet accumulation. AS cultures showed significantly higher dye content, particularly in outer

and horn MFCs (asterisk, $P < 0.001$ compared to CT; double dagger, $P < 0.03$ compared to CT and inner AS; $n = 9-15$). Bottom: Day 21 cell numbers normalized to initial cell number (dotted line). In all cultures, cell numbers increased four- to sevenfold. Asterisk, $P < 0.05$ compared to CT ($n = 6-9$). Data are from three replicate studies.

media, with significantly more cells resulting from AS treatment of inner MFCs compared to its control ($P < 0.05$; $n = 9$).

Osteogenesis

MFCs from each region of the meniscus were also maintained in monolayer culture and treated with osteogenic supplements (OS) for 21 days. In these cultures, cell number increased by five- to eightfold, with significantly more cells in OS-treated samples compared to control cultures for cells from all regions ($P < 0.04$; $n = 6-9$; Fig. 7), with outer zone and horn regions under OS condition having significantly more cells on day 21 compared to the inner region ($P < 0.05$). Osteogenesis under OS culture condition was indicated by increased *ALP* gene expression in both inner and outer MFCs, with a small decrease in horn MFCs (fold difference, I: 1.9; O: 3.5; H: 0.8; $n = 3$). Culture of MFCs under OS condition also showed a marked increase in mineralized matrix deposition, specifically in cells derived from the outer region of the meniscus. Minimal staining for matrix mineralization was observed in cultures of MFCs derived from the inner or horn regions (Fig. 7).

DISCUSSION

The results of this study demonstrate that culture-expanded meniscal fibrochondrocytes possess a marked

multilineage differentiation potential, including the capacity to undergo chondrogenesis, adipogenesis, and osteogenesis. To better understand the origin of these multipotential cells, and their relationship to meniscus repair, we isolated MFCs from the inner avascular, outer vascular, and mixed horn region of the meniscus. We hypothesized that a multipotential characteristic would be apparent only in cells derived from the outer region, where some degree of endogenous meniscus repair occurs. Of the three regions investigated, cells from the outer vascular zone did show the greatest capacity to differentiate along all three lineages examined. These findings suggest that the cells of the outer meniscus have a wide phenotypic range, a potential explanation for the increased integration observed when explants from this region are transplanted to avascular regions of the tissue (Kobayashi et al., 2004). Counter to our initial hypothesis, MFCs from the inner avascular region of the meniscus also showed some multipotential capacity, particularly toward the adipogenic and chondrogenic lineages. These inner MFCs, which show a more cartilaginous phenotype in situ (Sweigart and Athanasiou, 2001), had a reduced osteogenic potential when compared to MFCs from the outer region. In light of the reduced capacity of the inner regions to mount a reparative response in vivo, this diminished multipotential capacity of inner MFCs may account for some of the limitations in endogenous repair.

Given the wide array of tissues found to house cells that possess a multilineage differentiation potential, some con-




Fig. 7. Osteogenic differentiation of day 21 monolayer cultures of MFCs isolated from each region of the meniscus maintained in osteogenic medium. OS, osteogenic medium; CT, control in basal medium. Top: Day 21 cell numbers normalized to initial cell number (dotted line). In all cultures, cell numbers increased five- to eightfold, with higher levels seen in cells from the outer and horn regions (asterisk, $P < 0.05$

compared to CT; double dagger, $P < 0.05$ compared to CT and inner OS; $n = 6-9$). Bottom: Alizarin red staining of mineralized matrix in cultures of MFCs. High level of matrix mineralization is seen in OS cultures of MFCs isolated from the outer region of the meniscus. Bone marrow-derived human MSCs similarly cultured were also stained as positive control. Scale bar = 500 μm . Data are from three replicate studies.

sideration should be given to the nature of the adult stem cell. It has been suggested that the conceptualization of the a stem cell most accurately refers to a biological function, rather than a distinct cell type (Blau et al., 2001). It is further suggested that this functional capacity is not necessarily equal among all cells, but is likely present in most. In this study, MFCs from the meniscus were observed to possess sufficient plasticity to distinguish them as putative stem cells, i.e., they were capable of a minimum of three different lineage-specific differentiation profiles when presented with the appropriate culture and chemical environments. When compared to bone marrow-derived MSCs, differentiated MFCs retained a distinct appearance. For example, in chondrogenic pellets, Alcian blue staining filled the entirety of the MSC-laden pellets, while MFC-laden pellets showed pronounced staining only at the periphery. In adipogenic cultures, MSCs were generally larger than MFCs, and their lipid droplets were densely packed in individual cells compared to the wider but more diffuse distribution in MFCs. Under osteogenic conditions, only MFCs from the outer region showed pronounced mineralized matrix deposition, and they did so to a greater extent than the corresponding MSC cultures. The diminished capacity of MFCs from some regions to undergo osteogenesis may point to their distinct phenotype and/or origin. It should be noted that in this study a standard differentiation cocktail was used to induce each lineage. Recent work has shown that adult MSCs derived from different sources (bone marrow vs. adipose tissue) undergo

more or less robust differentiation with different media formulations and growth factors (Estes et al., 2006). Thus, the induction of multiple lineage differentiation in MFCs derived from different regions of the meniscus may be further optimized.

These findings also raised another question: if cells that are sufficiently plastic to participate in repair processes do reside in all zones of the meniscus, then why is endogenous repair of this tissue limited to only the outer zones? This same question may be posed for many of the dense tissues of the musculoskeletal system, such as articular cartilage, ligament, and tendon, all of which have been shown to contain a similar cell population, and yet fail to heal. One explanation for these observations may simply be the density of these tissues. To attain their mature load-bearing capacity, these tissues sacrifice vascular and neural invaginations that are present during development to achieve the requisite mechanical properties for adult function (Clark and Ogden, 1983). Without a route for cellular mobility, this dense tissue may impede the progression of progenitor cells to the point of injury. Furthermore, as mechanical loading is typically present during the regenerative process, mechanical stress at the newly forming union may modulate differentiation toward a fibrous phenotype (Altman et al., 2002), or simply result in micro- and macroscopic fractures within the forming neo-tissue prior to its functional maturation. Finally, it has been demonstrated in tissue culture that MFCs within the meniscus are mobilized by the chemical

signals found in fibrin clots (Webber et al., 1989). In vitro studies have also demonstrated a number of cytokines that differentially enhance the migration of MFCs from different meniscus zones (Bhargava et al., 1999). In other systems, homing cytokines, such as stromal cell-derived factor 1, have been implicated in the recruitment of MSCs to sites of injury (Ceradini et al., 2004; Ji et al., 2004; Ma et al., 2005). Without exposure to gradients of these signals, multipotential cells embedded within the meniscus (particularly the avascular inner region) may not migrate toward the site of damage. Local administration of such signals, either by protein (Imler et al., 2004) or gene delivery (Goto et al., 1999; Hidaka et al., 2002), may improve the migration and localized matrix deposition by endogenous MFCs and thereby accelerate meniscus repair.

While improving migration of MFCs to sites of injury may hasten meniscus repair, the exogenous delivery of multipotential MSCs may be a more direct approach. This is particularly relevant considering the limited differentiation spectrum of MFCs from the horn and inner regions observed in this study. To this end, one recent study reported on the delivery of MSCs to meniscus defects within a fibrin clot. These cells were shown to maintain residence in the defect for up to 8 weeks after implantation (Izuta et al., 2005; Yamasaki et al., 2005). In another study, MSCs delivered in fibrin glue enhanced filling of an avascular defect (Ishimura et al., 1997), although a similar study showed no improvement in tensile properties across the defect site compared to repair with sutures alone (Port et al., 1996). Clearly, MSCs can play a role in the reparative process, although their administration (and carrier materials) requires further optimization.

While the findings of this study are intriguing, several issues bear further consideration. First, the MFC populations used in this study were derived from young healthy bovine donors. In both animals and humans, meniscus vascularity decreases substantially with increasing age (Gardner and O'Rahilly, 1968). In the inner meniscal tissue used in this study, vascular channels were not apparent, although the persistence of cells from these vascular remnants may have contributed to the observation of cells with multipotential characteristics in this region. In addition to aging effects, differences have been noted in the menisci between species. For example, anatomic form (size and shape) varies, and in some species (rat, for example), a central ossicle is observed within the meniscus substance (Messner and Gao, 1998). Therefore, extrapolation of our findings in a young animal model to that of aged human tissues warrants further study. Finally, it has recently been demonstrated via flow cytometry that digestion of whole menisci yields a heterogeneous population of cells, with a main MFC group that is CD44⁺/CD105⁺, and a mixed subgroup containing CD34⁺ cells (Verdonk et al., 2005). It has been postulated that the CD34⁺ group, derived primarily from the vascular outer zone and a thin synovial sublining, may contain an endothelial progenitor population. These authors reported that this subpopulation of cells disappeared within 2 weeks of monolayer expansion and therefore would not be expected to play a significant role in our differentiated populations, although their presence and contribution to the patterns observed in the present study cannot fully be discounted.

Taken together, the results of this study suggest that fibrochondrocytes from all regions of the meniscus possess a multilineage differentiation capability and may therefore

contribute to endogenous repair processes. The finding that MFCs from the outer region are most plastic may underlie the recent experimental observation of improved integration of meniscus pieces from the outer zone transplanted to avascular defects in an organ culture model (Kobayashi et al., 2004). While multilineage differentiation is not per se a requirement for in vivo meniscus repair, these findings demonstrate the presence of a population of cells with a flexible phenotype throughout the tissue. Thus, in the inner region, where multipotential cells are present, a limitation in healing capacity is more likely due to the mechanical demands placed on the injured site or impediments to cellular migration within this dense fibrous structure rather than the local cellular composition. As such, more direct approaches involving the enhancement of MFC migration or the direct delivery of MSCs on scaffolds with the appropriate structural and mechanical properties (Mauck et al., 2005) will be required to improve meniscus repair. If successful, such cell-based tissue engineering therapies would rectify what is otherwise an untreatable and progressively debilitating orthopedic disorder.

ACKNOWLEDGMENT

The authors thank Dr. Paul Manner, George Washington University, for providing human bone marrow specimens.

LITERATURE CITED

- Adams ME, Hukins DWL. 1992. The extracellular matrix of the meniscus. In: Mow VC, Arnoczky SP, Jackson DW, editors. *Knee meniscus: basic and clinical foundations*. New York: Raven Press. p 15–28.
- Ahmed AM. 1992. The load-bearing role of the knee meniscus. In: Mow VC, Arnoczky SP, Jackson DW, editors. *Knee meniscus: basic and clinical foundations*. New York: Raven Press. p 59–73.
- Alsalameh S, Amin R, Gemba T, Lotz M. 2004. Identification of mesenchymal progenitor cells in normal and osteoarthritic human articular cartilage. *Arthritis Rheum* 50:1522–1532.
- Altman GH, Horan RL, Martin I, Farhadi J, Stark PR, Volloch V, Richmond JC, Vunjak-Novakovic G, Kaplan DL. 2002. Cell differentiation by mechanical stress. *FASEB J* 16:270–272.
- Arnoczky SP, Warren RF. 1983. The microvasculature of the meniscus and its response to injury: an experimental study in the dog. *Am J Sports Med* 11:131–141.
- Arnoczky SP, Warren RF, Kaplan N. 1985. Meniscal remodeling following partial meniscectomy: an experimental study in the dog. *Arthroscopy* 1:247–252.
- Arnoczky SP, Adams ME, DeHaven KE, Eyre DR, Mow VC. 1988a. Meniscus. In: Woo SL, Buckwalter JA, editors. *Injury and repair of the musculoskeletal soft tissues*. Park Ridge, IL: American Academy of Orthopedic Surgery. p 487–537.
- Arnoczky SP, Warren RF, Spivak JM. 1988b. Meniscal repair using an exogenous fibrin clot: an experimental study in dogs. *J Bone Joint Surg Am* 70:1209–1217.
- Arnoczky SP. 1992. Gross and vascular anatomy of the meniscus and its role in meniscal healing, regeneration, and remodeling. In: Mow VC, Arnoczky SP, Jackson DW, editors. *Knee meniscus: basic and clinical foundations*. New York: Raven Press. p 1–14.
- Baksh D, Song L, Tuan RS. 2004. Adult mesenchymal stem cells: characterization, differentiation, and application in cell and gene therapy. *J Cell Mol Med* 8:301–316.
- Baratz ME, Fu FH, Mengato R. 1986. Meniscal tears: the effect of meniscectomy and of repair on intraarticular contact areas and stress in the human knee—a preliminary report. *Am J Sports Med* 14:270–275.

- Barbero A, Ploegert S, Heberer M, Martin I. 2003. Plasticity of clonal populations of dedifferentiated adult human articular chondrocytes. *Arthritis Rheum* 48:1315–1325.
- Barry FP. 2003. Biology and clinical applications of mesenchymal stem cells. *Birth Defects Res C Embryo Today* 69:250–256.
- Benjamin M, Ralphs JR. 2004. Biology of fibrocartilage cells. *Int Rev Cytol* 233:1–45.
- Bhargava MM, Attia ET, Murrell GA, Dolan MM, Warren RF, Hannafin JA. 1999. The effect of cytokines on the proliferation and migration of bovine meniscal cells. *Am J Sports Med* 27:636–643.
- Blau HM, Brazelton TR, Weimann JM. 2001. The evolving concept of a stem cell: entity or function? *Cell* 105:829–841.
- Bullough PG, Munuera L, Murphy J, Weinstein AM. 1970. The strength of the menisci of the knee as it relates to their fine structure. *J Bone Joint Surg Br* 52:564–567.
- Buma P, Ramrattan NN, van Tienen TG, Veth RP. 2004. Tissue engineering of the meniscus. *Biomaterials* 25:1523–1532.
- Caplan AI. 1991. Mesenchymal stem cells. *J Orthop Res* 9:641–650.
- Ceradini DJ, Kulkarni AR, Callaghan MJ, Tepper OM, Bastidas N, Kleinman ME, Capla JM, Galiano RD, Levine JP, Gurtner GC. 2004. Progenitor cell trafficking is regulated by hypoxic gradients through HIF-1 induction of SDF-1. *Nat Med* 10:858–864.
- Cheung HS. 1987. Distribution of type I, II, III and V in the pepsin solubilized collagens in bovine menisci. *Connect Tissue Res* 16:343–356.
- Clark CR, Ogden JA. 1983. Development of the menisci of the human knee joint: morphological changes and their potential role in childhood meniscal injury. *J Bone Joint Surg Am* 65:538–547.
- Cravener EK, MacElroy DG. 1941. Injuries of the internal semilunar cartilage. *JAMA* 117:1695–1700.
- De Bari C, Dell'Accio F, Vandenabeele F, Vermeesch JR, Raymackers JM, Luyten FP. 2003. Skeletal muscle repair by adult human mesenchymal stem cells from synovial membrane. *J Cell Biol* 160:909–918.
- Elliott DM, Guilak F, Vail TP, Wang JY, Setton LA. 1999. Tensile properties of articular cartilage are altered by meniscectomy in a canine model of osteoarthritis. *J Orthop Res* 17:503–508.
- Erickson GR, Gimble JM, Franklin DM, Rice HE, Awad H, Guilak F. 2002. Chondrogenic potential of adipose tissue-derived stromal cells in vitro and in vivo. *Biochem Biophys Res Commun* 290:763–769.
- Estes BT, Wu AW, Guilak F. 2006. Potent induction of chondrocytic differentiation of human adipose-derived adult stem cells by bone morphogenetic protein 6. *Arthritis Rheum* 54:1222–1232.
- Eyre DR, Wu JJ. 1983. Collagen of fibrocartilage: a distinctive molecular phenotype in bovine meniscus. *FEBS Lett* 158:265–270.
- Farndale RW, Buttle DJ, Barrett AJ. 1986. Improved quantitation and discrimination of sulphated glycosaminoglycans by use of dimethylmethylene blue. *Biochim Biophys Acta* 883:173–177.
- Fithian DC, Kelly MA, Mow VC. 1990. Material properties and structure-function relationships in the menisci. *Clin Orthop* 252:19–31.
- Fitzgerald JB, Jin M, Dean D, Wood DJ, Zheng MH, Grodzinsky AJ. 2004. Mechanical compression of cartilage explants induces multiple time-dependent gene expression patterns and involves intracellular calcium and cyclic AMP. *J Biol Chem* 279:19502–19511.
- Gardner E, O'Rahilly R. 1968. The early development of the knee joint in staged human embryos. *J Anat* 102:289–299.
- Ghosh P, Taylor TK. 1987. The knee joint meniscus: a fibrocartilage of some distinction. *Clin Orthop Relat Res* 224:52–63.
- Goto H, Shuler FD, Lamsam C, Moller HD, Niyibizi C, Fu FH, Robbins PD, Evans CH. 1999. Transfer of lacZ marker gene to the meniscus. *J Bone Joint Surg Am* 81:918–925.
- Hellio Le Graverand MP, Ou Y, Schield-Yee T, Barclay L, Hart D, Natsume T, Rattner JB. 2001a. The cells of the rabbit meniscus: their arrangement, interrelationship, morphological variations and cytoarchitecture. *J Anat* 198:525–535.
- Hellio Le Graverand MP, Vignon E, Otterness IG, Hart DA. 2001b. Early changes in lapine menisci during osteoarthritis development: I, cellular and matrix alterations. *Osteoarthritis Cartil* 9:56–64.
- Hellio Le Graverand MP, Vignon E, Otterness IG, Hart DA. 2001c. Early changes in lapine menisci during osteoarthritis development: II, molecular alterations. *Osteoarthritis Cartil* 9:65–72.
- Hidaka C, Ibarra C, Hannafin JA, Torzilli PA, Quitoriano M, Jen SS, Warren RF, Crystal RG. 2002. Formation of vascularized meniscal tissue by combining gene therapy with tissue engineering. *Tissue Eng* 8:93–105.
- Hoch DH, Grodzinsky AJ, Koob TJ, Albert ML, Eyre DR. 1983. Early changes in material properties of rabbit articular cartilage after meniscectomy. *J Orthop Res* 1:4–12.
- Imler SM, Doshi AN, Levenston ME. 2004. Combined effects of growth factors and static mechanical compression on meniscus explant biosynthesis. *Osteoarthritis Cartil* 12:736–744.
- Ishimura M, Ohgushi H, Habata T, Tamai S, Fujisawa Y. 1997. Arthroscopic meniscal repair using fibrin glue: I, experimental study. *Arthroscopy* 13:551–557.
- Izuta Y, Ochi M, Adachi N, Deie M, Yamasaki T, Shinomiya R. 2005. Meniscal repair using bone marrow-derived mesenchymal stem cells: experimental study using green fluorescent protein transgenic rats. *Knee* 12:217–223.
- Ji JF, He BP, Dheen ST, Tay SS. 2004. Interactions of chemokines and chemokine receptors mediate the migration of mesenchymal stem cells to the impaired site in the brain after hypoglossal nerve injury. *Stem Cells* 22:415–427.
- Kambic HE, McDevitt CA. 2005. Spatial organization of types I and II collagen in the canine meniscus. *J Orthop Res* 23:142–149.
- King D. 1936a. The function of the semilunar cartilages. *J Bone Joint Surg* 18:1069.
- King D. 1936b. The healing of semilunar cartilages. *J Bone Joint Surg* 18:333.
- Kobayashi K, Fujimoto E, Deie M, Sumen Y, Ikuta Y, Ochi M. 2004. Regional differences in the healing potential of the meniscus-an organ culture model to eliminate the influence of microvasculature and the synovium. *Knee* 11:271–278.
- Krause WR, Pope MH, Johnson RJ, Wilder DG. 1976. Mechanical changes in the knee after meniscectomy. *J Bone Joint Surg Am* 58:599–604.
- Lanzer WL, Komenda G. 1990. Changes in articular cartilage after meniscectomy. *Clin Orthop* 252:41–48.
- Lekic P, McCulloch CA. 1996. Periodontal ligament cell population: the central role of fibroblasts in creating a unique tissue. *Anat Rec* 245:327–341.
- LeRoux MA, Arokoski J, Vail TP, Guilak F, Hyttinen MM, Kiviranta I, Setton LA. 2000. Simultaneous changes in the mechanical properties, quantitative collagen organization, and proteoglycan concentration of articular cartilage following canine meniscectomy. *J Orthop Res* 18:383–392.
- Levy IM, Torzilli PA, Warren RF. 1982. The effect of medial meniscectomy on anterior-posterior motion of the knee. *J Bone Joint Surg Am* 64:883–888.
- Li WJ, Danielson KG, Alexander PG, Tuan RS. 2003. Biological response of chondrocytes cultured in three-dimensional nanofibrous poly(epsilon-caprolactone) scaffolds. *J Biomed Mater Res* 67A:1105–1114.
- Ma J, Ge J, Zhang S, Sun A, Shen J, Chen L, Wang K, Zou Y. 2005. Time course of myocardial stromal cell-derived factor 1 expression and beneficial effects of intravenously administered bone marrow stem cells in rats with experimental myocardial infarction. *Basic Res Cardiol* 100:217–223.
- MacAusland WR. 1931. Derangements of the semilunar cartilages. *Ann Surg* 93:649–682.
- Mauck RL, Wang CC, Oswald ES, Ateshian GA, Hung CT. 2003. The role of cell seeding density and nutrient supply for articular cartilage tissue engineering with deformational loading. *Osteoarthritis Cartil* 11:879–890.
- Mauck RL, Li WJ, Yuan X, Tuan RS. 2005. Meniscus tissue engineering with anisotropic biodegradable nanofibrous poly-caprolactone scaffolds. Washington, DC: Proceedings of the 51st Annual Orthopaedic Research Society Meeting. p 30:1722.
- Mauck RL, Yuan X, Tuan RS. 2006. Chondrogenic differentiation and functional maturation of bovine mesenchymal stem cells in long-term agarose culture. *Osteoarthritis Cartil* 14:179–189.
- McDevitt CA, Webber RJ. 1990. The ultrastructure and biochemistry of meniscal cartilage. *Clin Orthop Relat Res* 252:8–18.
- Messner K, Gao J. 1998. The menisci of the knee joint: anatomical and functional characteristics, and a rationale for clinical treatment. *J Anat* 193(Pt 2):161–178.

- Mow VC, Ratcliffe A, Chern KY, Kelly MA. 1992. Structure and function relationships of the menisci of the knee. In: Mow VC, Arnoczky SP, Jackson DW, editors. *Knee meniscus: basic and clinical foundations*. New York: Raven Press. p 37–57.
- Nishimura K, Solchaga LA, Caplan AI, Yoo JU, Goldberg VM, Johnstone B. 1999. Chondroprogenitor cells of synovial tissue. *Arthritis Rheum* 42:2631–2637.
- O'Connor BL. 1976. The histological structure of dog knee menisci with comments on its possible significance. *Am J Anat* 147:407–417.
- Osycka AM, Noth U, Danielson KG, Tuan RS. 2002. Different osteochondral potential of clonal cell lines derived from adult human trabecular bone. *Ann NY Acad Sci* 961:73–77.
- Petersen W, Tillmann B. 1998. Collagenous fibril texture of the human knee joint menisci. *Anat Embryol (Berl)* 197:317–324.
- Petrosini AV, Sherman OH. 1996. A historical perspective on meniscal repair. *Clin Sports Med* 15:445–453.
- Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, Moorman MA, Simonetti DW, Craig S, Marshak DR. 1999. Multilineage potential of adult human mesenchymal stem cells. *Science* 284:143–147.
- Port J, Jackson DW, Lee TQ, Simon TM. 1996. Meniscal repair supplemented with exogenous fibrin clot and autogenous cultured marrow cells in the goat model. *Am J Sports Med* 24:547–555.
- Prockop DJ. 1997. Marrow stromal cells as stem cells for nonhematopoietic tissues. *Science* 276:71–74.
- Rath E, Richmond JC. 2000. The menisci: basic science and advances in treatment. *Br J Sports Med* 34:252–257.
- Sen A, Lea-Currie YR, Sujkowska D, Franklin DM, Wilkison WO, Halvorsen YD, Gimble JM. 2001. Adipogenic potential of human adipose derived stromal cells from multiple donors is heterogeneous. *J Cell Biochem* 81:312–319.
- Seo BM, Miura M, Sonoyama W, Coppe C, Stanyon R, Shi S. 2005. Recovery of stem cells from cryopreserved periodontal ligament. *J Dent Res* 84:907–912.
- Setton LA, Guilak F, Hsu EW, Vail TP. 1999. Biomechanical factors in tissue engineered meniscal repair. *Clin Orthop* 367 Suppl:S254–S272.
- Shin SJ, Fermor B, Weinberg JB, Pisetsky DS, Guilak F. 2003. Regulation of matrix turnover in meniscal explants: role of mechanical stress, interleukin-1, and nitric oxide. *J Appl Physiol* 95:308–313.
- Song L, Tuan RS. 2004. Transdifferentiation potential of human mesenchymal stem cells derived from bone marrow. *FASEB J* 18:980–982.
- Sweigart MA, Athanasiou KA. 2001. Toward tissue engineering of the knee meniscus. *Tissue Eng* 7:111–129.
- Tallheden T, Dennis JE, Lennon DP, Sjogren-Jansson E, Caplan AI, Lindahl A. 2003. Phenotypic plasticity of human articular chondrocytes. *J Bone Joint Surg* 85A:93–100.
- Tuli R, Tuli S, Nandi S, Wang ML, Alexander PG, Haleem-Smith H, Hozack WJ, Manner PA, Danielson KG, Tuan RS. 2003. Characterization of multipotential mesenchymal progenitor cells derived from human trabecular bone. *Stem Cells* 21:681–693.
- Upton ML, Chen J, Guilak F, Setton LA. 2003. Differential effects of static and dynamic compression on meniscal cell gene expression. *J Orthop Res* 21:963–969.
- Verdonk PC, Forsyth RG, Wang J, Almqvist KF, Verdonk R, Veys EM, Verbruggen G. 2005. Characterisation of human knee meniscus cell phenotype. *Osteoarthritis Cartil* 13:548–560.
- Warejcka DJ, Harvey R, Taylor BJ, Young HE, Lucas PA. 1996. A population of cells isolated from rat heart capable of differentiating into several mesodermal phenotypes. *J Surg Res* 62:233–242.
- Webber RJ, Harris MG, Hough AJ Jr. 1985. Cell culture of rabbit meniscal fibrochondrocytes: proliferative and synthetic response to growth factors and ascorbate. *J Orthop Res* 3:36–42.
- Webber RJ, York JL, Vanderschelden JL, Hough AJ Jr. 1989. An organ culture model for assaying wound repair of the fibrocartilaginous knee joint meniscus. *Am J Sports Med* 17:393–400.
- Webber RJ. 1990. In vitro culture of meniscal tissue. *Clin Orthop Relat Res* 252:114–120.
- Williams JT, Southerland SS, Souza J, Calcutt AF, Cartledge RG. 1999. Cells isolated from adult human skeletal muscle capable of differentiating into multiple mesodermal phenotypes. *Am Surg* 65:22–26.
- Yamasaki T, Deie M, Shinomiya R, Izuta Y, Yasunaga Y, Yanada S, Sharman P, Ochi M. 2005. Meniscal regeneration using tissue engineering with a scaffold derived from a rat meniscus and mesenchymal stromal cells derived from rat bone marrow. *J Biomed Mater Res A* 75:23–30.
- Young HE, Steele TA, Bray RA, Hudson J, Floyd JA, Hawkins K, Thomas K, Austin T, Edwards C, Cuzzourt J, Duenzl M, Lucas PA, Black AC Jr. 2001. Human reserve pluripotent mesenchymal stem cells are present in the connective tissues of skeletal muscle and dermis derived from fetal, adult, and geriatric donors. *Anat Rec* 264:51–62.
- Zuk PA, Zhu M, Mizuno H, Huang J, Futrell JW, Katz AJ, Benhaim P, Lorenz HP, Hedrick MH. 2001. Multilineage cells from human adipose tissue: implications for cell-based therapies. *Tissue Eng* 7:211–228.