

Asymmetric Cyclopropanation

# Highly Diastereo- and Enantioselective Cyclopropanation of 1,2-Disubstituted Alkenes\*\*


Jun Li, Sai-Hu Liao, Hu Xiong, You-Yun Zhou, Xiu-Li Sun,\* Yue Zhang, Xiao-Guang Zhou, and Yong Tang\*

Since Nozaki et al. reported the first enantioselective synthesis of cyclopropanes, a reaction that involved copper-catalyzed transfer of a carbene moiety from diazo compounds to alkenes,<sup>[1]</sup> much effort has been devoted to the area of transition-metal-catalyzed asymmetric cyclopropanation reactions because it is a straightforward method for accessing optically active cyclopropanes.<sup>[2–7]</sup> However, there are only a few examples where 1,2-disubstituted alkenes have been transformed through a transition-metal-catalyzed asymmetric cyclopropanation reaction with high levels of diastereo- and enantioselectivity;<sup>[8]</sup> these reactions usually involve cyclic alkenes<sup>[8c]</sup> and trisubstituted alkenes.<sup>[8a,e]</sup> In 1991, Masamune et al. reported a double-asymmetric-induction approach in which *cis*- $\beta$ -methyl styrene was transformed using a Cu<sup>I</sup>/BOX-catalyzed cyclopropanation reaction involving L-menthol-derived diazoacetate to give product in 92% *ee* and 76% *de*.<sup>[8a]</sup> High enantioselectivity was achieved by Ito and Katsuki when they used chiral bipyridine ligands in the cyclopropanation of *trans*- $\beta$ -methyl styrene, although the diastereoselectivity was low (*trans/cis* 40:60).<sup>[8b]</sup> Recently, Katsuki and co-workers reported the use of an aryliridium/salen catalyst, which led to remarkably high levels of enantio- and diastereoselectivity (favoring the *cis* product) in the cyclopropanation of terminal and cyclic alkenes. However, when *cis*- $\beta$ -methyl styrene was used as a substrate, a relatively low yield of product (29%) was obtained and for *trans*- $\beta$ -methyl styrene only a trace amount of cyclopropanation product was obtained.<sup>[8c]</sup> The unsatisfactory results obtained in the cases of 1,2-disubstituted alkenes can be mainly ascribed to the high sensitivity of metallocarbenes to the steric hindrance and geometry of the alkene.<sup>[8]</sup> Therefore, a cyclopropanation catalyst that is efficient and applicable to the highly stereoselective cyclopropanation of both *cis*- and *trans*-1,2-disubstituted alkenes, especially simple *trans* alkenes, is still in high demand. Herein, we report that the use

of bis(oxazoline) (BOX) ligands that contain C<sub>2</sub>-symmetry-breaking pendant groups in the copper-catalyzed cyclopropanation of both *cis*- and *trans*-1,2-disubstituted alkenes can lead to high levels of diastereo- and enantioselectivity.

We commenced our study by screening copper salts in combination with several BOX ligands in the cyclopropanation reaction of *cis*- $\beta$ -methyl styrene (Table 1). When using


**Table 1:** Screening of ligands for the copper-catalyzed cyclopropanation reaction.<sup>[a]</sup>



Entry	Ligand	Yield [%] <sup>[b]</sup>	<i>Trans/cis</i> <sup>[c]</sup>	<i>ee</i> <sub><i>trans</i></sub> [%] <sup>[d]</sup>
1	<b>L1</b>	25	75/25	7
2	<b>L2</b>	0	—	—
3	<b>L3</b>	45	86/14	7
4	<b>L4a</b>	61	95/5	89
5	<b>L5a</b>	53	97/3	87
6	<b>L6a</b>	50	94/6	80

[a] **1a** (0.5 mmol), CH<sub>3</sub>CO<sub>2</sub>*i*Bu (3.5 mL). [b] Yield of isolated product.

[c] Determined by <sup>1</sup>H NMR analysis. [d] Determined by HPLC using a chiral stationary phase.



[\*] J. Li, Dr. S.-H. Liao, H. Xiong, Dr. Y.-Y. Zhou, Dr. X.-L. Sun, Y. Zhang, X.-G. Zhou, Prof. Dr. Y. Tang

State Key Laboratory of Organometallic Chemistry  
Shanghai Institute of Organic Chemistry  
Chinese Academy of Sciences  
345 Lingling Lu, Shanghai 200032 (P.R. China)  
E-mail: xlsun@sioc.ac.cn  
tangy@mail.sioc.ac.cn


[\*\*] We are grateful for the financial support from the National Natural Sciences Foundation of China (No. 21121062, 20932008, and 21072207), the Major State Basic Research Development Program (Grant No. 2009CB825300) and the Chinese Academy of Sciences.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201203218>.

2,6-dimethylphenyl diazoacetate as the carbene source,<sup>[9]</sup> all ligands tested exhibited the required activity, except Ph-DBFOX (**L2**; Table 1, entry 2); only low levels of enantioselectivity were obtained with **L1** and **L3** (Table 1, entries 1 and 3). Because *i*Pr-BOX (**L4a**) gave the most promising enantioselectivity (89% *ee*; Table 1, entry 4), we prepared and tested ligands **L5a** and **L6a**, which have different substituents on the C4 atom. Unfortunately, the levels of enantioselectivity that were obtained using these ligands were not an improvement on that obtained using **L4a** and were thus impractical (Table 1, entries 5 and 6).



**Table 3:** Asymmetric cyclopropanation of *Z* alkenes.<sup>[a]</sup>



Entry	Alkene	Yield [%] <sup>[b]</sup>	Trans/cis <sup>[c]</sup>	ee <sub>trans</sub> [%] <sup>[d]</sup>
1	<b>1a</b> :	84	96/4	92
2	<b>1c</b> :	78	96/4	93
3	<b>1d</b> :	72	95/5	94
4	<b>1e</b> :	72	95/5	93
5	<b>1f</b> :	60	97/3	89
6	<b>1g</b> :	95	97/3	89
7	<b>1h</b> :	60	97/3	86
8 <sup>[e]</sup>	<b>1i</b> :	66	93/7	86


[a] Alkene (0.5 mmol), CH<sub>3</sub>CO<sub>2</sub>iBu (3.5 mL). [b] Yield of isolated product. [c] Determined by <sup>1</sup>H NMR analysis. [d] Determined by HPLC using a chiral stationary phase. [e] 4 equivalents of diazo acetate were used.

entries 6 and 7). Although a relatively low yield was obtained in the reaction of the more sterically demanding *cis*-β-ethyl styrene, high diastereoselectivity (93:7) and enantioselectivity (86% *ee*) were observed (Table 3, entry 8).

The cyclopropanation reaction of 1,2-*trans*-disubstituted alkenes had broad substrate scope and high levels of stereoselectivity (Table 4). Using **L7b**/CuOTf as the catalyst, various *trans* β-methyl styrene derivatives bearing electron-donating and electron-withdrawing substituents on the phenyl ring, underwent cyclopropanation with perfect levels of diastereoselectivity (> 99:1) and high levels of enantioselectivity (94–97% *ee*; Table 4, entries 1–5). 1-Naphthyl and 1-cinnamyl alkene were also suitable substrates and were transformed into the corresponding cyclopropanes with high levels of *trans* selectivity and enantioselectivity (Table 4, entries 6–7). Notably, under the optimized reaction conditions, more hindered *trans* alkenes, such as cinnamyl-alcohol derivative **1p** and β-ethyl styrene **1q**, can also be converted into the desired cyclopropanes with high levels of stereoselectivity (Table 4, entries 8 and 9), and a Sommelet–Hauser rearrangement product was not observed in the reaction of **1p**. Moreover, the reaction of trisubstituted alkene **1r** also proceeded well, affording fused bicyclic product **2r** in 82% yield with greater than 99:1 *trans/cis* and 96% *ee* (Table 4, entry 10).


Considering the high efficiency of the **L7b**/CuOTf catalyzed cyclopropanation of *trans* alkenes, a scaled-up reaction (50 mmol) was performed. In the event, high yields and high levels of enantioselectivity were obtained when 0.5 mol% of catalyst was used (Scheme 1). The cyclopropanation of *trans*-β-methyl styrene also proceeded well even with only 0.05 mol% catalyst loading, giving the desired propane **2b**

**Table 4:** Asymmetric cyclopropanation of *E* alkenes.<sup>[a]</sup>



Entry	Alkene	Yield [%] <sup>[b]</sup>	Trans/cis <sup>[c]</sup>	ee <sub>trans</sub> [%] <sup>[d]</sup>
1	<b>1b</b> :	89	> 99/1	96
2	<b>1j</b> :	99	> 99/1	96
3	<b>1k</b> :	96	> 99/1	94
4	<b>1l</b> :	96	> 99/1	97
5 <sup>[e]</sup>	<b>1m</b> :	73	> 99/1	96
6	<b>1n</b> :	60	> 99/1	96
7	<b>1o</b> :	97	93/7	96
8	<b>1p</b> :	64	> 99/1	98
9	<b>1q</b> :	84	> 99/1	97
10	<b>1r</b> :	82	> 99/1	96

[a] Alkene (0.5 mmol), CH<sub>3</sub>CO<sub>2</sub>iBu (3.5 mL). [b] Yield of isolated product. [c] Determined by <sup>1</sup>H NMR analysis. [d] Determined by HPLC using a chiral stationary phase. [e] The absolute configuration was determined to be (1*R*,2*R*,3*R*) by X-ray crystallography.<sup>[14]</sup>



**Scheme 1.** Reactions with reduced catalyst loading.

in 44% yield with 98% *ee*. To our knowledge, 0.05 mol% is the lowest catalyst loading used in a copper-catalyzed cyclopropanation reaction.

In summary, an efficient BOX/Cu<sup>I</sup>-catalyzed cyclopropanation reaction has been designed and developed. *Cis*- and *trans*-1,2-substituted alkenes can be converted into the corresponding trisubstituted cyclopropanes with high levels of diastereo- and enantioselectivity (> 99:1 *trans/cis* and up to 98% *ee*). The effect of the pendant group of BOX ligands was investigated, thus leading to the discovery of the BOX ligand **L4b** and **L7b**, the copper complexes of which were the best catalysts. This reaction features high catalytic efficiency and excellent stereoselectivity, especially for *trans* alkenes, the generality and high diastereoselectivity of which are unprecedented. Further investigations of this catalytic reaction are underway.

## Experimental Section

A typical procedure using the reaction that gives the product **2a** as an example: a mixture of CuOTf·0.5PhCH<sub>3</sub> (0.025 mmol), **L4b** (0.0275 mmol), and activated 4 Å MS (300 mg) in CH<sub>3</sub>CO<sub>2</sub>iBu (1 mL) was stirred at 30 °C for 1 h under N<sub>2</sub> atmosphere. **1a** (0.5 mmol) in CH<sub>3</sub>CO<sub>2</sub>iBu (0.5 mL) were added, followed by the diazo acetate (1.0 mmol) in CH<sub>3</sub>CO<sub>2</sub>iBu (0.5 mL) which was added

dropwise using a syringe pump over 8 h. After the reaction was complete, the mixture was filtered through a thin layer of silica gel, eluting with  $\text{CH}_2\text{Cl}_2$ , and the filtrate was concentrated. The residue was purified by column chromatography over silica gel using  $\text{CH}_2\text{Cl}_2$ /petroleum ether 6:1 as eluent to afford **2a** (118 mg, 84% yield).

Received: April 26, 2012

Revised: July 2, 2012

Published online: July 24, 2012

**Keywords:** alkenes · copper · cyclopropanation · diazo compounds · homogeneous catalysis

- [1] a) H. Nozaki, H. Takaya, R. Noyori, *Tetrahedron Lett.* **1965**, 6, 2563–2567; b) H. Nozaki, H. Takaya, S. Moriuti, R. Noyori, *Tetrahedron* **1968**, 24, 3655–3669.
- [2] For reviews, see: a) A. Padwa, S. F. Hornbuckle, *Chem. Rev.* **1991**, 91, 263–309; b) A. Pfaltz, *Acc. Chem. Res.* **1993**, 26, 339–345; c) T. Ye, M. A. McKervey, *Chem. Rev.* **1994**, 94, 1091–1160; d) M. P. Doyle, D. C. Forbes, *Chem. Rev.* **1998**, 98, 911–936; e) H. Lebel, J. F. Marcoux, C. Molinaro, A. B. Charette, *Chem. Rev.* **2003**, 103, 977–1050; f) H. Pellissier, *Tetrahedron* **2008**, 64, 7041–7095; g) Z. Zhang, J. Wang, *Tetrahedron* **2008**, 64, 6577–6605.
- [3] For selected recent examples of Rh-catalyzed reactions, see: a) D. Marcoux, A. B. Charette, *Angew. Chem.* **2008**, 120, 10309–10312; *Angew. Chem. Int. Ed.* **2008**, 47, 10155–10158; b) J. R. Denton, K. Cheng, H. M. L. Davies, *Chem. Commun.* **2008**, 1238–1240; c) A. DeAngelis, O. Dmitrenko, G. P. A. Yap, J. M. Fox, *J. Am. Chem. Soc.* **2009**, 131, 7230–7231; d) V. N. G. Lindsay, W. Lin, A. B. Charette, *J. Am. Chem. Soc.* **2009**, 131, 16383–16385; e) S. Chuprakov, S. W. Kwok, L. Zhang, L. Lercher, V. V. Fokin, *J. Am. Chem. Soc.* **2009**, 131, 18034–18035; f) T. Nishimura, Y. Maeda, T. Hayashi, *Angew. Chem.* **2010**, 122, 7482–7485; *Angew. Chem. Int. Ed.* **2010**, 49, 7324–7327; g) A. Ghanem, M. G. Gardiner, R. M. Williamson, P. Müller, *Chem. Eur. J.* **2010**, 16, 3291–3295; h) P. Pelphrey, J. Hansen, H. M. L. Davies, *Chem. Sci.* **2010**, 1, 254–257; i) C. Qin, V. Boyarskikh, J. H. Hansen, K. I. Hardcastle, D. G. Musaev, H. M. L. Davies, *J. Am. Chem. Soc.* **2011**, 133, 19198–19204; j) V. N. G. Lindsay, C. Nicolas, A. B. Charette, *J. Am. Chem. Soc.* **2011**, 133, 8972–8981; k) M. Zibinsky, V. V. Fokin, *Org. Lett.* **2011**, 13, 4870–4872.
- [4] For selected recent examples of Ru-catalyzed reactions, see: a) I. Nicolas, P. L. Maux, G. Simonneaux, *Tetrahedron Lett.* **2008**, 49, 2111–2113; b) Z.-J. Xu, R. Fang, C. Zhao, J.-S. Huang, G.-Y. Li, N. Zhu, C.-M. Che, *J. Am. Chem. Soc.* **2009**, 131, 4405–4417; c) C. S. Gill, K. Venkatasubbaiah, C. W. Jones, *Adv. Synth. Catal.* **2009**, 351, 1344–1354; d) J. Ito, S. Ujiiie, H. Nishiyama, *Chem. Eur. J.* **2010**, 16, 4986–4990; e) A.-M. Abu-Elfotouh, K. Phomkeona, K. Shibatomi, S. Iwasa, *Angew. Chem.* **2010**, 122, 8617–8621; *Angew. Chem. Int. Ed.* **2010**, 49, 8439–8443; f) A. Mezzetti, *Dalton Trans.* **2010**, 39, 7851–7869; g) J. M. Falkowski, C. Wang, S. Liu, W. Lin, *Angew. Chem.* **2011**, 123, 8833–8837; *Angew. Chem. Int. Ed.* **2011**, 50, 8674–8678; h) C. del Pozo, A. Corma, M. Iglesias, F. Sánchez, *Green Chem.* **2011**, 13, 2471–2481.
- [5] For selected recent examples of Cu-catalyzed reactions, see: a) A. Mandoli, M. Lessi, D. Pini, C. Evangelisti, P. Salvadori, *Adv. Synth. Catal.* **2008**, 350, 375–379; b) A. Mandoli, R. Garzelli, S. Orlandi, D. Pini, M. Lessi, P. Salvadori, *Catal. Today* **2009**, 140, 51–57; c) D. R. Boyd, N. D. Sharma, L. Sbircea, D. Murphy, T. Belhocine, J. F. Malone, S. L. James, C. C. R. Allen, J. T. G. Hamilton, *Chem. Commun.* **2008**, 5535–5537; d) T. Minuth, M. Irmak, A. Groschner, T. Lehnert, M. M. K. Boysen, *Eur. J. Org. Chem.* **2009**, 997–1008; e) E. P. Carreiro, J. P. P. Ramalho, A. J. Burke, *Tetrahedron* **2011**, 67, 4640–4648.
- [6] For selected recent examples of Co-catalyzed reactions, see: a) S. Fantauzzi, E. Gallo, E. Rose, N. Raoul, A. Caselli, S. Issa, F. Ragaini, S. Cenini, *Organometallics* **2008**, 27, 6143–6151; b) S. Zhu, J. A. Perman, X. P. Zhang, *Angew. Chem.* **2008**, 120, 8588–8591; *Angew. Chem. Int. Ed.* **2008**, 47, 8460–8463; c) S. Zhu, J. V. Ruppel, H. Lu, L. Wojtas, X. P. Zhang, *J. Am. Chem. Soc.* **2008**, 130, 5042–5043; d) J. V. Ruppel, T. J. Gauthier, N. L. Snyder, J. A. Perman, X. P. Zhang, *Org. Lett.* **2009**, 11, 2273–2276; e) S. Zhu, X. Xu, J. A. Perman, X. P. Zhang, *J. Am. Chem. Soc.* **2010**, 132, 12796–12799; f) X. Xu, H. Lu, J. V. Ruppel, X. Cui, S. L. de Mesa, L. Wojtas, X. P. Zhang, *J. Am. Chem. Soc.* **2011**, 133, 15292–15295.
- [7] For recent examples, see: a) M. Ichinose, H. Suematsu, T. Katsuki, *Angew. Chem.* **2009**, 121, 3167–3169; *Angew. Chem. Int. Ed.* **2009**, 48, 3121–3123; b) I. Nicolas, T. Roisnel, P. L. Maux, G. Simonneaux, *Tetrahedron Lett.* **2009**, 50, 5149–5151; c) C.-T. Yeung, K.-C. Sham, W.-S. Lee, W.-T. Wong, W.-Y. Wong, H.-L. Kwong, *Inorg. Chim. Acta* **2009**, 362, 3267–3273; d) T. Osswald, I. S. Mikhel, H. Rüegger, P. Butti, A. Mezzetti, *Inorg. Chim. Acta* **2010**, 363, 474–480.
- [8] a) R. E. Lowenthal, S. Masamune, *Tetrahedron Lett.* **1991**, 32, 7373–7376; b) K. Ito, T. Katsuki, *Synlett* **1993**, 638–640; c) H. Suematsu, S. Kanchiku, T. Uchida, T. Katsuki, *J. Am. Chem. Soc.* **2008**, 130, 10327–10337; d) H. M. L. Davies, M. G. Coleman, D. L. Ventura, *Org. Lett.* **2007**, 9, 4971–4974; e) N. Østergaard, J. F. Jensen, D. Tanner, *Tetrahedron* **2001**, 57, 6083–6088.
- [9] D. A. Evans, K. A. Woerpel, M. M. Hinman, M. M. Faul, *J. Am. Chem. Soc.* **1991**, 113, 726–728. See also the Supporting Information.
- [10] H. Fritschi, U. Leutenegger, A. Pfaltz, *Helv. Chim. Acta* **1988**, 71, 1553–1565.
- [11] J. M. Fraile, J. I. García, V. Martínez-Merino, J. A. Mayoral, L. Salvatella, *J. Am. Chem. Soc.* **2001**, 123, 7616–7625.
- [12] a) M. Seitz, C. Capacchione, S. Bellemin-Lapponnaz, H. Wade-pohl, B. D. Ward, L. H. Gade, *Dalton Trans.* **2006**, 193–202; b) C. Foltz, M. Enders, S. Bellemin-Lapponnaz, H. Wade-pohl, L. H. Gade, *Chem. Eur. J.* **2007**, 13, 5994.
- [13] CCDC 878058 (**2m**), CCDC 883866 (**16d**/CuBr<sub>2</sub>) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).
- [14] a) J. Zhou, Y. Tang, *J. Am. Chem. Soc.* **2002**, 124, 9030–9131; b) J. Zhou, Y. Tang, *Chem. Soc. Rev.* **2005**, 34, 664–676; c) Z.-H. Xu, S.-N. Zhu, X.-L. Sun, Y. Tang, L.-X. Dai, *Chem. Commun.* **2007**, 1960–1962; d) Y.-B. Kang, X.-L. Sun, Y. Tang, *Angew. Chem.* **2007**, 119, 3992; *Angew. Chem. Int. Ed.* **2007**, 46, 3918; e) P. Cao, C. Deng, Y.-Y. Zhou, X.-L. Sun, J.-C. Zheng, Z.-W. Xie, Y. Tang, *Angew. Chem.* **2010**, 122, 4565; *Angew. Chem. Int. Ed.* **2010**, 49, 4463.