

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/11184437>

# The role of Alu repeat clusters as mediators of recurrent chromosomal aberrations in tumors

ARTICLE *in* GENES CHROMOSOMES AND CANCER · OCTOBER 2002

Impact Factor: 4.04 · DOI: 10.1002/gcc.10111 · Source: PubMed

---

CITATIONS

198

---

READS

44

4 AUTHORS, INCLUDING:



[Elena Kolomietz](#)

University of Toronto

16 PUBLICATIONS 473 CITATIONS

SEE PROFILE



[Ajay Pandita](#)

University of Toronto

49 PUBLICATIONS 2,113 CITATIONS

SEE PROFILE



[Jeremy Squire](#)

University of São Paulo

395 PUBLICATIONS 23,706 CITATIONS

SEE PROFILE

# The Role of *Alu* Repeat Clusters as Mediators of Recurrent Chromosomal Aberrations in Tumors

Elena Kolomietz,<sup>1</sup> M. Stephen Meyn,<sup>2</sup> Ajay Pandita,<sup>3</sup> and Jeremy A. Squire<sup>1\*</sup>

<sup>1</sup>Department of Laboratory Medicine and Pathobiology and Department of Medical Biophysics, University of Toronto, Princess Margaret Hospital and Ontario Cancer Institute, Toronto, Canada

<sup>2</sup>Genetics and Genomic Biology Program, Department of Molecular and Medical Genetics and Pediatrics, University of Toronto, The Hospital for Sick Children, Toronto, Canada

<sup>3</sup>The Arthur and Sonia Labatt Brain Tumor Research Centre, The Hospital for Sick Children, Ontario, Canada

There is increasing evidence for the involvement of repetitive DNA sequences as facilitators of some of the recurrent chromosomal rearrangements observed in human tumors. The high densities of repetitive DNA, such as *Alu* elements, at some chromosomal translocation breakpoint regions has led to the suggestion that these sequences could provide hot spots for homologous recombination, and could mediate the translocation process and elevate the likelihood of other types of chromosomal rearrangements taking place. The *Alu* core sequence itself has been suggested to promote DNA strand exchange and genomic rearrangement, and it has striking sequence similarity to  $\chi$  (which has been shown to stimulate recBCD-mediated recombination in *Escherichia coli*). *Alu* repeats have been shown to be involved in the generation of many constitutional gene mutations in meiotic cells, attributed to unequal homologous recombination and consequent deletions and/or duplication events. It has recently been demonstrated that similar deletion events can take place in neoplasia because several types of leukemia-associated chromosomal rearrangements frequently have submicroscopic deletions immediately adjacent to the translocation breakpoint regions. Significantly, these types of deletions appear to be more likely to take place when the regions subject to rearrangement contain a high density of *Alu* repeats. With the completion of the Human Genome Project, it will soon be possible to create more comprehensive maps of the distribution and densities of repetitive sequences, such as *Alu*, throughout the genome. Such maps will offer unique insights into the relative distribution of cancer translocation breakpoints and the localization of clusters of repetitive DNA. © 2002 Wiley-Liss, Inc.

## INTRODUCTION

Cytogenetic analysis of tumor cells has revealed that recurring chromosomal abnormalities such as translocations, deletions, and inversions are present in many tumors. In leukemias, lymphomas, and sarcomas, these specific chromosomal aberrations are frequently associated with specific morphologic subtypes. There have been a limited number of attempts to understand the underlying molecular mechanisms leading to the formation of the nonrandom genomic rearrangements that cause recurrent chromosomal aberrations in neoplasms. For hematologic neoplasms, several molecular mechanisms have been proposed to explain these chromosomal rearrangements, including illegitimate V(D)J recombination, homologous recombination mediated by *Alu* elements, translin activity, cleavage at sites of Z-DNA structure, topoisomerase II subunit exchange, repair of DNA breaks with nonhomologous chromosomes by nonhomologous end joining (NHEJ), and fragile sites. These mechanisms are not mutually exclusive, and it seems probable that the generation of a chromosomal rearrangement is the end result of a sequence of complex molecular interactions (Aplan, 1999). In this

review, we discuss the evidence for the involvement of *Alu* elements and repetitive DNA in general as facilitators of chromosomal rearrangement.

## SEQUENCE-DEPENDENT MECHANISMS OF REARRANGEMENT

Most of the proposed mechanisms that mediate nonrandom chromosomal rearrangements are sequence-dependent, and it has been suggested that various recombination-promoting sequences play a role. In lymphoid neoplasms, this view is supported by the presence of recombination signal sequence–heptamer/nonamer sequences characteristic of V(D)J recombination at the chromosome breakpoints (Finger et al., 1986; Tycko and Sklar, 1990).

A. Pandita is currently at Department of Experimental Pathology, Mayo Clinic, Rochester, MN.

\*Correspondence to: Dr. Jeremy A. Squire, Ontario Cancer Institute (University Health Network), Division of Cellular and Molecular Biology, 610 University Avenue, Room 9-721, Toronto, Ontario, Canada M5G 2M9. E-mail: jeremy.squire@utoronto.ca

Received 8 November 2001; Accepted 28 February 2002

DOI 10.1002/gcc.10111

The rearrangements observed in lymphoid neoplasms represent a specific dysregulation of the V(D)J recombinase system. However, a more general mechanism must be responsible for the translocations seen in other classes of neoplasms. It has been proposed that high densities of repetitive DNA such as *Alu* elements could provide hot spots for homologous recombination and mediate chromosomal translocation. More specific sequence elements have also been identified at translocation breakpoint regions. A number of chromosomal breakpoints have been mapped to the translin consensus sequence-binding site of the translin protein that is involved in DNA unwinding, which makes these regions more susceptible to nuclease cleavage (Aoki et al., 1995).

Other investigators have noticed some unusual DNA characteristics in the general regions of rearrangement. It has been reported that alternating purine and pyrimidine residues surround chromosomal breakpoints in a number of chromosomal translocations (Boehm et al., 1989; Adachi and Tsujimoto, 1990). These repeated tracts lead to the formation of a left-handed helical structure known as Z-DNA, altering chromatin organization and making the region involved more susceptible to DNA recombination events. Similarly, the presence of site-specific sequences that are uniquely susceptible to the DNA double-strand breaks induced by the treatment with topoisomerase II are thought to lead to translocations (Aplan et al., 1996; Strissel et al., 1998). More general properties of eukaryotic chromatin may also make some regions more susceptible to rearrangement. It has frequently been suggested that fragile sites predispose to intrachromosomal recombination (Glover and Stein, 1988) because cancer breakpoints are often associated with fragile sites (Le Beau and Rowley, 1984; Yunis and Soreng, 1984; Wenger, 1992). Interestingly, it has also been reported that the experimental induction of chromosomal fragile sites can lead to asynchronous replication timing and gene amplification (Debatisse et al., 1998). DNA, in general, is likely to be more susceptible to rearrangement when it is undergoing replication. It follows that the presence of repetitive sequences may, by virtue of high homology, be an additional destabilizing factor that could increase the likelihood of a rearrangement taking place within the developing replication fork (Flores-Rozas and Kolodner, 2000; Michel, 2000).

#### EFFECTS OF DNA DAMAGE

Various published studies have shown that DNA damage induces homologous recombination. DNA-damaging agents constantly challenge the integrity

of chromosomes. To cope with DNA breakage, several different repair mechanisms have evolved, including NHEJ, homologous recombination (HR), and single-strand annealing. Chromosomal DNA breaks may be either single- or double-stranded. These breaks are thought to be potent inducers of recombination and can increase the frequency of HR (Rouet et al., 1994) and rearrangement between loci on nonhomologous chromosomes (Richardson et al., 1998) by 3 orders of magnitude or more. Double-stranded breaks (DSBs) can be generated by exposing cells to ionizing radiation or other types of DNA-damaging agents or from endogenous sources (e.g., nicks caused by free radicals arising from intermediary metabolism). In addition, some cellular processes such as DNA replication can also generate DSBs. It is also known that certain chromosomal regions, by the nature of DNA structure, are prone to spontaneous or induced breakage (Le Beau and Rowley, 1984; Yunis et al., 1987; Koduru and Chaganti, 1988). Certain sequences such as fragile sites are prone to DSBs, increasing the potential for recombination. It seems likely that genomic rearrangements could arise from misrepair of DSBs. Our basic knowledge of these processes in eukaryotes is derived largely by extension from studies in more amenable prokaryotic DNA-repair systems.

#### PROKARYOTIC SYSTEMS OF REPAIR INDICATE THE IMPORTANCE OF HOMOLOGOUS RECOMBINATION

In *Escherichia coli*, at least 25 different proteins are involved in HR. These include RecA, RecBCD, RecF, -G, -J, -N, and so forth; DNA polymerases; DNA topoisomerases; and DNA ligase. Many of these proteins have functional homologs in other bacteria, eukarya, and some phages. The RecA and RecBCD proteins interact with  $\chi$  ("chi") motifs, crossover recombination hot spots that are short, orientation-dependent, strand-specific sequences implicated in Rec-mediated DNA recombination. The RecBCD enzyme is essential for 99% of recombination events occurring at DSBs in wild-type *Escherichia coli*, and for the repair of DNA damage and DSBs. RecBCD enzyme possesses a number of enzymatic activities, including single-stranded (ss) DNA exonuclease, ssDNA endonuclease, double-stranded (ds) DNA exonuclease, and DNA helicase activity. The RecBCD endonuclease unwinds dsDNA and simultaneously cleaves the ssDNA produced until it associates with a  $\chi$  element (Kowalczykowski, 2000).

Upon recognition of the  $\chi$  sequence, the 3'-5' nuclease activity is attenuated, whereas its helicase activity remains unaltered (Dixon and Kowalczykowski, 1991). The  $\chi$  sequence itself causes RecBCD to switch its functional interactions with DNA from exonuclease to recombinase activity. This change in functional state allows  $\chi$  to act as a unique regulatory element by controlling the degradative function of RecBCD enzyme and, at the same time, enhancing its recombination function (Dixon and Kowalczykowski, 1993). The recombination-proficient RecBC enzyme (without a RecD subunit) has the ability to load RecA protein onto the  $\chi$ -containing ssDNA sequences, ensuring incorporation of this ssDNA into a recombinationally proficient nucleoprotein complex (Kowalczykowski, 2000). This step is followed by the ssDNA searching, invading, and pairing with homologous DNA sequences and thus achieving DSB repair. The RecA protein is required for all HR in *Escherichia coli* and preferentially binds to  $\chi$  sequences (Tracy and Kowalczykowski, 1996). The pairing of DNA by RecA protein is homology-dependent, suggesting that the basis of the elevated recombinational activity of  $\chi$  and the increased genomic rearrangements between homologous DNA sequences may be greatly enhanced by local DNA-pairing activity. The specific mechanism by which RecA protein finds DNA sequences with homology is presently unclear, but it is thought to involve association primarily by passive diffusion (von Hippel and Berg, 1989).

#### HOMOLOGOUS RECOMBINATION IN EUKARYOTES

There is now growing evidence that HR has a similar function associated with repair in eukaryotes. Many of the proteins involved in the repair of DSBs by HR are encoded by the *RAD* genes, which were first identified in yeast (Flores-Rozas and Kolodner, 2000). It has been shown that the two critical steps in HR, the search for homologous DNA and the DNA strand-exchange process, comprise a remarkably phylogenetically conserved mechanism. Using the sequences of *E. coli* RecA and *S. cerevisiae* Rad51, *RAD51* homologs from higher eukaryotes were isolated (Shinohara et al., 1993), suggesting that the main process of HR is carried out by recA-like proteins in all living cells. RecA, Rad51p, and human RAD51 polymerize on DNA to form a nucleoprotein filament that searches for homologous DNA (Swagemakers et al., 1998). Further characterization of *RAD51* homologs, however, demonstrated that eukaryotic RAD51 has several functional differences from pro-

karyotic recA (Shinohara and Ogawa, 1999). First, eukaryotic cells contain multiple recA-like homologs. Second, Rad51 protein interacts with various proteins in the recombination reaction. At least seven RecA homologs have been identified in man, and, based on the recent genetic and biochemical analysis of the genes and proteins, these homologs can be divided into three subgroups: Rad51, Dmc1, and Rad55-57 (Shinohara and Ogawa, 1999). Whereas the Rad51 subfamily plays a pivotal role in both mitotic and meiotic recombination (Shinohara et al., 1992), the Dmc1 subfamily is a meiosis-specific homolog of Rad51/RecA (Bishop et al., 1992; Habu et al., 1996). The third group of proteins has an accessory role in the above two families in recombination and DNA repair.

For the enhancement of the Rad51 activities, other proteins, such as the replication protein-A RPA, Rad52, Rad54, Mre11, Rad50; the Nibrin/p95/NBS complex; and Rad55-57 are necessary for optimal formation of Rad51 filaments. It has been suggested that RPA stimulates Rad51-mediated DNA pairing and strand exchange by removal of secondary structures of the ssDNA for the formation of the continuous Rad51 filament (Shinohara and Ogawa, 1999). This RPA function is regulated by Rad52 (Sung, 1997; Benson et al., 1998; New et al., 1998), and it has been shown (Gasior et al., 1998) that Rad51 binding to DNA requires the preceding binding of Rad52 to DNA. Interaction of Rad51 with Rad54, a dsDNA-dependent ATPase of the Swi2/Snf2 superfamily (Petukhova et al., 1998; Swagemakers et al., 1998), is necessary for efficient Rad51-mediated pairing between homologous DNA molecules in vitro because Rad54 facilitates unwinding of a DNA duplex before its interaction with the ssDNA within the Rad51-DNA nucleoprotein filament. Rad51, similar to RecA, requires ssDNA as a substrate on which to form the nucleoprotein filament to carry out its homology search and strand-exchange activities. Exonucleolytic processing of the broken dsDNA is performed by the heterotrimeric complex composed of Mre11, Rad50, and Nibrin/p95 [so called because it is mutated in the human chromosomal instability disorder Nijmegen breakage syndrome (NBS)]/p95 (Dolganov et al., 1996; Trujillo et al., 1998; Paull and Gellert, 1999).

One of the most distinctive features of a eukaryotic Rad51 protein is that the protein interacts with multiple proteins such as *BRCA1* and *BRCA2* gene products (Scully et al., 1997; Wong et al., 1997), ABL (Yuan et al., 1998; Chen et al., 1999), and TP53 (Sturzbecher et al., 1996; Buchhop et al.,

1997) proteins, suggesting that they work in the same DNA-repair pathway for keeping genome integrity.

Rad51 plays a central role in HR and DNA repair. Eukaryotes have evolved a variety of proteins that regulate the Rad51 functions positively and negatively (Shinohara and Ogawa, 1999). The precise functions of these proteins and their role in orchestrating the repair of DSBs are poorly understood and are currently the subject of extensive analyses. For reasons outlined below, we suggest that repetitive sequences may facilitate the process of HR in mammalian DNA, and that this may be instrumental in the process of chromosomal rearrangement.

#### **INTERACTION OF HR AND NHEJ PATHWAYS OF DNA DOUBLE-STRAND REPAIR**

To maintain the integrity of the genome and to prevent the formation of chromosomal rearrangements, fidelity of the repair of DSBs is crucial for each organism (Pastink et al., 2001). In eukaryotes, the repair of DSBs is carried out by two main pathways: the NHEJ and HR mechanisms, with single-strand annealing playing a relatively minor role. The fundamental difference between these pathways is their dependence on DNA homology and the concomitant accuracy of repair (Essers et al., 2000). In general, HR ensures accurate repair by using the undamaged sister chromatid as a template. However, homologous chromosomes or sequence repeats on nonhomologous chromosomes (such as repetitive *Alu* elements) can also serve as homologous repair templates, thereby increasing the risk of loss of heterozygosity by gene conversion or generating chromosomal rearrangements, respectively. Generally, nonmutagenic outcomes of homologous repair in mammalian cells can be explained by the preferred use of sister chromatids as repair templates (Kadyk and Hartwell, 1992; Johnson and Jasin, 2000), with the rate of recombination in mammals between sister chromatids (intrachromosomal recombination) in somatic cells being almost 5,000-fold higher than that which occurs between homologous chromosomes (interchromosomal recombination) (Shulman et al., 1995; Pastink et al., 2001).

In contrast to HR, NHEJ uses no or extremely limited sequence homology to rejoin ends in a manner that is intrinsically error-prone. At present, two multicomponent complexes of NHEJ pathways have been identified. The first is a DNA-dependent protein kinase complex (DNA-PK), containing three components: Ku 70, Ku 80, and

the catalytic subunit of DNA-PK (DNA-PKcs). The second complex is composed of DNA Ligase IV and XRCC4 (for reviews, see Critchlow and Jackson, 1998; Featherstone and Jackson, 1999; Smith and Jackson, 1999). The Ku complex is composed of a heterodimer of Ku 70 and Ku 80, which binds to DNA ends and recruits DNA-PKcs to the DSB, activating its kinase function. It also facilitates ligation by Ligase IV/XRCC4 complex by aligning the ends (Ramsden and Gellert, 1998).

Both HR and NHEJ are important in preserving genomic integrity and preventing the formation of large chromosomal rearrangements and the concomitant risk of neoplasia (Hoeijmakers, 2001). It has been suggested that HR and NHEJ pathways of DNA DSB repair have overlapping roles in the maintenance of chromosomal integrity and each predominates in the different phases of the cell cycle, such that the HR pathway functions during late S–G<sub>2</sub> phase, whereas the NHEJ pathway is preferentially used during G<sub>1</sub>–early S phase (Sonoda et al., 1998; Takata et al., 1998). Indeed, it would be selectively advantageous for cells to restrict HR to the G<sub>2</sub> phase of the cell cycle when the genome has duplicated and the sister chromatid can be used as a template (Pastink et al., 2001). On the other hand, HR in G<sub>1</sub> may be more mutational than NHEJ (Ferguson and Alt, 2001) because of gene conversion events. Convincing evidence has recently been presented that NHEJ and HR can interact in the repair of a single DSB and presumably reduce the risk of chromosomal rearrangements (Richardson and Jasin, 2000).

#### **DENSITY, DISTRIBUTION, AND STRUCTURE OF *Alu* REPEATS**

More than a third of the human genome consists of repetitive sequences that are ubiquitously interspersed with single-copy sequences throughout the genome. A survey of human interspersed repeats has been updated recently (Smit, 1999). Almost all of these interspersed repeats have arisen by retroposition of an RNA intermediate followed by insertion of the resulting cDNA into the genome. *Alu* elements are the most abundant class of interspersed repeat sequences. It has been estimated that there are more than one million copies of the *Alu* core element per haploid human genome (Smit, 1996). *Alu* elements comprise 5 to 10% of the human genome, occurring on the average once every 4 kb, although in practice certain regions are much more densely populated with these repeats (Smit, 1999). Originally detected by hybridization (Korenberg and Rykowski, 1988), *Alu* repeats have






Figure 1. Positions of the cancer breakpoints of recurrent chromosome aberrations (summarized by the Cancer Genome Anatomy Project: <http://www.ncbi.nlm.nih.gov/CCAP/mitelsum.cgi>) map to areas of accumulation of *Alu* repeats within R-bands.

been shown to hybridize preferentially to reverse bands (R-bands) of metaphase chromosomes (Holmquist, 1992; Craig and Bickmore, 1993). Differential distribution of *Alu* repeats in AT- and GC-rich DNA was later confirmed by computer-based analysis (Smit, 1996, 1999), demonstrating accumulation of *Alu* repeats within R-bands (see Fig. 1).

Human *Alu* repeats can be divided into subfamilies whose relative age can be estimated by analyzing sequence divergence away from the core functional sequence associated with transposition (Smit, 1996). The GC content of "young" *Alu* elements is ninefold greater than the average DNA content, and this class of recently transposed *Alu* elements also offers the potential for genomic methylation (Schmid, 1998). Individual *Alu* family members are highly homologous to each other, and 70–98% homologous to the *Alu* consensus sequence. The 282-bp consensus *Alu* structure has a precisely defined 5' end, shared by most of its members, and is a divergent tandem dimer in

which two monomers are separated by an A-rich region (Schmid, 1996). The two monomers are, in turn, homologous to 7SL RNA, an integral component of the signal recognition particle that governs the attachment of ribosomes to membranes of the endoplasmic reticulum. *Alu* elements contain internal A-box and B-box promoter elements for RNA polymerase III and are transcribed by Pol III *in vitro*.

Sequence analysis of the genomic DNA directly adjacent to translocation breakpoints has shown that the conserved 26-base pair (bp) *Alu* core sequence was present at or close (within 20–50 bp downstream or upstream) to the sites of recombination (Rudiger et al., 1995). These data strongly suggest that the *Alu* core sequence itself can promote DNA strand exchange and genomic rearrangement. The similarity of the *Alu* core sequence to  $\chi$ , which has been shown to stimulate recBCD-mediated recombination in *E. coli*, is strongly supportive of a model in which the common 26-bp core sequence could, in some situations, provide the

binding site for the proteins catalyzing HR. In support of this hypothesis, it was shown recently that sequences closely homologous to translin DNA-binding sequences were identified within *Alu* consensus sequences (Jeffs et al., 1998). The translin protein associates with specific sequences and is involved in local DNA unwinding. Significantly, it has been shown that translin binding makes chromosomal regions more susceptible to both nuclease and recombinase activities (Aoki et al., 1995), making its close sequence identity within the *Alu* consensus provocative. In addition, the sequence similarity within the consensus sequence and the established prokaryotic model of recombination and DNA DSB repair provides further support to implicate a more direct role for *Alu* elements in chromosomal rearrangements. If such a hypothesis is correct, then it seems reasonable that *Alu* repeats are frequent sites for recombination, not only because of their abundance but also because they can serve as binding sites for RAD proteins. In such a model, *Alu* may thus both mediate and facilitate HR.

#### THE ROLE OF *Alu* ELEMENTS IN EVOLUTION

The ubiquitous presence of *Alu* elements in the primate genome and some of their sequence-dependent properties have led various investigators to ascribe functional roles for *Alu* elements in evolution: *Alu* repeats play a role as ubiquitous DNA methylation signals regulating gene expression and imprinting in the human genome (Schmid, 1996). The *Alu* transcript may activate protein translation by inhibiting dsRNA-regulated protein kinase PRK in response to cell stress, such as viral infection and heat shock (Chu et al., 1998; Schmid, 1998). It has been shown that retrotransposition of *Alu* elements can insert novel sequences within or near genes, leading to altered transcriptional expression (Norris et al., 1995; Vansant and Reynolds, 1995; Britten, 1996). Insertion can also lead to dysregulation of post-transcriptional events such as polyadenylation, splicing, and the creation of new exonic sequences (Schmid, 1996).

#### *Alu* REPEATS ARE IMPLICATED AS MEDIATORS OF GENOMIC REARRANGEMENT

When *Alu* elements are present on recombinant DNA constructs, they have been shown to increase the recombination frequency between vector DNA and the host genomic loci (Kato et al., 1986; Wallenburg et al., 1987; Kang et al., 1999), which is in

keeping with this sequence element's role as a "hot spot" for recombination events. If unequal crossing over occurs between *Alu* elements intrachromosomally (in cis), it can result in deletion or duplication of intervening sequences. When *Alu*-mediated recombination occurs interchromosomally (in trans), it can lead to more complex chromosomal rearrangements such as translocations (Deininger and Batzer, 1999). Table 1 is an updated summary of the constitutive and acquired diseases in man that have been attributed to the presence of *Alu* repeats, insertions, or unequal recombination [the original summary was published by Labuda et al. (1995)].

*Alu* repeat-mediated mutations in meiotic cells, the result of unequal HR leading to deletions or duplications, bring about a number of constitutional diseases (Labuda, 1995; Hartevelde et al., 1997; Tvrdik et al., 1998; Huie et al., 1999). Sixteen *Alu*-based insertion mutations were identified in the Human Genetic Mutation Database (Deininger and Batzer, 1999) (14,374 entries), suggesting that *Alu* repeats could contribute to 0.1% of human genetic diseases attributed to insertion alone (Janicic et al., 1995; Labuda, 1995; Rowe et al., 1995; Miki et al., 1996). *Alu* repeats have been implicated in the etiology of some recurrent somatic chromosome rearrangements such as the Philadelphia translocation in some cases of chronic myelogenous leukemia (CML) and complex *BCR/ABL1* rearrangements (Rogers, 1985; de Klein et al., 1986; Koduru et al., 1993; Chisoe et al., 1995; Zhang et al., 1995; Morris et al., 1996; Jeffs et al., 1998), Ph-positive ALL (Chen et al., 1989a,b; van der Feltz et al., 1989; Papadopoulos et al., 1990), as well as partial duplication of the *MLL* gene in acute myeloid leukemia, given that *Alu* repetitive sequences have been identified at or near the junction of translocation breakpoints (in most of the cases on both sides of the breakpoints).

Based on the preceding observations, it seems probable that *Alu* repeats may facilitate genomic rearrangement in some in vivo situations. The very high densities of *Alu* repeats at regions most frequently involved in chromosomal rearrangements are particularly compelling in this regard (Figs. 1 and 2).

#### MECHANISTIC IMPLICATIONS OF DELETIONS ASSOCIATED WITH RECURRENT CHROMOSOMAL REARRANGEMENTS

We and others have recently demonstrated that in about 10% of leukemia-associated chromosomal rearrangements, there are deletions immediately

TABLE I. Disease-Causing Chromosomal Rearrangements Resulting From *Alu*-Mediated Recombination

Locus	Disease	References
Disease-causing deletions due to <i>Alu</i> -mediated recombination		
<i>BRCA1</i>	Breast and ovarian cancer	Swensen et al., 1997; Montagna et al., 1999; Rohlfis et al., 2000
<i>RB1</i>	Glioma brain tumors	Rothberg et al., 1997
<i>PDGFB</i>	Meningioma	Smidt et al., 1990
<i>MLH1</i>	Familial colorectal cancer	Mauillon et al., 1996
<i>MSH2</i> human DNA mismatch repair gene	Hereditary nonpolyposis colorectal cancer	Marshall et al., 1996
<i>N</i> -Acetylgalactosamine-6-sulfatase gene ( <i>GALNS</i> )	Mucopolysaccharidosis type IVA	Hori et al., 1995
Alpha-galactosidase A	Fabry disease	Kornreich et al., 1990
$\alpha$ -Globin gene cluster	$\alpha$ -Thalassemia	Nicholls et al., 1987; Hartevelde et al., 1997; Hattori et al., 1999
$\beta$ -Globin gene cluster	$\delta\beta$ -Thalassemia, $\gamma\delta\beta$ -thalassemia, HPFH2	Henthorn et al., 1990
C1 inhibitor gene	Hereditary angioedema	Stoppa-Lyonnet et al., 1990
Adenosine deaminase ( <i>ADA</i> )	<i>ADA</i> -deficient SCID	Markert et al., 1988
Low-density lipoprotein receptor	Familial hypercholesterolemia	Hobbs et al., 1986; Lehrman et al., 1986
Platelet fibrinogen receptor	Glanzmann's thrombasthenia (GT)	Burk et al., 1991
Apolipoprotein B gene	Hypobetalipoproteinemia	Huang et al., 1989
von Willebrand factor A	von Willebrand disease	Bernardi et al., 1993; Ferreira et al., 1993; Zhang et al., 1994
<i>FANCA</i>	Fanconi anemia	Levrin et al., 1998; Morgan et al., 1999
$\beta$ -Hexosaminidase $\alpha$ -chain gene	Tay-Sachs disease	Myerowitz and Hogikyan, 1987
<i>HEXB</i>	Sandhoff	Neote et al., 1990
Hypoxanthine phosphoribosyltransferase	Lesch-Nyhan (LN)	Tvrdek et al., 1998
$\alpha$ -Glucosidase	Glycogen storage disease type II (GSDII)	Huie et al., 1999
Disease-causing duplication due to <i>Alu</i> -mediated recombination		
C1 inhibitor gene	Hereditary angioedema	Stoppa-Lyonnet et al., 1990
Dystrophin	Duchenne muscular dystrophy	Hu et al., 1991
Lysyl hydroxylase	Ehlers-Danlos syndrome	Pousi et al., 1994
<i>MLL</i> dup(11)(q23)	Acute myeloid and lymphoid leukemia	Bernard et al., 1995; Schnittger et al., 1998; Strout et al., 1998; Wiedemann et al., 1999
Disease-causing translocations due to <i>Alu</i> -mediated recombination		
t(11;22)	Ewing sarcoma	Obata et al., 1999
Ph translocation t(9;22)	CML, Ph-positive acute leukemias	de Klein et al., 1986; Chen et al., 1989a; Papadopoulos et al., 1990; Toth and Jurka, 1994; Chissoe et al., 1995; Martinelli et al., 2000
Variant Ph translocation	CML	Koduru et al., 1993; Morris et al., 1996; Jeffs et al., 1998
t(4;11)	Treatment-related acute lymphoblastic leukemia	Megonigal et al., 1997
t(14;18)	Follicular lymphomas (FLs)	Buchonnet et al., 2000
t(9;11)	De novo acute myeloid leukemia	Super et al., 1997
t(2;8)	Burkitt lymphoma cell line	Kato et al., 1991
t(14;19)	Chronic lymphocytic leukemia (CLL)	Ohno et al., 1993
t(11;22)c	der(22) syndrome	Hill et al., 2000*

\*In contrast to the findings of Hill et al. (2000), it has been suggested recently that palindromic AT-rich repeats are responsible for the constitutional t(11;22) (Kurahashi et al., 2000)

adjacent to the breakpoint regions (Sinclair et al., 2000; Kolomietz et al., 2001). Our retrospective study of 250 cases of CML patients in chronic phase and blast crisis and 13 cases of Ph+ ALL patients indicated that 23 CML patients (9.2%) and 1 ALL patient (7.7%) exhibited a hybridization

pattern consistent with deletion of two regions: proximal to the rearranged *ABL* gene and distal to the *BCR* gene on the der(9) chromosome. The observation that deletions are associated with both Ph-positive CML and ALL leukemias suggested that other leukemia-associated genomic rearrange-






Figure 2. Results of computer-based DNA sequence analysis with the purpose of identification of repeat density for chromosome 22 were correlated with breakpoints of recurrent chromosomal aberrations to confirm genome-wide analysis based on in situ hybridization methods of mapping of *Alu* repeats (Korenberg and Rykowski, 1988). Cancer breakpoints on chromosome 22 map to the regions of clusters of *Alu* repeats with *Alu* density higher than 20%. In silico analysis of the complete genome sequence now permits detailed mapping of repetitive se-

quences throughout the human genome to improve our understanding of its role. A number of computer-aided programs such as Repeat Masker (<http://ftp.genome.washington.edu/c/s.dll/RepeatMasker>), Censor (<http://www.girinst.org/>), and BLAST (*alu* db) are widely used for the identification of repeats. These programs allow the user to identify and characterize different classes of repeats within a given genomic sequence and to map precisely their relative distributions and densities.

ments may also be subject to deletion in the break point flanking regions. This idea was confirmed by the detection of a deletion of the 3' region of the

*CBFB* gene in 20% ( $n = 10$ ) of AML-M4 patients with *inv(16)(p13q22)*. However, fluorescence in situ hybridization (FISH) analysis of two groups of




Figure 3. Correlation between the presence of the deletions associated with chromosomal rearrangements and the density of the *Alu* repeats in the genes involved in this rearrangement and their flanking regions.

patients, one with AML-M3 ( $n = 30$ ) and the other with AML-M2 ( $n = 14$ ), showed no deletions associated with the  $t(15;17)(q22;q11.2)$  and  $t(8;21)(q22;q22)$ , respectively. In contrast, others (Godon et al., 2000) have identified deletions associated with the  $t(8;21)$ . These findings suggest that such deletions are not disease-specific per se, but are more likely determined by the factors that influence the somatic process of chromosomal rearrangement itself. The fact that deletions are associated with different chromosomal rearrangements suggests that there might be a common mechanism for deletion formation, which is sequence-specific.

To investigate this possibility, DNA sequences of the genes involved in the chromosomal rearrangements and the adjacent flanking regions were submitted to the repeat identification programs Censor ([http://www.girinst.org/Censor\\_Server-Data\\_Entry\\_Forms.html](http://www.girinst.org/Censor_Server-Data_Entry_Forms.html)) and Repeat Masker (<http://ftp.genome.washington.edu/cgi-bin/RepeatMasker>), to identify the density of *Alu* repeats within the regions. Our findings suggest a strong association between the propensity to undergo deletion and an elevated density of *Alu* repeats in the chromosomal regions subject to rearrangement. A much higher density of *Alu* repeats was observed in the DNA

sequences of the genes involved in the  $t(9;22)$  (*ABL* and *BCR*) and  $inv(16)$  (*CBFB* and *MYH11*) (over 20%) than in the genes involved in the  $t(15;17)$  (*PML* and *RARA*) (5–10%) (Fig. 3). It has recently been confirmed by computer-based analysis of *BCR* and *ABL* genes that *Alu* elements are the predominant repeat type in the *BCR* and *ABL1* genes and their distribution is nonrandom (Jeffs et al., 2001). This analysis revealed significant clusters of *Alu* elements at the 5' end of both the *BCR* and the *ABL* genes. Sequence variation of *Alu* elements within subfamilies and a nucleotide difference between individuals have been reported (Chisoe et al., 1995). Concerning this observation, it has been suggested that certain sequence configurations of the *Alu* repeats are more susceptible to recombination than others. Although it remains unproved, it is tempting to speculate that in some individuals such nonrandom distribution may, in fact, play an important role in facilitating deletion formation, associated with chromosomal rearrangements, in neighboring genes that have close proximity to a flanking high density of *Alu* repeats.

Sinclair et al. (2000) have provided a detailed map of the regions that are subject to deletion on the derivative chromosomes 9 and 22. The dele-

tions vary in size from several hundred kilobases to 3–4 megabases and have variable breakpoints. It is well established that deletions of a few thousand base pairs will often be created by the action of endogenous endo- and exonucleases on the broken ends following DSBs (Szostak et al., 1983; Zucman-Rossi et al., 1998). Other studies have suggested that the small deletions ranging from several to 1,500 nucleotides that accompany translocation formation result from the base-pairing of fortuitously located direct repeats (microhomology spanned from 11–18 bp) flanking the DNA broken ends. When the direct repeats are joined together, the intervening DNA fragment will be deleted (Salganik and Dianov, 1992). *Alu* elements might serve as sites for unequal homologous crossing over, consequently scrambling their flanking genomic sequences (Schmid, 1996), leading to their deletion and/or amplification and possibly leading to the creation of the very large deletions reported by our group (Kolomietz et al., 2001) and others (Sinclair et al., 2000). Alternatively, *Alu* elements could partially recombine and hold chromosomal regions near each other. Consequently, rearrangements would be more likely to occur in the regions where *Alu* elements are present at higher densities (Chen et al., 1989a). Translocations may result from HR between *Alu* elements within the genomic sequences (Morris et al., 1996). However, it is more likely that the chance of rearrangement increases because attraction between *Alu* elements on nonhomologous chromosomes initially brings two genomic sites together. *Alu* sequences may base-pair following DSBs, either by means of physical juxtaposition arising from nonrandom nuclear compartmentalization (Neves et al., 1999) or by serving as substrates for HR after DSB. Rearrangement between homologous sequences could take place, even if there is no homology at the breakpoints because of spatial configuration of chromatin in nuclei (Neves et al., 1999).

It has also been suggested that transcriptionally active chromatin is more open in structure and thereby is prone to recombine more frequently than inactive chromatin (Datta and Jinks-Robertson, 1995). If this model is correct, then the open chromatin structure associated with the active transcription of the *BCR* and *ABL* genes in myeloid cells could increase the probability that these genes will associate, so that *Alu*-mediated recombination can then take place (Martinelli et al., 2000).

## GENETIC INSTABILITY AND DISEASES WITH INCREASED FREQUENCY OF HOMOLOGOUS RECOMBINATION

The high density of *Alu* sequences in the human genome and the evidence that sequence-dependent HR is a major DNA repair pathway indicate that there must be a mechanism capable of regulating HR and preventing it from causing chromosomal instability. The fact that several genetic diseases have a DNA instability phenotype and a high frequency of carcinogenesis suggests a correlation between those mutations that result in a higher recombination frequency, genetic instability, and cancer predisposition (Bishop and Schiestl, 2000; Hoeijmakers, 2001). These diseases include ataxia telangiectasia (AT), Li–Fraumeni syndrome, Nijmegen breakage syndrome, Bloom syndrome (BS), Werner syndrome (WS), Fanconi anemia, Lynch syndromes I and II, Wiscott–Aldrich syndrome, and xeroderma pigmentosum.

*BLM* and *WRN*, the genes that are mutated in the cancer-prone disorders BS and WS, respectively, belong to the family of RecQ helicase homologs (Ellis et al., 1995; Yu et al., 1996; Enomoto, 2001). These two syndromes are characterized by a high degree of genomic instability, including chromosomal breaks, multiple large deletions, and translocations; moreover, cells derived from BS and WS patients show defects in DNA replication (Enomoto, 2001). It has become clear that DSBs could arise by active processing of stalled replication forks by a specific enzyme system (Michel et al., 1997; Flores-Rozas and Kolodner, 2000). HR plays an important role in the repair of stalled or broken replication forks, leading to the reinitiation of replication (Shinohara and Ogawa, 1995; Osman and Subramani, 1998; Flores-Rozas and Kolodner, 2000). Defects in the processing of stalled replication forks could lead to aberrant recombination events resulting in genetic instability (Enomoto, 2001). Recent studies on *BLM*, *WRN*, and the RecQ homolog of *S. cerevisiae*, *Sgs1*, indicate that these RecQ homologs function in a pathway from the DNA replication checkpoint to HR (Karow et al., 2000b). It has also been shown that *BLM* disrupts potentially recombinogenic molecules that arise at sites of stalled replication forks, suggesting a role of *BLM* in suppression of hyperrecombination (Karow et al., 2000a).

AT cells have an abnormally high rate of recombination and defective repair around the site of DNA strand breakage (Meyn, 1993; Luo et al., 1996). It is noteworthy that the knockout *Atm* gene

mouse model displays genetic instability and a significantly elevated frequency of intrachromosomal HR, resulting in deletion events (Bishop et al., 2000). These authors concluded that, because 25% of our genome consists of repeated elements, genomic instability attributed to an increased number of repeats and error-prone HR between such repeats might contribute to carcinogenesis in AT patients. The controlling role of *ATM* in HR-mediated repair (Morrison et al., 2000) provides the link between the recombinational repair machinery and the DNA damage checkpoint apparatus.

Mismatch repair (MMR) machinery is also known to suppress HR and microsatellite instability in both prokaryotic and eukaryotic cells (Raysiguier et al., 1989; de Wind et al., 1995; Prolla, 1998; Harfe and Jinks-Robertson, 2000a). Recently, it was shown that MMR is also involved in suppression of other genomic instabilities such as gene amplification and chromosomal DNA integration (Lin et al., 2001).

HR involves the pairing of complementary single strands derived from two different duplexes, resulting in a heteroduplex DNA molecule. If the recombining duplexes are not identical in sequence, mismatches contained in the resulting heteroduplex are recognized and processed by MMR machinery, triggering one of two events: simple mismatch correction or complete abortion of the recombination event. In addition, MMR proteins are involved in the removal of the nonhomologous single-strand tails that are generated during the recombination processes (Harfe and Jinks-Robertson, 2000a).

The anti-recombination activity of the eukaryotic MMR machinery has been shown to inhibit recombination between diverged sequences both in mitosis and in meiosis (Harfe and Jinks-Robertson, 2000b). It has been demonstrated that a single mismatch is sufficient to inhibit recombination in an MMR-dependent manner, with additional mismatches having a cumulative negative effect on recombination rates (Datta et al., 1997; Chen and Jinks-Robertson, 1998). Moreover, sequence divergence not only inhibits HR but also reduces the efficiency of the single-strand annealing recombination pathway (Sugawara et al., 1997). Due to mismatch-triggered anti-recombination activity, eukaryotic MMR proteins inhibit ectopic interactions between divergent, repetitive DNA sequences such as *Alu* repeats during both mitosis and meiosis, and thereby promote genome stability by preventing deleterious genome rearrangements (Harfe and Jinks-Robertson, 2000a). Interestingly,

mitotic recombination in yeast is more sensitive (in an MMR-dependent manner) to low levels of sequence divergence than is meiotic recombination (Chen and Jinks-Robertson, 1999), which may help reinforce the strong bias for sister chromatid vs. homologous chromosome interactions during mitosis. The observations made in yeast are likely to be relevant to the mismatch-associated regulation of recombination in higher eukaryotes (Harfe and Jinks-Robertson, 2000a,b).

Recent studies suggest that normal cells have stringent mechanisms for controlling HR, and that the altered expression of genes involved in DNA repair and DNA-damage sensing can lead to error-prone recombination and chromosomal rearrangement. Such changes may permit cells that have undergone these events to survive by escaping apoptosis (Richardson et al., 1998).

#### INFLUENCE OF CHROMATIN STRUCTURE AND REPEAT-MEDIATED REARRANGEMENT

Given the large number of repetitive elements in the human genome, it would be expected that recombination between these sequences could lead to gross genomic rearrangements, loss of genetic information, and general genetic instability (Bishop and Schiestl, 2000). Indeed, it has been shown that regions enriched in *Alu* repeat sequences do display an elevated incidence of genomic instability (Calabretta et al., 1982). However, cells must possess the mechanisms that limit and control the recombination process. The exquisite sensitivity of the HR process to any deviation away from perfect sequence homology allows cells to suppress recombination between divergent repetitive elements. The likelihood that HR will take place is a function of both the length of uninterrupted homology and the extent of perfect homology between two repetitive sequence elements (Wallenburg et al., 1987). Efficient recombination appears to require between 134 and 232 bp of uninterrupted homology; the recombination rate is primarily governed by the availability of contiguous uninterrupted homology (Waldman and Liskay, 1988).

It has also been shown that the frequency of *Alu*-mediated deletions in undamaged DNA is relatively low, with deletion rates in sperm of  $3 \times 10^{-7}$  and  $7 \times 10^{-7}$  and maximum frequencies of somatic mutation of less than  $10^{-6}$  per cell (Hollies et al., 2001). It seems that chromosomal DNA breaks may be necessary inducers of recombination and therefore increase the frequency of HR (Rouet et al., 1994) and chromosomal rearrangements.

Chromatin structure creates an additional level of complexity for HR-mediated DSB repair. Both local and topologic differences in chromatin structure might strongly influence the likelihood of the search for homology being successful (Kanaar et al., 1998). Chromatin structure associated with *A/u* repeats may play a role in the suppression of HR between *A/u* repeats by complicating the process of homology search. Computational analysis of the organization of chromatin associated with *A/u* repeats indicated that the sequences are typically associated with rotationally positioned nucleosomes. Several lines of evidence (Englander et al., 1993; Englander and Howard, 1995) indicate that *A/u* repeat monomer elements are capable of setting the rotational phase of nucleosomes or nucleosome-like particles. Rotational positioning for *A/u* elements as a group reflects DNA sequence features intrinsic to these repeats (Englander and Howard, 1995). It is reasonable to speculate that nucleosome positioning over the *A/u* A-box promoter element might have a contributory role in suppression of HR between *A/u* repeats.

The relative position of different chromatin domains within the three-dimensional structure of the nucleus may play a role in mediating the recombination process. Nuclear topographic analysis of the relative position of the *ABL* and *BCR* genes has provided evidence for their close proximity in specific phases of the cell cycle (Koduru and Chaganti, 1988; Kozubek et al., 1997; Neves et al., 1999). Thus, physical juxtaposition arising from nonrandom nuclear compartmentalization may strongly facilitate the passive process of homology search and increase the chance of *A/u*-mediated rearrangement taking place.

Normal cells have mechanisms to control and suppress the occurrence of chromosomal rearrangements created by inappropriate repair of DSBs. Recent data suggest that the combination of factors—including the presence of sufficient uninterrupted homologous sequences of repeated elements, their topologic proximity resulting from the nuclear compartmentalization of the genes, and mutations in genes involved in DNA repair and DNA-damage sensing—may create the conditions in which error-prone recombination may lead to chromosomal rearrangement. *A/u* repeats play an important role in initiating such recurrent chromosomal rearrangements, in spite of their role in suppressing “promiscuous” recombination.

## FUTURE DIRECTIONS

With the near completion of the entire Human Genome Sequence, it will be possible to create more detailed maps of the distribution and the density of repetitive sequences. Such maps will offer opportunities for better comparisons between the developing cancer breakpoints map [Breakpoint Map of Recurrent Chromosome Aberrations at CGAP (the Cancer Genome Anatomy Project) <http://www.ncbi.nlm.nih.gov/CCAP/mitelsum.cgi>] and the localization of repetitive DNA. In this way, it will be possible to develop a better appreciation of the role of sequences such as *A/u* in chromosomal rearrangement and tumorigenesis. In addition, an improved understanding of the topologic architecture of chromatin and the role of both homologous and nonhomologous DNA repair will help to determine why chromosomal rearrangements preferentially take place in certain regions of the human genome.

## ACKNOWLEDGMENTS

We are indebted to Dr. Mark Minden, Paula Marrano, and Shawn Brennan for critically reviewing this manuscript.

## REFERENCES

- Adachi M, Tsujimoto Y. 1990. Potential Z-DNA elements surround the breakpoints of chromosome translocation within the 5' flanking region of bcl-2 gene. *Oncogene* 5:1653–1657.
- Aoki K, Suzuki K, Sugano T, Tasaka T, Nakahara K, Kuge O, Omori A, Kasai M. 1995. A novel gene, Translin, encodes a recombination hotspot binding protein associated with chromosomal translocations. *Nat Genet* 10:167–174.
- Aplan PD. 1999. Mechanism of leukemogenesis: Chromosomal translocations. American Society of Hematology annual meeting.
- Aplan PD, Chervinsky DS, Stanulla M, Burhans WC. 1996. Site-specific DNA cleavage within the MLL breakpoint cluster region induced by topoisomerase II inhibitors. *Blood* 87:2649–2658.
- Benson FE, Baumann P, West SC. 1998. Synergistic actions of Rad51 and Rad52 in recombination and DNA repair. *Nature* 391:401–404.
- Bernard OA, Romana SP, Schichman SA, Mauchauffe M, Jonveaux P, Berger R. 1995. Partial duplication of HRX in acute leukemia with trisomy 11. *Leukemia* 9:1487–1490.
- Bernardi F, Patracchini P, Gemmati D, Pinotti M, Schwienbacher C, Ballerini G, Marchetti G. 1993. In-frame deletion of von Willebrand factor A domains in a dominant type of von Willebrand disease. *Hum Mol Genet* 2:545–548.
- Bishop AJ, Schiestl RH. 2000. Homologous recombination as a mechanism for genome rearrangements: environmental and genetic effects. *Hum Mol Genet* 9:2427–2434.
- Bishop AJ, Barlow C, Wynshaw-Boris AJ, Schiestl RH. 2000. Atm deficiency causes an increased frequency of intrachromosomal homologous recombination in mice. *Cancer Res* 60:395–399.
- Bishop DK, Park D, Xu L, Kleckner N. 1992. DMC1: a meiosis-specific yeast homolog of E. coli recA required for recombination, synaptonemal complex formation, and cell cycle progression. *Cell* 69:439–456.
- Boehm T, Mengle-Gaw L, Kees UR, Spurr N, Lavenir I, Forster A, Rabbitts TH. 1989. Alternating purine-pyrimidine tracts may promote chromosomal translocations seen in a variety of human lymphoid tumours. *EMBO J* 8:2621–2631.
- Britten RJ. 1996. DNA sequence insertion and evolutionary variation in gene regulation. *Proc Natl Acad Sci USA* 93:9374–9377.
- Buchhop S, Gibson MK, Wang XW, Wagner P, Sturzbecher HW,



- Harris CC. 1997. Interaction of p53 with the human Rad51 protein. *Nucleic Acids Res* 25:3868–3874.
- Buchonnet G, Lenain P, Ruminy P, Lepretre S, Stamatoullas A, Parmentier F, Jardin F, Duval C, Tilly H, Bastard C. 2000. Characterisation of BCL2-JH rearrangements in follicular lymphoma: PCR detection of 3' BCL2 breakpoints and evidence of a new cluster. *Leukemia* 14:1563–1569.
- Burk CD, Newman PJ, Lyman S, Gill J, Collier BS, Poncz M. 1991. A deletion in the gene for glycoprotein IIb associated with Glanzmann's thrombasthenia. *J Clin Invest* 87:270–276.
- Calabretta B, Robberson DL, Barrera-Saldana HA, Lambrou TP, Saunders GF. 1982. Genome instability in a region of human DNA enriched in Alu repeat sequences. *Nature* 296:219–225.
- Chen G, Yuan SS, Liu W, Xu Y, Trujillo K, Song B, Cong F, Goff SP, Wu Y, Arlinghaus R, Baltimore D, Gasser PJ, Park MS, Sung P, Lee EY. 1999. Radiation-induced assembly of Rad51 and Rad52 recombination complex requires ATM and c-Abl. *J Biol Chem* 274:12748–12752.
- Chen SJ, Chen Z, d'Auriol L, Le Coniat M, Grausz D, Berger R. 1989a. Ph1+ bcr-acute leukemias: implication of Alu sequences in a chromosomal translocation occurring in the new cluster region within the BCR gene. *Oncogene* 4:195–202.
- Chen SJ, Chen Z, Font MP, d'Auriol L, Larsen CJ, Berger R. 1989b. Structural alterations of the BCR and ABL genes in Ph1 positive acute leukemias with rearrangements in the BCR gene first intron: further evidence implicating Alu sequences in the chromosome translocation. *Nucleic Acids Res* 17:7631–7642.
- Chen W, Jinks-Robertson S. 1998. Mismatch repair proteins regulate heteroduplex formation during mitotic recombination in yeast. *Mol Cell Biol* 18:6525–6537.
- Chen W, Jinks-Robertson S. 1999. The role of the mismatch repair machinery in regulating mitotic and meiotic recombination between diverged sequences in yeast. *Genetics* 151:1299–1313.
- Chissoc SL, Bodenteich A, Wang YF, Wang YP, Burian D, Clifton SW, Crabtree J, Freeman A, Iyer K, Jian L, Ma Y, McLaury HJ, Pan HQ, Sarhan OH, Toth S, Wang Z, Zhang G, Heisterkamp N, Groffen J, Ro BA. 1995. Sequence and analysis of the human ABL gene, the BCR gene, and regions involved in the Philadelphia chromosomal translocation. *Genomics* 27:67–82.
- Chu WM, Ballard R, Carpick BW, Williams BR, Schmid CW. 1998. Potential Alu function: regulation of the activity of double-stranded RNA-activated kinase PKR. *Mol Cell Biol* 18:58–68.
- Craig JM, Bickmore WA. 1993. Chromosome bands—flavours to savour. *BioEssays* 15:349–354.
- Critchlow SE, Jackson SP. 1998. DNA end-joining: from yeast to man. *Trends Biochem Sci* 23:394–398.
- Datta A, Jinks-Robertson S. 1995. Association of increased spontaneous mutation rates with high levels of transcription in yeast. *Science* 268:1616–1619.
- Datta A, Hendrix M, Lipsitch M, Jinks-Robertson S. 1997. Dual roles for DNA sequence identity and the mismatch repair system in the regulation of mitotic crossing-over in yeast. *Proc Natl Acad Sci USA* 94:9757–9762.
- Debatisse M, Coquelle A, Toledo F, Buttin G. 1998. Gene amplification mechanisms: the role of fragile sites. *Recent Results Cancer Res* 154:216–226.
- Deininger PL, Batzer MA. 1999. Alu repeats and human disease. *Mol Genet Metab* 67:183–193.
- de Klein A, van Agthoven T, Groffen C, Heisterkamp N, Groffen J, Grosveld G. 1986. Molecular analysis of both translocation products of a Philadelphia-positive CML patient. *Nucleic Acids Res* 14:7071–7082.
- de Wind N, Dekker M, Berns A, Radman M, te Riele H. 1995. Inactivation of the mouse Msh2 gene results in mismatch repair deficiency, methylation tolerance, hyperrecombination, and predisposition to cancer. *Cell* 82:321–330.
- Dixon DA, Kowalczykowski SC. 1991. Homologous pairing in vitro stimulated by the recombination hotspot, chi. *Cell* 66:361–371.
- Dixon DA, Kowalczykowski SC. 1993. The recombination hotspot chi is a regulatory sequence that acts by attenuating the nuclease activity of the E. coli RecBCD enzyme. *Cell* 73:87–96.
- Dolganov GM, Maser RS, Novikov A, Tosto L, Chong S, Bressan DA, Petrini JH. 1996. Human Rad50 is physically associated with human Mre11: identification of a conserved multiprotein complex implicated in recombinational DNA repair. *Mol Cell Biol* 16:4832–4841.
- Ellis NA, Groden J, Ye TZ, Straughen J, Lennon DJ, Ciocci S, Proytcheva M, German J. 1995. The Bloom's syndrome gene product is homologous to RecQ helicases. *Cell* 83:655–666.
- Englander EW, Howard BH. 1995. Nucleosome positioning by human Alu elements in chromatin. *J Biol Chem* 270:10091–10096.
- Englander EW, Wolffe AP, Howard BH. 1993. Nucleosome interactions with a human Alu element. Transcriptional repression and effects of template methylation. *J Biol Chem* 268:19565–19573.
- Enomoto T. 2001. Functions of RecQ family helicases: possible involvement of Bloom's and Werner's syndrome gene products in guarding genome integrity during DNA replication. *J Biochem* 129:501–507.
- Essers J, van Steeg H, de Wit J, Swagemakers SM, Vermeij M, Hoeijmakers JH, Kanaar R. 2000. Homologous and non-homologous recombination differentially affect DNA damage repair in mice. *EMBO J* 19:1703–1710.
- Featherstone C, Jackson SP. 1999. Ku, a DNA repair protein with multiple cellular functions. *Mutat Res* 434:3–15.
- Ferguson DO, Alt FW. 2001. DNA double strand break repair and chromosomal translocation: lessons from animal models. *Oncogene* 20:5572–5579.
- Ferreira V, Assouline Z, Schwachtgen JL, Bahnak BR, Meyer D, Kerbiriou-Nabias D. 1993. The role of the 5'-flanking region in the cell-specific transcription of the human von Willebrand factor gene. *Biochem J* 293:641–648.
- Finger LR, Harvey RC, Moore RC, Showe LC, Croce CM. 1986. A common mechanism of chromosomal translocation in T- and B-cell neoplasia. *Science* 234:982–985.
- Flores-Rozas H, Kolodner RD. 2000. Links between replication, recombination and genome instability in eukaryotes. *Trends Biochem Sci* 25:196–200.
- Gasior SL, Wong AK, Kora Y, Shinohara A, Bishop DK. 1998. Rad52 associates with RPA and functions with rad55 and rad57 to assemble meiotic recombination complexes. *Genes Dev* 12:2208–2221.
- Glover TW, Stein CK. 1988. Chromosome breakage and recombination at fragile sites. *Am J Hum Genet* 43:265–273.
- Godon C, Proffitt J, Dastugue N. 2000. Large deletions 5' to the ETO breakpoint are observed in 10% of patients with t(8;21)-positive AML. American Society of Hematology annual meeting.
- Habu T, Taki T, West A, Nishimune Y, Morita T. 1996. The mouse and human homologs of DMC1, the yeast meiosis-specific homologous recombination gene, have a common unique form of exon-skipped transcript in meiosis. *Nucleic Acids Res* 24:470–477.
- Harfe BD, Jinks-Robertson S. 2000a. DNA mismatch repair and genetic instability. *Annu Rev Genet* 34:359–399.
- Harfe BD, Jinks-Robertson S. 2000b. Mismatch repair proteins and mitotic genome stability. *Mutat Res* 451:151–167.
- Harteveld KL, Losekoot M, Fodde R, Giordano PC, Bernini LF. 1997. The involvement of Alu repeats in recombination events at the alpha-globin gene cluster: characterization of two alpha-zero-thalassaemia deletion breakpoints. *Hum Genet* 99:528–534.
- Hattori Y, Okayama N, Ohba Y, Yamashiro Y, Yamamoto K, Tsukimoto I, Kohakura M. 1999. The precise breakpoints of a Filipino-type alpha-thalassemia-1 deletion found in two Japanese. *Hemoglobin* 23:239–248.
- Henthorn PS, Smithies O, Mager DL. 1990. Molecular analysis of deletions in the human beta-globin gene cluster: deletion junctions and locations of breakpoints. *Genomics* 6:226–237.
- Hill AS, Foot NJ, Chaplin TL, Young BD. 2000. The most frequent constitutional translocation in humans, the t(11;22)(q23;q11) is due to a highly specific alu-mediated recombination. *Hum Mol Genet* 9:1525–1532.
- Hobbs HH, Brown MS, Goldstein JL, Russell DW. 1986. Deletion of exon encoding cysteine-rich repeat of low density lipoprotein receptor alters its binding specificity in a subject with familial hypercholesterolemia. *J Biol Chem* 261:13114–13120.
- Hoeijmakers JH. 2001. Genome maintenance mechanisms for preventing cancer. *Nature* 411:366–374.
- Hollies CR, Monckton DG, Jeffreys AJ. 2001. Attempts to detect retrotransposition and de novo deletion of Alus and other dispersed repeats at specific loci in the human genome. *Eur J Hum Genet* 9:143–146.
- Holmquist GP. 1992. Chromosome bands, their chromatin flavors, and their functional features. *Am J Hum Genet* 51:17–37.
- Hori T, Tomatsu S, Nakashima Y, Uchiyama A, Fukuda S, Sukegawa K, Shimozawa N, Suzuki Y, Kondo N, Horiuchi T. 1995. Mucopolysaccharidosis type IVA: common double deletion in the N-acetylgalactosamine-6-sulfatase gene (GALNS). *Genomics* 26:535–542.
- Hu XY, Ray PN, Worton RG. 1991. Mechanisms of tandem duplication in the Duchenne muscular dystrophy gene include both

- homologous and nonhomologous intrachromosomal recombination. *EMBO J* 10:2471–2477.
- Huang LS, Ripps ME, Korman SH, Deckelbaum RJ, Breslow JL. 1989. Hypobetalipoproteinemia due to an apolipoprotein B gene exon 21 deletion derived by Alu-Alu recombination. *J Biol Chem* 264:11394–11400.
- Huic ML, Shanske AL, Kasper JS, Marion RW, Hirschhorn R. 1999. A large Alu-mediated deletion, identified by PCR, as the molecular basis for glycogen storage disease type II (GSDII). *Hum Genet* 104:94–98.
- Janicic N, Pausova Z, Cole DE, Hendy GN. 1995. Insertion of an Alu sequence in the Ca(2+)-sensing receptor gene in familial hypocalciuric hypercalcemia and neonatal severe hyperparathyroidism. *Am J Hum Genet* 56:880–886.
- Jeffs AR, Benjes SM, Smith TL, Sowerby SJ, Morris CM. 1998. The BCR gene recombines preferentially with Alu elements in complex BCR-ABL translocations of chronic myeloid leukaemia. *Hum Mol Genet* 7:767–776.
- Jeffs AR, Wells E, Morris CM. 2001. Nonrandom distribution of interspersed repeat elements in the BCR and ABL1 genes and its relation to breakpoint cluster regions. *Genes Chromosomes Cancer* 32:144–154.
- Johnson RD, Jasin M. 2000. Sister chromatid gene conversion is a prominent double-strand break repair pathway in mammalian cells. *EMBO J* 19:3398–3407.
- Kadyk LC, Hartwell LH. 1992. Sister chromatids are preferred over homologs as substrates for recombinational repair in *Saccharomyces cerevisiae*. *Genetics* 132:387–402.
- Kanaar R, Hoeijmakers JH, van Gent DC. 1998. Molecular mechanisms of DNA double strand break repair. *Trends Cell Biol* 8:483–489.
- Kang YK, Park JS, Lee CS, Yeom YI, Chung AS, Lee KK. 1999. Efficient integration of short interspersed element-flanked foreign DNA via homologous recombination. *J Biol Chem* 274:36585–36591.
- Karow JK, Constantinou A, Li JL, West SC, Hickson ID. 2000a. The Bloom's syndrome gene product promotes branch migration of holliday junctions. *Proc Natl Acad Sci USA* 97:6504–6508.
- Karow JK, Wu L, Hickson ID. 2000b. RecQ family helicases: roles in cancer and aging. *Curr Opin Genet Dev* 10:32–38.
- Kato S, Anderson RA, Camerini-Otero RD. 1986. Foreign DNA introduced by calcium phosphate is integrated into repetitive DNA elements of the mouse L cell genome. *Mol Cell Biol* 6:1787–1795.
- Kato S, Tachibana K, Takayama N, Kataoka H, Yoshida MC, Takano T. 1991. Genetic recombination in a chromosomal translocation t(2;8)(p11;q24) of a Burkitt's lymphoma cell line, KOBK101. *Gene* 97:239–244.
- Koduru P, Chaganti R. 1988. Congenital chromosome breakage clusters within Giemsa-light bands and identifies sites of chromatin instability. *Cytogenet Cell Genet* 49:269–274.
- Koduru PR, Goh JC, Pergolizzi RG, Lichtman SM, Broome JD. 1993. Molecular characterization of a variant Ph1 translocation t(9;22)(q34;q11;q13) in chronic myelogenous leukemia (CML) reveals the translocation of the 3'-part of BCR gene to the chromosome band 11q13. *Oncogene* 8:3239–3247.
- Kolomietz E, Al-Maghribi J, Brennan S, Karaskova J, Minkin S, Lipton J, Squire JA. 2001. Primary chromosomal rearrangements of leukemia are frequently accompanied by extensive submicroscopic deletions and may lead to altered prognosis. *Blood* 97:3581–3588.
- Korenberg JR, Rykowski MC. 1988. Human genome organization: Alu, lines, and the molecular structure of metaphase chromosome bands. *Cell* 53:391–400.
- Kornreich R, Bishop DF, Desnick RJ. 1990. Alpha-galactosidase A gene rearrangements causing Fabry disease. Identification of short direct repeats at breakpoints in an Alu-rich gene. *J Biol Chem* 265:9319–9326.
- Kowalczykowski SC. 2000. Initiation of genetic recombination and recombination-dependent replication. *Trends Biochem Sci* 25:156–165.
- Kozubek S, Lukasova E, Ryznar L, Kozubek M, Liskova A, Govorun RD, Krasavin EA, Horneck G. 1997. Distribution of ABL and BCR genes in cell nuclei of normal and irradiated lymphocytes. *Blood* 89:4537–4545.
- Kurahashi H, Shaikh TH, Emanuel BS. 2000. Alu-mediated PCR artifacts and the constitutional t(11;22) breakpoint. *Hum Mol Genet* 9:2727–2732.
- Labuda D, Zietkiewicz E, Mitchell GA. 1995. Alu elements as a source of genomic variation: deleterious effects and evolutionary novelties. In: Maraia RJ, editor. The impact of short interspersed elements (SINEs) on the host genome. R.G. Landes. 1–22.
- Le Beau MM, Rowley JD. 1984. Heritable fragile sites in cancer. *Nature* 308:607–608.
- Lehrman MA, Russell DW, Goldstein JL, Brown MS. 1986. Exon-Alu recombination deletes 5 kilobases from the low density lipoprotein receptor gene, producing a null phenotype in familial hypercholesterolemia. *Proc Natl Acad Sci USA* 83:3679–3683.
- Levrin O, Doggett NA, Auerbach AD. 1998. Identification of Alu-mediated deletions in the Fanconi anemia gene FAA. *Hum Mutat* 12:145–152.
- Lin CT, Lyu YL, Xiao H, Lin WH, Whang-Peng J. 2001. Suppression of gene amplification and chromosomal DNA integration by the DNA mismatch repair system. *Nucleic Acids Res* 29:3304–3310.
- Luo CM, Tang W, Mekeel KL, DeFrank JS, Anne PR, Powell SN. 1996. High frequency and error-prone DNA recombination in ataxia telangiectasia cell lines. *J Biol Chem* 271:4497–4503.
- Markert ML, Hutton JJ, Wiginton DA, States JC, Kaufman RE. 1988. Adenosine deaminase (ADA) deficiency due to deletion of the ADA gene promoter and first exon by homologous recombination between two Alu elements. *J Clin Invest* 81:1323–1327.
- Marshall B, Isidro G, Boavida MG. 1996. Insertion of a short Alu sequence into the hMSH2 gene following a double cross over next to sequences with chi homology [published erratum appears in *Gene* 197 Sep 15;197(1–2):413]. *Gene* 174:175–179.
- Martinelli G, Terragna C, Amabile M, Montefusco V, Testoni N, Ottaviani E, de Vivo A, Mianulli A, Saglio G, Tura S. 2000. Alu and translin recognition site sequences flanking translocation sites in a novel type of chimeric bcr-abl transcript suggest a possible general mechanism for bcr-abl breakpoints. *Haematologica* 85:40–46.
- Mauillon JL, Michel P, Limacher JM, Latouche JB, Dechelotte P, Charbonnier F, Martin C, Moreau V, Metayer J, Paillot B, Frebourg T. 1996. Identification of novel germline hMLH1 mutations including a 22 kb Alu-mediated deletion in patients with familial colorectal cancer. *Cancer Res* 56:5728–5733.
- Megonigal MD, Rappaport EF, Jones DH, Kim CS, Nowell PC, Lange BJ, Felix CA. 1997. Panhandle PCR strategy to amplify MLL genomic breakpoints in treatment-related leukemias. *Proc Natl Acad Sci USA* 94:11583–11588.
- Meyn MS. 1993. High spontaneous intrachromosomal recombination rates in ataxia-telangiectasia. *Science* 260:1327–1330.
- Michel B. 2000. Replication fork arrest and DNA recombination. *Trends Biochem Sci* 25:173–178.
- Michel B, Ehrlich SD, Uzzest M. 1997. DNA double-strand breaks caused by replication arrest. *EMBO J* 16:430–438.
- Miki Y, Katagiri T, Kasumi F, Yoshimoto T, Nakamura Y. 1996. Mutation analysis in the BRCA2 gene in primary breast cancers. *Nat Genet* 13:245–247.
- Montagna M, Santacatterina M, Torri A, Menin C, Zullato D, Chicco-Bianchi L, D'Andrea E. 1999. Identification of a 3 kb Alu-mediated BRCA1 gene rearrangement in two breast/ovarian cancer families. *Oncogene* 18:4160–4165.
- Morgan NV, Tipping AJ, Joenje H, Mathew CG. 1999. High frequency of large intragenic deletions in the Fanconi anemia group A gene. *Am J Hum Genet* 65:1330–1341.
- Morris C, Jeffs A, Smith T, McDonald M, Board P, Kennedy M, Fitzgerald P. 1996. BCR gene recombines with genomically distinct sites on band 11q13 in complex BCR-ABL translocations of chronic myeloid leukemia. *Oncogene* 12:677–685.
- Morrison C, Sonoda E, Takao N, Shinohara A, Yamamoto K, Takeda S. 2000. The controlling role of ATM in homologous recombination repair of DNA damage. *EMBO J* 19:463–471.
- Mycerowitz R, Hogikyan ND. 1987. A deletion involving Alu sequences in the beta-hexosaminidase alpha-chain gene of French Canadians with Tay-Sachs disease. *J Biol Chem* 262:15396–15399.
- Neote K, McInnes B, Mahuran DJ, Gravel RA. 1990. Structure and distribution of an Alu-type deletion mutation in Sandhoff disease. *J Clin Invest* 86:1524–1531.
- Neves H, Ramos C, da Silva MG, Parreira A, Parreira L. 1999. The nuclear topography of ABL, BCR, PML, and RARalpha genes: evidence for gene proximity in specific phases of the cell cycle and stages of hematopoietic differentiation. *Blood* 93:1197–1207.
- New JH, Sugiyama T, Zaitseva E, Kowalczykowski SC. 1998. Rad52 protein stimulates DNA strand exchange by Rad51 and replication protein A. *Nature* 391:407–410.
- Nicholls RD, Fischel-Ghodsian N, Higgs DR. 1987. Recombination

- at the human alpha-globin gene cluster: sequence features and topological constraints. *Cell* 49:369–378.
- Norris J, Fan D, Aleman C, Marks JR, Futreal PA, Wiseman RW, Iglehart JD, Deininger PL, McDonnell DP. 1995. Identification of a new subclass of Alu DNA repeats which can function as estrogen receptor-dependent transcriptional enhancers. *J Biol Chem* 270:22777–22782.
- Obata K, Hiraga H, Nojima T, Yoshida MC, Abe S. 1999. Molecular characterization of the genomic breakpoint junction in a t(11;22) translocation in Ewing sarcoma. *Genes Chromosomes Cancer* 25:6–15.
- Ohno H, Doi S, Yabumoto K, Fukuhara S, McKeithan TW. 1993. Molecular characterization of the t(14;19)(q32;q13) translocation in chronic lymphocytic leukemia. *Leukemia* 7:2057–2063.
- Osman F, Subramani S. 1998. Double-strand break-induced recombination in eukaryotes. *Prog Nucleic Acid Res Mol Biol* 58:263–299.
- Papadopoulos PC, Greenstein AM, Gaffney RA, Westbrook CA, Wiedemann LM. 1990. Characterization of the translocation breakpoint sequences in Philadelphia-positive acute lymphoblastic leukemia. *Genes Chromosomes Cancer* 1:233–239.
- Pastink A, Eeken JC, Lohman PH. 2001. Genomic integrity and the repair of double-strand DNA breaks. *Mutat Res* 480/481:37–50.
- Paull TT, Gellert M. 1999. Nbs1 potentiates ATP-driven DNA unwinding and endonuclease cleavage by the Mre11/Rad50 complex. *Genes Dev* 13:1276–1288.
- Petukhova G, Stratton S, Sung P. 1998. Catalysis of homologous DNA pairing by yeast Rad51 and Rad54 proteins. *Nature* 393:91–94.
- Pousi B, Hautala T, Heikkinen J, Pajunen L, Kivirikko KI, Myllylä R. 1994. Alu-Alu recombination results in a duplication of seven exons in the lysyl hydroxylase gene in a patient with the type VI variant of Ehlers–Danlos syndrome. *Am J Hum Genet* 55:899–906.
- Prolla TA. 1998. DNA mismatch repair and cancer. *Curr Opin Cell Biol* 10:311–316.
- Ramsden DA, Gellert M. 1998. Ku protein stimulates DNA end joining by mammalian DNA ligases: a direct role for Ku in repair of DNA double-strand breaks. *EMBO J* 17:609–614.
- Rayssiguier C, Thaler DS, Radman M. 1989. The barrier to recombination between *Escherichia coli* and *Salmonella typhimurium* is disrupted in mismatch-repair mutants. *Nature* 342:396–401.
- Richardson C, Jasin M. 2000. Coupled homologous and nonhomologous repair of a double-strand break preserves genomic integrity in mammalian cells. *Mol Cell Biol* 20:9068–9075.
- Richardson C, Moynahan ME, Jasin M. 1998. Double-strand break repair by interchromosomal recombination: suppression of chromosomal translocations. *Genes Dev* 12:3831–3842.
- Rogers J. 1985. Oncogene chromosome breakpoints and alu sequences. *Nature* 317:559.
- Rohlf EM, Puget N, Graham ML, Weber BL, Garber JE, Skrzynia C, Halperin JL, Lenoir GM, Silverman LM, Mazoyer S. 2000. An Alu-mediated 7.1 kb deletion of BRCA1 exons 8 and 9 in breast and ovarian cancer families that results in alternative splicing of exon 10. *Genes Chromosomes Cancer* 28:300–307.
- Rothberg PG, Ponnuru S, Baker D, Bradley JF, Freeman AI, Cibis GW, Harris DJ, Heruth DP. 1997. A deletion polymorphism due to Alu-Alu recombination in intron 2 of the retinoblastoma gene: association with human gliomas. *Mol Carcinog* 19:69–73.
- Rouet P, Smith F, Jasin M. 1994. Expression of a site-specific endonuclease stimulates homologous recombination in mammalian cells. *Proc Natl Acad Sci USA* 91:6064–6068.
- Rowe SM, Coughlan SJ, McKenna NJ, Garrett E, Kieback DG, Carney DN, Headon DR. 1995. Ovarian carcinoma-associated TaqI restriction fragment length polymorphism in intron G of the progesterone receptor gene is due to an Alu sequence insertion. *Cancer Res* 55:2743–2745.
- Rudiger NS, Gregersen N, Kielland-Brandt MC. 1995. One short well conserved region of Alu-sequences is involved in human gene rearrangements and has homology with prokaryotic chi. *Nucleic Acids Res* 23:256–260.
- Salganik RI, Dianov GL. 1992. Molecular mechanisms of the formation of DNA double-strand breaks and induction of genomic rearrangements. *Mutat Res* 266:163–170.
- Schmid CW. 1996. Alu: structure, origin, evolution, significance and function of one-tenth of human DNA. *Prog Nucleic Acid Res Mol Biol* 53:283–319.
- Schmid CW. 1998. Does SINE evolution preclude Alu function. *Nucleic Acids Res* 26:4541–4550.
- Schnittger S, Wormann B, Hiddemann W, Griesinger F. 1998. Partial tandem duplications of the MLL gene are detectable in peripheral blood and bone marrow of nearly all healthy donors. *Blood* 92:1728–1734.
- Scully R, Chen J, Plug A, Xiao Y, Weaver D, Feunteun J, Ashley T, Livingston DM. 1997. Association of BRCA1 with Rad51 in mitotic and meiotic cells. *Cell* 88:265–275.
- Shinohara A, Ogawa T. 1995. Homologous recombination and the roles of double-strand breaks. *Trends Biochem Sci* 20:387–391.
- Shinohara A, Ogawa T. 1999. Rad51/RecA protein families and the associated proteins in eukaryotes. *Mutat Res* 435:13–21.
- Shinohara A, Ogawa H, Ogawa T. 1992. Rad51 protein involved in repair and recombination in *S. cerevisiae* is a RecA-like protein. *Cell* 69:457–470.
- Shinohara A, Ogawa H, Matsuda Y, Ushio N, Ikeo K, Ogawa T. 1993. Cloning of human, mouse and fission yeast recombination genes homologous to RAD51 and recA. *Nat Genet* 4:239–243.
- Shulman MJ, Collins C, Connor A, Read LR, Baker MD. 1995. Interchromosomal recombination is suppressed in mammalian somatic cells. *EMBO J* 14:4102–4107.
- Sinclair PB, Nacheva EP, Leversha M, Telford N, Chang J, Reid A, Bench A, Champion K, Huntly B, Green AR. 2000. Large deletions at the t(9;22) breakpoint are common and may identify a poor-prognosis subgroup of patients with chronic myeloid leukemia. *Blood* 95:738–743.
- Smidt M, Kirsch I, Ratner L. 1990. Deletion of Alu sequences in the fifth c-sis intron in individuals with meningiomas. *J Clin Invest* 86:1151–1157.
- Smit AF. 1996. The origin of interspersed repeats in the human genome. *Curr Opin Genet Dev* 6:743–748.
- Smit AF. 1999. Interspersed repeats and other mementos of transposable elements in mammalian genomes. *Curr Opin Genet Dev* 9:657–663.
- Smith GC, Jackson SP. 1999. The DNA-dependent protein kinase. *Genes Dev* 13:916–934.
- Sonoda E, Sasaki MS, Buerstedde JM, Bezzubova O, Shinohara A, Ogawa H, Takata M, Yamaguchi-Iwai Y, Takeda S. 1998. Rad51-deficient vertebrate cells accumulate chromosomal breaks prior to cell death. *EMBO J* 17:598–608.
- Stoppa-Lyonnet D, Carter PE, Meo T, Tosi M. 1990. Clusters of intragenic Alu repeats predispose the human C1 inhibitor locus to deleterious rearrangements. *Proc Natl Acad Sci USA* 87:1551–1555.
- Strissel PL, Strick R, Rowley JD, Zeleznik-Le NJ. 1998. An in vivo topoisomerase II cleavage site and a DNase I hypersensitive site colocalize near exon 9 in the MLL breakpoint cluster region. *Blood* 92:3793–3803.
- Strout MP, Marcucci G, Bloomfield CD, Caligiuri MA. 1998. The partial tandem duplication of ALL1 (MLL) is consistently generated by Alu-mediated homologous recombination in acute myeloid leukemia. *Proc Natl Acad Sci USA* 95:2390–2395.
- Sturzbecher HW, Donzelmann B, Henning W, Knippschild U, Buchhop S. 1996. p53 is linked directly to homologous recombination processes via RAD51/RecA protein interaction. *EMBO J* 15:1992–2002.
- Sugawara N, Paques F, Colaiacovo M, Haber JE. 1997. Role of *Saccharomyces cerevisiae* Msh2 and Msh3 repair proteins in double-strand break-induced recombination. *Proc Natl Acad Sci USA* 94:9214–9219.
- Sung P. 1997. Function of yeast Rad52 protein as a mediator between replication protein A and the Rad51 recombinase. *J Biol Chem* 272:28194–28197.
- Super HG, Strissel PL, Sobulo OM, Burian D, Reshmi SC, Roe B, Zeleznik-Le NJ, Diaz MO, Rowley JD. 1997. Identification of complex genomic breakpoint junctions in the t(9;11) MLL-AF9 fusion gene in acute leukemia. *Genes Chromosomes Cancer* 20:185–195.
- Swagemakers SM, Essers J, de Wit J, Hoeijmakers JH, Kanaar R. 1998. The human RAD54 recombinational DNA repair protein is a double-stranded DNA-dependent ATPase. *J Biol Chem* 273:28292–28297.
- Swensen J, Hoffman M, Skolnick MH, Neuhausen SL. 1997. Identification of a 14 kb deletion involving the promoter region of BRCA1 in a breast cancer family. *Hum Mol Genet* 6:1513–1517.
- Szostak JW, Orr-Weaver TL, Rothstein RJ, Stahl FW. 1983. The double-strand-break repair model for recombination. *Cell* 33:25–35.
- Takata M, Sasaki MS, Sonoda E, Morrison C, Hashimoto M, Utsumi H, Yamaguchi-Iwai Y, Shinohara A, Takeda S. 1998. Homologous recombination and non-homologous end-joining pathways of DNA double-strand break repair have overlapping roles in the

- maintenance of chromosomal integrity in vertebrate cells. *EMBO J* 17:5497–5508.
- Toth G, Jurka J. 1994. Repetitive DNA in and around translocation breakpoints of the Philadelphia chromosome. *Gene* 140:285–288.
- Tracy RB, Kowalczykowski SC. 1996. In vitro selection of preferred DNA pairing sequences by the *Escherichia coli* RecA protein. *Genes Dev* 10:1890–1903.
- Trujillo KM, Yuan SS, Lee EY, Sung P. 1998. Nuclease activities in a complex of human recombination and DNA repair factors Rad50, Mre11, and p95. *J Biol Chem* 273:21447–21450.
- Tvrđik T, Marcus S, Hou SM, Falt S, Noori P, Podlutska N, Hanefeld F, Stromme P, Lambert B. 1998. Molecular characterization of two deletion events involving Alu-sequences, one novel base substitution and two tentative hotspot mutations in the hypoxanthine phosphoribosyltransferase (HPRT) gene in five patients with Lesch–Nyhan syndrome. *Hum Genet* 103:311–318.
- Tycko B, Sklar J. 1990. Chromosomal translocations in lymphoid neoplasia: a reappraisal of the recombinase model. *Cancer Cells* 2:1–8.
- van der Feltz MJ, Shivji MK, Allen PB, Heisterkamp N, Groffen J, Wiedemann LM. 1989. Nucleotide sequence of both reciprocal translocation junction regions in a patient with Ph positive acute lymphoblastic leukaemia, with a breakpoint within the first intron of the BCR gene. *Nucleic Acids Res* 17:1–10.
- Vansant G, Reynolds WF. 1995. The consensus sequence of a major Alu subfamily contains a functional retinoic acid response element. *Proc Natl Acad Sci USA* 92:8229–8233.
- von Hippel PH, Berg OG. 1989. Facilitated target location in biological systems. *J Biol Chem* 264:675–678.
- Waldman AS, Liskay RM. 1988. Dependence of intrachromosomal recombination in mammalian cells on uninterrupted homology. *Mol Cell Biol* 8:5350–5357.
- Wallenburg JC, Nepveu A, Chartrand P. 1987. Integration of a vector containing rodent repetitive elements in the rat genome. *Nucleic Acids Res* 15:7849–7863.
- Wenger SL. 1992. Relationship between fragile sites and cancer breakpoints [letter]. *Cancer Genet Cytogenet* 61:213.
- Wiedemann LM, MacGregor A, Caldas C. 1999. Analysis of the region of the 5' end of the MLL gene involved in genomic duplication events. *Br J Haematol* 105:256–264.
- Wong AK, Pero R, Ormonde PA, Tavtigian SV, Bartel PL. 1997. RAD51 interacts with the evolutionarily conserved BRC motifs in the human breast cancer susceptibility gene *brca2*. *J Biol Chem* 272:31941–31944.
- Yu CE, Oshima J, Fu YH, Wijsman EM, Hisama F, Alisch R, Matthews S, Nakura J, Miki T, Ouais S, Martin GM, Mulligan J, Schellenberg GD. 1996. Positional cloning of the Werner's syndrome gene. *Science* 272:258–262.
- Yuan ZM, Huang Y, Ishiko T, Nakada S, Utsugisawa T, Kharbanda S, Wang R, Sung P, Shinohara A, Weichselbaum R, Kufe D. 1998. Regulation of Rad51 function by c-Abl in response to DNA damage. *J Biol Chem* 273:3799–3802.
- Yunis JJ, Soreng AL. 1984. Constitutive fragile sites and cancer. *Science* 226:1199–1204.
- Yunis JJ, Soreng AL, Bowe AE. 1987. Fragile sites are targets of diverse mutagens and carcinogens. *Oncogene* 1:59–69.
- Zhang JG, Goldman JM, Cross NC. 1995. Characterization of genomic BCR-ABL breakpoints in chronic myeloid leukaemia by PCR. *Br J Haematol* 90:138–146.
- Zhang ZP, Blomback M, Egberg N, Falk G, Anvret M. 1994. Characterization of the von Willebrand factor gene (VWF) in von Willebrand disease type III patients from 24 families of Swedish and Finnish origin. *Genomics* 21:188–193.
- Zucman-Rossi J, Legoix P, Victor JM, Lopez B, Thomas G. 1998. Chromosome translocation based on illegitimate recombination in human tumors. *Proc Natl Acad Sci USA* 95:11786–11791.