

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/243859996>

# ChemInform Abstract: Copper-Catalyzed Three-Component Synthesis of 2-Iminodihydrocoumarins and 2-Iminocoumarins.

ARTICLE in CHEMINFORM · MAY 2010

Impact Factor: 0.74 · DOI: 10.1002/adsc.200900890

---

CITATIONS

33

---

READS

21

6 AUTHORS, INCLUDING:



Jing Wang

University of Southampton

910 PUBLICATIONS 9,705 CITATIONS

SEE PROFILE



Ping Lu

Zhejiang University

334 PUBLICATIONS 6,884 CITATIONS

SEE PROFILE



Yanguang Wang

Zhejiang University

154 PUBLICATIONS 2,762 CITATIONS

SEE PROFILE

# Copper-Catalyzed Three-Component Synthesis of 2-Iminodihydrocoumarins and 2-Iminocoumarins

Yang Shen,<sup>a</sup> Sunliang Cui,<sup>a</sup> Jing Wang,<sup>a</sup> Xiaopeng Chen,<sup>a</sup> Ping Lu,<sup>a,\*</sup> and Yanguang Wang<sup>a,b,\*</sup>

<sup>a</sup> Department of Chemistry, Zhejiang University, Hangzhou 310027, People's Republic of China  
Fax: (+86)-571-8795-1512; e-mail: pinglu@zju.edu.cn or orgwyg@zju.edu.cn

<sup>b</sup> State Key Laboratory of Applied Organic Chemistry, Lanzhou University, Lanzhou 730000, People's Republic of China

Received: December 22, 2009; Revised: March 30, 2010; Published online: April 23, 2010

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/adsc.200900890>.

**Abstract:** An efficient synthesis of *N*-sulfonyl-substituted 2-imino-3,4-dihydrocoumarins and 2-iminocoumarins *via* a copper-catalyzed multicomponent reaction of sulfonyl azides with terminal alkynes and  $\beta$ -(*ortho*-hydroxyphenyl)- $\alpha,\beta$ -unsaturated ketones or *ortho*-hydroxyphenylpropiolates has been developed. The cascade process involves trapping the ketimine by a nucleophilic addition and an intramolecular Michael addition. This methodology could well be extended to the concise synthesis of the polysubstituted piperidine scaffold.

**Keywords:** cascade reactions; copper; coumarins; ketenimines; multicomponent reactions


addition (CuAAC) and the generation of ketenimine intermediate by nucleophiles. Instead of sulfonyl azides, phosphoryl azides could also participate in the reaction and afford three component adducts.<sup>[6]</sup> As an extension of our previous work, here we use  $\beta$ -(*ortho*-hydroxyphenyl)- $\alpha,\beta$ -unsaturated ketones as substrates, with both nucleophilicity and electrophilicity, for trapping the ketenimine intermediates. Fortunately, 2-iminocoumarins were ideally obtained in moderate yields and with *trans* as the predominant configuration.

The CuI-catalyzed reaction of toluenesulfonyl azide (**1a**) with phenylacetylene (**2a**) and chalcone (**3a**) was used to screen the reaction conditions (Table 1). Triethylamine (TEA, 1 equiv.) was proven to be the

Iminocoumarins are one type of protein tyrosine kinase (PTK) inhibitors that are most valuable for the treatment of diseases involving excess cell proliferation as well as the antitumor process.<sup>[1]</sup> They are also useful fluorescent dyes in fields as various as cell biology, medical analysis and sensors.<sup>[2]</sup> Classic methods for synthesis of iminocoumarins, such as Knoevenagel condensation and modification from coumarins, suffer from several shortcomings including limited substituents and troublesome chemical process management.<sup>[3]</sup> In order to overcome these shortages, here we would like to report a concise synthesis of iminocoumarins *via* multicomponent reactions that are versatile with regard to the substrate scope and the synthetic applicability.

Recently, we<sup>[4]</sup> and others<sup>[5]</sup> have developed Cu-catalyzed three-component reactions of sulfonyl azides with terminal alkynes and a number of nucleophiles, such as amines, alcohols as well as water. These reactions involve a copper-catalyzed azide-alkyne cyclo-


**Table 1.** Optimization of reaction conditions for the synthesis of iminodihydrocoumarin **4a**.<sup>[a]</sup>

					
Entry	Base	Solvent	Time [h]	Yield <sup>[b]</sup> [%]	<i>dr</i> <sup>[c]</sup>
1	Et <sub>3</sub> N	CH <sub>2</sub> Cl <sub>2</sub>	8	86	95:5
2	Et <sub>3</sub> N	CH <sub>3</sub> CN	8	72	95:5
3	pyridine	CH <sub>2</sub> Cl <sub>2</sub>	8	83	92:8
4	pyridine	CH <sub>3</sub> CN	12	55	90:10
5	2,6-lutidine	CHCl <sub>3</sub>	8	75	95:5
6	K <sub>2</sub> CO <sub>3</sub>	CH <sub>2</sub> Cl <sub>2</sub>	12	47	90:10

<sup>[a]</sup> Reaction conditions: **1a** (0.6 mmol), **2a** (0.6 mmol), **3a** (0.5 mmol), CuI (0.06 mmol), the indicated base (0.6 mmol) and solvent (5 mL).

<sup>[b]</sup> Yield of the isolated product.

<sup>[c]</sup> The ratio was determined by <sup>1</sup>H NMR spectroscopy of the crude product.

**Table 2.** Three-component reaction of azides **1** with alkynes **2** and chalcones **3**.<sup>[a]</sup>

Entry	<b>1</b> ( $R^1$ )	<b>2</b> ( $R^2$ )	<b>3</b> ( $R^3/R^4$ )	Product <b>4</b> /Yield [%] <sup>[b]</sup>
1	<b>1a</b> (4-MeC <sub>6</sub> H <sub>4</sub> )	<b>2a</b> (Ph)	<b>3a</b> (H/Ph)	<b>4a</b> /86 <sup>[c]</sup>
2	<b>1b</b> (Ph)	<b>2b</b> (4-MeOC <sub>6</sub> H <sub>4</sub> )	<b>3a</b>	<b>4b</b> /73 <sup>[c]</sup>
3	<b>1c</b> (4-ClC <sub>6</sub> H <sub>4</sub> )	<b>2c</b> (4-MeC <sub>6</sub> H <sub>4</sub> )	<b>3a</b>	<b>4c</b> /81 <sup>[c]</sup>
4	<b>1d</b> (5-Me <sub>2</sub> N-naphth-1-yl)	<b>2d</b> (4-FC <sub>6</sub> H <sub>4</sub> )	<b>3a</b>	<b>4d</b> /75 <sup>[c]</sup>
5	<b>1a</b>	<b>2e</b> ( <i>n</i> -Bu)	<b>3a</b>	<b>4e</b> /97 <sup>[d]</sup>
6	<b>1a</b>	<b>2f</b> (1-cyclohexenyl)	<b>3a</b>	<b>4f</b> /88 <sup>[c]</sup>
7	<b>1e</b> (Me)	<b>2a</b>	<b>3b</b> (H/4-ClC <sub>6</sub> H <sub>4</sub> )	<b>4g</b> /83 <sup>[c]</sup>
8	<b>1b</b>	<b>2f</b>	<b>3b</b>	<b>4h</b> /85 <sup>[c]</sup>
9	<b>1c</b>	<b>2a</b>	<b>3c</b> (H/2-MeC <sub>6</sub> H <sub>4</sub> )	<b>4i</b> /71 <sup>[c]</sup>
10	<b>1b</b>	<b>2c</b>	<b>3c</b>	<b>4j</b> /79 <sup>[c]</sup>
11	<b>1a</b>	<b>2b</b>	<b>3d</b> (4-Br/Ph)	<b>4k</b> /69 <sup>[c]</sup>
12	<b>1c</b>	<b>2e</b>	<b>3d</b>	<b>4l</b> /74 <sup>[c]</sup>
13	<b>1a</b>	<b>2g</b> (4-EtC <sub>6</sub> H <sub>4</sub> )	<b>3d</b>	<b>4m</b> /81 <sup>[c]</sup>
14	<b>1a</b>	<b>2a</b>	<b>3e</b> (3-MeO/Ph)	<b>4n</b> /80 <sup>[c]</sup>
15	<b>1a</b>	<b>2f</b>	<b>3e</b>	<b>4o</b> /85 <sup>[c]</sup>
16	<b>1a</b>	<b>2a</b>	<b>3f</b> (H/Me)	<b>4p</b> /46 <sup>[c]</sup>

<sup>[a]</sup> Reaction conditions: **1** (0.6 mmol), **2** (0.6 mmol), **3** (0.5 mmol), CuI (0.06 mmol), TEA (0.6 mmol), CH<sub>2</sub>Cl<sub>2</sub> (5 mL), room temperature, 8 h.

<sup>[b]</sup> Isolated yield.

<sup>[c]</sup> *dr* > 95:5 based on <sup>1</sup>H NMR spectroscopy of the crude product.

<sup>[d]</sup> *dr* ≈ 88:12.

<sup>[e]</sup> *dr* ≈ 82:18.


most effective base among those others tested, such as pyridine, 2,6-lutidine and potassium carbonate (Table 1, entries 1, 3–6). As to the solvent, in comparison with acetonitrile, dichloromethane not only provided a higher isolated yield but also a higher diastereomeric ratio (*dr* = 95:5) in favour of the *trans* isomer (Table 1, entries 1 and 2).


With the optimized reaction conditions in hand, we tested the substrate diversity with a variety of azides, alkynes and chalcones (Table 2). In all cases, the reactions afforded the corresponding *N*-toluenesulfonyl-aminodihydrocoumarins **4a–4p** in 46–97% yields. The stereoselectivity of the reaction was related to the character of alkynes. Arylacetylenes **2a–2d** and **2g** (Table 2, entries 1–4, 7, 9–11, 13, 14, and 16) and cyclohexenylacetylene (**2f**) (Table 2, entries 6, 8, and 15) underwent this cascade reaction smoothly with excellent diastereomeric ratios (*dr* > 95:5) while 1-hexyne (**2e**) gave a slightly decreased diastereomeric ratio (Table 2, entries 5 and 12).

The structures of the products was fully characterized by <sup>1</sup>H NMR and <sup>13</sup>C NMR spectroscopy as well as high-resolution mass spectrometry (see Supporting Information). The structure of **4h** was further estab-

lished by a single crystal X-ray analysis (Figure 1),<sup>[7]</sup> which clearly presented the *trans* configuration.

Based on our previous works, a possible mechanism was proposed and is shown in Scheme 1. Firstly, ketenimine **A** could be formed by the Cu-catalyzed cycloaddition reaction of azides **1a** and alkynes **2a** via a click reaction and a subsequent release of N<sub>2</sub>.<sup>[4,5]</sup> Then nucleophile **B**, deprotonated by triethylamine, could then attack the central carbon of **A** to form **C**.

**Figure 1.** X-ray crystal structure of **4h**.




**Scheme 1.** Possible mechanism for the formation of **4a**.


Finally, **C** could undergo an intramolecular Michael addition and a sequential proton transfer to afford the iminodihydrocoumarin **4a**.

As shown in Figure 1, the iminodihydrocoumarin ring exists in a twisted planar form. The vicinal coupling constant ( $^3J$ ) for two protons on C-1 and C-2 is not clearly shown in the  $^1\text{H}$  NMR spectrum because the dihedral angle of H-C-1-C-2-H is  $92.77^\circ$ . Based on the Karplus relation,  $^3J$  is calculated to be 2.07 Hz. Meanwhile, the structures of both *trans*-**4h** and *cis*-**4h** have been optimized by the RHF/6-31G(d) method. The dihedral angles of H-C-1-C-2-H in *trans*-**4h** and *cis*-**4h** are calculated to be  $96^\circ$  and  $48^\circ$ , respectively. Hence, the  $^3J$  values are calculated to be 2.21 Hz for *trans*-**4h** and 8.2 Hz for *cis*-**4h**. Based on these calculated results and  $^1\text{H}$  NMR data, we assigned all compounds **4** as having the *trans* configuration. The observed diastereoselectivity could be rationalized by two factors. One is the steric hindrance (Scheme 2) and the second is the result of the thermodynamic equilibration. By calculation of the energy potentials of *trans*-**4h** and *cis*-**4h** [RHF/6-31G(d) method], *trans*-**4h** is more stable than *cis*-**4h** and the energy difference is  $3.41\text{ kcal}\cdot\text{mol}^{-1}$ . In comparison to aromatic acetylenes (**2a–2d**) and cyclohexenylacetylene (**2g**), 1-hexyne (**2f**) gave a relatively lower diastereoselectivity due to the fact that the steric hindrance has been released a little when the *cis* configuration is adopted in the step of the Michael addition.

In order to extend the substrate diversity, *ortho*-hydroxyphenyl- $\alpha,\beta$ -unsaturated ester **5** was tested for this cascade reaction. Unfortunately, although trapping the ketenimine by phenolic anion was observed in this case, we did not detect the sequential intramolecular Michael addition product. The three-compo-



**Scheme 2.** Depiction of the *trans* stereocontrol of **4a**.




**Scheme 3.** Three-component reaction of **1a**, **2a** and **5**.

nent adduct product **6** was obtained in 90% isolated yield (Scheme 3).

Rather than altering the ketone to an ester, we tried to change C=C into C≡C. However, many attempts at synthesis of 3-(2-hydroxyphenyl)-1-phenylprop-2-yn-1-one failed because this expected compound was too unstable and would intramolecularly cyclize to benzofuran-2-yl(phenyl)methanone in high yield.


Based on the above survey, we used *ortho*-hydroxyphenylpropiolates **7**, which couple an ester group and a C≡C bond, to test the reality of this cascade reaction. Fortunately, the anticipated iminocoumarins **8** were obtained in high yields (Table 3). Both aromatic sulfonyl azides (Table 3, entries 1–8 and 10–17) and aliphatic sulfonyl azide (Table 3, entry 9) gave iminocoumarins in good to excellent yields. An electronic effect, either by a conjugative effect or by an inductive effect, on the aromatic sulfonyl azides **1**, aromatic acetylenes **2** and phenylpropiolates **7** did not affect the reaction. Hex-1-yne (**2e**) and 3,3-dimethylbut-1-yne (**2j**) afforded **9a** and **9b** in 87% and 95% yields, respectively (Scheme 4). In these cases, the basicity of intermediate **E** was increased. Therefore it preferred abstraction of a proton instead of a sequential Michael addition.

**Table 3.** Three-component reaction of azides **1** with alkynes **2** and phenylpropiolates **7**.<sup>[a]</sup>


Entry	<b>1</b> (R <sup>1</sup> )	<b>2</b> (R <sup>2</sup> )	<b>7</b> (R <sup>3</sup> /R <sup>4</sup> )	Product <b>8</b> /Yield <sup>[b]</sup> [%]
1	<b>1a</b> (4-MeC <sub>6</sub> H <sub>4</sub> )	<b>2a</b> (Ph)	<b>7a</b> (H/Me)	<b>8a</b> /92
2	<b>1b</b> (Ph)	<b>2a</b>	<b>7a</b>	<b>8b</b> /91
3	<b>1a</b>	<b>2b</b> (4-MeOC <sub>6</sub> H <sub>4</sub> )	<b>7a</b>	<b>8c</b> /90
4	<b>1a</b>	<b>2c</b> (4-MeC <sub>6</sub> H <sub>4</sub> )	<b>7a</b>	<b>8d</b> /96
5	<b>1a</b>	<b>2g</b> (4-EtC <sub>6</sub> H <sub>4</sub> )	<b>7a</b>	<b>8e</b> /98
6	<b>1a</b>	<b>2a</b>	<b>7a</b>	<b>8f</b> /95
7	<b>1f</b> (4-O <sub>2</sub> NC <sub>6</sub> H <sub>4</sub> )	<b>2a</b>	<b>7a</b>	<b>8g</b> /92
8	<b>1g</b> (4-AcNHC <sub>6</sub> H <sub>4</sub> )	<b>2a</b>	<b>7a</b>	<b>8h</b> /93
9	<b>1e</b> (Me)	<b>2a</b>	<b>7a</b>	<b>8i</b> /87
10	<b>1c</b> (4-ClC <sub>6</sub> H <sub>4</sub> )	<b>2c</b>	<b>7a</b>	<b>8j</b> /92
11	<b>1f</b>	<b>2b</b>	<b>7a</b>	<b>8k</b> /81
12	<b>1a</b>	<b>2h</b> (4- <i>t</i> -BuC <sub>6</sub> H <sub>4</sub> )	<b>7a</b>	<b>8l</b> /96
13	<b>1a</b>	<b>2i</b> (3-ClC <sub>6</sub> H <sub>4</sub> )	<b>7a</b>	<b>8m</b> /90
14	<b>1a</b>	<b>2a</b>	<b>7b</b> (H/Et)	<b>8n</b> /94
15	<b>1a</b>	<b>2a</b>	<b>7c</b> (H/Bn)	<b>8o</b> /93
16	<b>1a</b>	<b>2a</b>	<b>7d</b> (4-Cl/Me)	<b>8p</b> /91
17	<b>1a</b>	<b>2a</b>	<b>7e</b> (4-Me/Me)	<b>8q</b> /93

<sup>[a]</sup> Reaction conditions: **1** (1.2 mmol), **2** (1.2 mmol), **7** (1.0 mmol), CuI (0.1 mmol), CH<sub>2</sub>Cl<sub>2</sub> (10 mL), TEA (1.2 mmol), room temperature, 24 h.


<sup>[b]</sup> Isolated yield.

**Scheme 4.** Three-component reaction of propiolate **7a** with azide **1a** and alkynes **2**.

To highlight the utility of this methodology, typically with high diastereoselectivity in the case of (*ortho*-hydroxyphenyl)- $\alpha,\beta$ -unsaturated ketones, we synthesized **4a** on a gram scale and converted it into synthetically and biologically useful molecules such as the substituted piperidine **12** (Scheme 5).<sup>[8]</sup> In this procedure, **4a** was firstly reduced by NaBH<sub>4</sub> to afford **10**. Oxidation of secondary alcohol **10** to the ketone, followed by protection of the phenolic OH and then reduction by NaBH<sub>4</sub> furnished **11** smoothly. Finally, *via* an intramolecular Mitsunobu reaction, **12** was obtained. Selective protection of the phenolic OH of **10** failed. As a net result, starting from chalcone (**3a**), pi-

peridine **12** could be prepared in six steps and with a total yield of 43%.

In conclusion, we have developed an efficient and highly diastereoselective synthesis of *N*-sulfonyl-substituted 2-iminodihydrocoumarins and 2-iminocoumarins *via* a copper-catalyzed three-component reaction of sulfonyl azides with terminal alkynes and  $\beta$ -(*ortho*-hydroxyphenyl)- $\alpha,\beta$ -unsaturated ketones or *ortho*-hydroxyphenylpropiolates. The *trans*-iminodihydrocoumarins could be applied for the stereoselective synthesis of chiral piperidine scaffolds. Investigations on more synthetic applications of the present approach are underway in our laboratory.

Scheme 5. Synthesis of piperidine **12** from **4a**.

## Experimental Section

### General Procedure for the Synthesis of Iminodihydrocoumarins **4**

Triethylamine (0.6 mmol) was slowly added into a solution of azide **1** (0.6 mmol), alkyne **2** (0.6 mmol), chalcone **3** (0.5 mmol) and CuI (0.06 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) over 3–5 min under a nitrogen atmosphere at room temperature. After the reaction mixture had been stirred at the same temperature for 8 h, the solution was diluted with CH<sub>2</sub>Cl<sub>2</sub> (20 mL). The organic phase was then washed with water (10 mL) and brine (10 mL), and dried over sodium sulfate. After filtration, the solvent was removed under vacuum and the residue was subject to flash chromatography on silica gel (petroleum ether/ethyl acetate, 3:1 to 1:2).

### General Procedure for the Synthesis of Iminocoumarins **8**

Triethylamine (1.2 mmol) was slowly added into a solution of azide **1** (1.2 mmol), alkyne **2** (1.2 mmol), phenylpropiolate **7** (1.0 mmol) and CuI (0.1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) over 3–5 min under a nitrogen atmosphere at room temperature. After the reaction mixture had been stirred at the same temperature for 24 h, the solution was diluted with CH<sub>2</sub>Cl<sub>2</sub> (20 mL). The organic phase was washed with water (10 mL) and brine (10 mL), and dried over sodium sulfate. After filtration, the solvent was removed under vacuum and the residue was subject to flash chromatography on silica gel (petroleum ether/ethyl acetate, 3:1 to 1:2) to afford pure **8**.

**Data for compound 4a:** white solid; mp 173–175 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz): δ = 7.90 (m, 4H), 7.59 (m, 1H), 7.46 (m, 2H), 7.31–7.15 (m, 7H), 7.07 (m, 4H), 4.19 (s, 1H), 4.03 (m, 1H), 3.34 (dd, *J* = 8.0, 18.0 Hz, 1H), 3.17 (dd, *J* = 5.2, 18.4 Hz, 1H), 2.42 (s, 3H); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz): δ = 196.6, 165.4, 149.9, 143.7, 138.2, 136.2, 136.1,

133.7, 129.3, 129.0, 128.8, 128.7, 128.0, 127.61, 127.57, 126.9, 126.0, 123.7, 116.6, 48.5, 43.4, 37.1, 21.6; IR (KBr): ν = 2950, 1686, 1639, 1598, 1332, 1157, 769 cm<sup>−1</sup>; MS (ESI): *m/z* = 494.3 ([M−H]<sup>−</sup>); HR-MS (ESI): *m/z* = 518.1385, calcd for C<sub>30</sub>H<sub>25</sub>NO<sub>5</sub>S + Na<sup>+</sup> ([M+Na]<sup>+</sup>): 518.1397.

**Data for compound 8a:** white solid; mp 205–206 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 400 MHz): δ = 7.81 (d, *J* = 8.4 Hz, 2H), 7.60–7.55 (m, 2H), 7.50–7.48 (m, 2H), 7.44–7.39 (m, 2H), 7.39–7.35 (t, *J* = 8.0 Hz, 1H), 7.28–7.26 (m, 2H), 7.23 (d, *J* = 8.4 Hz, 2H), 3.68 (s, 5H), 2.38 (s, 3H); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 100 MHz): δ = 169.4, 158.1, 151.7, 142.8, 142.7, 139.2, 133.2, 131.9, 130.8, 129.5, 129.0, 128.6, 128.5, 126.9, 125.7, 124.8, 119.7, 117.0, 52.7, 35.8, 21.4; IR (KBr): ν = 1742, 1620, 1544, 1450, 1318, 1088, 1157, 777 cm<sup>−1</sup>; MS (ESI): *m/z* = 470.0 ([M+Na]<sup>+</sup>); HR-MS (ESI): *m/z* = 470.1022, calcd for C<sub>25</sub>H<sub>21</sub>NO<sub>5</sub>S + Na<sup>+</sup> ([M+Na]<sup>+</sup>): 470.1033.

## Acknowledgements

The authors gratefully acknowledge the financial support of the National Natural Science Foundation of China (No. 20872128; J0830413).

## References

- [1] a) T. R. Burke, B. Lim, V. E. Marquez, Z.-H. Li, J. B. Bolen, I. Stefanova, I. D. Horak, *J. Med. Chem.* **1993**, *36*, 425–432; b) C. N. O'Callaghan, M. L. Conalty, *Proc. Royal Irish Acad. B* **1979**, *6*, 87–98.
- [2] a) K. Komatsu, Y. Urano, H. Kojima, T. Nagano, *J. Am. Chem. Soc.* **2007**, *129*, 13447–13454; b) F. Liepouri, T. G. Deligeorgiev, Z. Veneti, C. Savakis, H. E. Katerinopoulos, *Cell Calcium* **2002**, *31*, 221–227; c) F. Liepouri, E. Foukaraki, T. G. Deligeorgiev, H. E. Katerinopoulos, *Cell Calcium* **2001**, *30*, 331–335; d) H. Turki, S. Abid, S. Fery-Forgues, R. E. Gharbi, *Dyes Pigm.* **2007**, *73*, 311–316.
- [3] a) J. Volmajer, R. Toplak, S. Bittner, I. Leban, A. M. Le Marechal, *Tetrahedron Lett.* **2003**, *44*, 2363–2366; b) T. I. EI-Emary, A. Khalil, G. A. M. EI-Hag Ali, A. A. A. M. EI-Adasy, *Phosphorus, Sulfur Silicon Related Elements* **2005**, *180*, 19–30.
- [4] a) S. L. Cui, X. F. Lin, Y. G. Wang, *Org. Lett.* **2006**, *8*, 4517–4520; b) S. L. Cui, J. Wang, Y. G. Wang, *Org. Lett.* **2007**, *9*, 5023–5025; c) S. L. Cui, J. Wang, Y. G. Wang, *Org. Lett.* **2008**, *10*, 1267–1269; d) S. L. Cui, J. Wang, Y. G. Wang, *Tetrahedron* **2008**, *64*, 487–492; e) W. Lu, W. Z. Song, D. Hong, P. Lu, Y. G. Wang, *Adv. Synth. Catal.* **2009**, *351*, 1768–1772; f) J. She, Z. Jiang, Y. G. Wang, *Synlett* **2009**, 2023–2027; g) P. Lu, Y. G. Wang, *Synlett* **2010**, 165–173.
- [5] a) I. Bae, H. Han, S. Chang, *J. Am. Chem. Soc.* **2005**, *127*, 2038–2039; b) S. H. Cho, E. J. Yoo, I. Bae, S. Chang, *J. Am. Chem. Soc.* **2005**, *127*, 16046–16047; c) E. J. Yoo, I. Bae, S. H. Cho, H. Han, S. Chang, *Org. Lett.* **2006**, *8*, 1347–1350; d) M. P. Cassidy, J. Raushel, V. V. Fokin, *Angew. Chem.* **2006**, *118*, 3226–3229; *Angew. Chem. Int. Ed.* **2006**, *45*, 3154–3157; e) M. Whiting, V. V. Fokin, *Angew. Chem.* **2006**, *118*, 3229–3233;



- Angew. Chem. Int. Ed.* **2006**, *45*, 3157–3161; f) E. J. Yoo, S. Chang, *Org. Lett.* **2008**, *10*, 1163–1166; g) S. H. Cho, S. Chang, *Angew. Chem.* **2008**, *120*, 2878–2881; *Angew. Chem. Int. Ed.* **2008**, *47*, 2836–2839; h) S. H. Cho, S. Chang, *Angew. Chem.* **2007**, *119*, 1929–1932; *Angew. Chem. Int. Ed.* **2007**, *46*, 1897–1900. E. J. Yoo, M. Ahlquist, I. Bae, V. V. Fokin, S. Chang, K. B. Sharpless, *J. Org. Chem.* **2008**, *73*, 5520–5528.
- [6] S. H. Kim, D. Y. Jung, S. Chang, *J. Org. Chem.* **2007**, *72*, 9769–9771.
- [7] CCDC 695148 contains the supplementary crystallographic data for compound **4h**. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing data request @ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12, Union Road, Cambridge CB2 1EZ, UK; fax: (+44)-1223-336033.
- [8] For recent piperidine syntheses, see: a) Y. Hayashi, H. Gotoh, R. Masui, H. Ishikawa, *Angew. Chem.* **2008**, *120*, 4076–4079; *Angew. Chem. Int. Ed.* **2008**, *47*, 4012–4015; b) B. B. Touré, D. G. Hall, *Angew. Chem.* **2004**, *116*, 2035–2038; *Angew. Chem. Int. Ed.* **2004**, *43*, 2001–2004; c) G. A. Cortez, R. R. Schrock, A. H. Hoveyda, *Angew. Chem.* **2007**, *119*, 4618–4622; *Angew. Chem. Int. Ed.* **2007**, *46*, 4534–4538; d) M. Ahari, A. Perez, C. Menant, J. L. Vasse, J. Szymoniak, *Org. Lett.* **2008**, *10*, 2473–2476; e) M. Terada, K. Machioka, K. Sorimachi, *J. Am. Chem. Soc.* **2007**, *129*, 10336–10337; f) J. F. Bower, P. Szeto, T. Gallagher, *Org. Lett.* **2007**, *9*, 4909–4912.