

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51178659>

ChemInform Abstract: Direct Synthesis of Aryl Ketones by Palladium-Catalyzed Desulfonative Addition of Sodium Sulfinates to Nitriles.

ARTICLE *in* CHEMISTRY - A EUROPEAN JOURNAL · NOVEMBER 2011

Impact Factor: 5.73 · DOI: 10.1002/chem.201101252 · Source: PubMed

CITATIONS

59

READS

19

6 AUTHORS, INCLUDING:



Honghua Rao

Capital Normal University

22 PUBLICATIONS 761 CITATIONS

SEE PROFILE



Fuhong Xiao

Xiangtan University

26 PUBLICATIONS 482 CITATIONS

SEE PROFILE



Chao-Jun Li

Stanford University

477 PUBLICATIONS 20,361 CITATIONS

SEE PROFILE



Guo-Jun Deng

Xiangtan University

111 PUBLICATIONS 2,201 CITATIONS

SEE PROFILE

Direct Synthesis of Aryl Ketones by Palladium-Catalyzed Desulfative Addition of Sodium Sulfinates to Nitriles

Jing Liu,^[a] Xianya Zhou,^[a] Honghua Rao,^[b] Fuhong Xiao,^[a] Chao-Jun Li,^{*[b]} and Guo-Jun Deng^{*[a]}


Aryl ketones are fundamental intermediates in many types of organic compound, for example, pharmaceuticals, fragrances, and natural products.^[1] Classical routes to synthesize aryl ketones mainly rely on the Friedel–Crafts acylation of aromatic compounds in the presence of corrosive AlCl_3 ^[2] and the oxidation of the corresponding secondary alcohols.^[3] Transition-metal-catalyzed reactions provide many opportunities for the synthesis of aryl ketones.^[4] Among them, hydroacylation is a useful synthetic method for preparing aryl ketones from aldehydes and olefines through C–H bond activation and subsequent addition.^[5]

Nitriles are normally very stable and widely commercially available. The nucleophilic addition of organometallic reagents to nitriles is an old and well-known reaction for ketone preparation. However, application of this addition reaction has been limited by poor functional group tolerance.^[6] In recent years, the transition-metal-catalyzed addition reaction has resulted in great progress in the synthesis of aryl ketones from nitriles. One of the pioneering works involving the use of nitriles as the starting materials for aryl ketone synthesis was developed by Larock et al. by using palladium as the catalyst. Arylboronic acids and even arene C–H bonds were successfully added to nitriles to generate various aryl ketones by the hydrolysis of the intermediate ketimine in trifluoroacetic acid.^[7] Later, Lu and Zhao,^[8] Miura et al.,^[9] and others^[10] developed various catalytic systems for aryl ketone synthesis from nitriles and arylboronic acids, catalyzed by palladium or rhodium. Cheng et al. developed a nickel-catalyzed addition of arylboronic acids to nitriles by using ZnCl_2 as the Lewis acid, under mild conditions.^[11] However, arylboronic acids normally require several

steps to synthesize from arenes (via the halide and then reaction with borate by using stoichiometric magnesium),^[12] although newer methods of direct boronic acid formation from C–H bonds have recently been explored.^[13] In 2010, Larhed et al. developed a palladium-catalyzed decarboxylative addition of benzoic acids to nitriles, and various aryl ketones can be synthesized with this method.^[14] One of the main advantages of this method is that the carboxylic acids used for this transformation are comparably cheap and are commercially available.^[15] However, this method is mainly suitable for *ortho*-functionalized electron-rich benzoic acids and aliphatic nitriles.

Arenesulfonyl chlorides are active compounds and are used as starting materials for preparing compounds containing a sulfonyl group.^[16] They are also used as aryl sources for C–C bond-forming reactions through desulfative Heck-type reactions under harsh conditions ($> 130^\circ\text{C}$).^[17] Compared to the active and moisture sensitive arenesulfonyl chloride, sulfinic acid sodium salts are comparably stable and easy to handle. Thus, sulfinic acid sodium salts have the potential to serve as the ideal aryl source for C–C bond-forming reactions through release of SO_2 . However, sulfinic acids (or their sodium salts) are mainly used as sulfonylation reagents^[18] and rarely used as the aryl source in desulfative reactions.^[19] Recently, we developed a palladium-catalyzed desulfative Heck-type reaction of aromatic sulfinic acid sodium salts with various olefins under mild conditions.^[20] Herein, we report the direct synthesis of aryl ketones by palladium-catalyzed desulfative addition of aryl sulfinic acid sodium salts to nitriles under mild conditions (Scheme 1).

We began our investigation by screening various palladium salts for the desulfative addition reaction between benzonitrile (**1a**) and *p*-toluenesulfinic acid sodium salt (**2a**) in the presence of trifluoroacetic acid, and the results are summarized in Table 1. If benzonitrile was reacted with **2a** (2 equiv) in the presence of $\text{Pd}(\text{OAc})_2$ (5 mol %) and 1,10-phenanthroline (1,10-phen; 10 mol %) in 1,4-dioxane/water (3:2), the desired product was observed in 53 % yield, as de-




Scheme 1. General reaction scheme.

[a] J. Liu, X. Zhou, F. Xiao, Prof. Dr. G.-J. Deng
Key Laboratory of Environmental Friendly Chemistry and
Application of Ministry of Education
College of Chemistry, Xiangtan University
Xiangtan 411105 (China)
Fax: (+86) 731-58292477
E-mail: gjdeng@xtu.edu.cn

[b] Dr. H. Rao, Prof. Dr. C.-J. Li
Department of Chemistry, McGill University
Montreal, QC, H3 A2K6 (Canada)
Fax: (+1) 514-398-3797
E-mail: cj.li@mcgill.ca

Supporting information for this article is available on the WWW
under <http://dx.doi.org/10.1002/chem.201101252>.

Table 1. Optimization of the reaction conditions.^[a]

				
	Catalyst	Ligand	Solvent	Yield [%] ^[b]
1	Pd(OAc) ₂	1,10-phen	1,4-dioxane	53
2	Pd(O ₂ CCF ₃) ₂	1,10-phen	1,4-dioxane	54
3	PdCl ₂	1,10-phen	1,4-dioxane	41
4	[Pd(acac) ₂]	1,10-phen	1,4-dioxane	42
5	[Pd ₂ (dba) ₃]	1,10-phen	1,4-dioxane	58
6	[Pd(cod)Cl ₂]	1,10-phen	1,4-dioxane	52
7	[Pd(NH ₃) ₄ Cl ₂]	1,10-phen	1,4-dioxane	40
8	PdO	1,10-phen	1,4-dioxane	70
9	Pd(OH) ₂	1,10-phen	1,4-dioxane	91
10	Pd(OH) ₂	dipyridine	1,4-dioxane	60
11	Pd(OH) ₂	DMAP	1,4-dioxane	2
12	Pd(OH) ₂	DBU	1,4-dioxane	trace
13	Pd(OH) ₂	PPh ₃	1,4-dioxane	trace
14	Pd(OH) ₂	1,10-phen	DMF	3
15	Pd(OH) ₂	1,10-phen	anisole	45
16	Pd(OH) ₂	1,10-phen	diglyme	56
17	Pd(OH) ₂	1,10-phen	THF	67
18 ^[c]	Pd(OH) ₂	1,10-phen	1,4-dioxane	71
19 ^[d]	Pd(OH) ₂	1,10-phen	1,4-dioxane	63
20 ^[e]	Pd(OH) ₂	1,10-phen	1,4-dioxane	47

[a] Reaction conditions: **1a** (0.2 mmol), **2a** (0.4 mmol), catalyst (5 mol %), ligand (10 mol %), CF₃CO₂H (0.2 mL), solvent (0.3 mL), H₂O (0.2 mL), 110 °C, 20 h, under air; acac = acetylacetonate, dba = dibenzylideneacetone, cod = 1,5-cyclooctadiene, DMAP = 4-dimethylaminopyridine, DBU = 1,8-diazabicyclo[5.4.0]undec-7-ene. [b] GC yield based on **1a**. [c] 1,4-Dioxane/H₂O = 1:1. [d] At 90 °C. [e] No CF₃CO₂H was used.

tested by GC-MS and ¹H NMR methods (Table 1, entry 1). Subsequently, various palladium salts were tested for this desulfinate addition reaction and moderate yields were observed (Table 1, entries 2–7). The reaction yield can be improved to 70 % if PdO was used (Table 1, entry 8). However, the best yield was obtained with Pd(OH)₂ as the catalyst, for which the desired product was obtained in 91 % yield (Table 1, entry 9). Other ligands were also investigated for this transformation but were less efficient (Table 1, entries 10–13). The effect of solvents on this reaction was also investigated. A low yield was obtained if the reaction was carried out in DMF (Table 1, entry 14) although the reactions in anisole, diglyme, and THF gave the desired product in moderate yields (Table 1, entries 15–17). The ratio of 1,4-dioxane to water also affected the reaction yield significantly, decreasing it to 71 % if the ratio of was changed to 1:1 (Table 1, entry 18). Decreasing the reaction temperature also decreased the yield (Table 1, entry 19). Trifluoroacetic acid played an important role in this addition reaction and only 47 % yield was achieved in its absence (Table 1, entry 20).^[21]

Next, the desulfinate addition of *p*-toluenesulfinic acid sodium salt **2a** to various nitriles was conducted under the optimized conditions, and the results are summarized in Table 2. The palladium-catalyzed desulfinate addition reaction was successfully extended to various substituted aryl

nitriles **1b–g**, and various asymmetric diaryl ketones were obtained in moderate to good yields (Table 2, entries 2–7). A series of functional groups, including methyl, methoxy, fluoro, chloro, and acetyl, were tolerated under the optimized reaction conditions. A moderate yield was obtained when 2-methylbenzonitrile was reacted with **2a** (Table 2, entry 8). Aliphatic nitriles also worked well for this reaction. Phenylacetone nitrile and its substituted derivatives were

Table 2. Desulfinate addition of **2a** with various nitriles.^[a]

	Nitrile	Product	Yield [%] ^[b]
1	Ph-CN		3a 78
2			3b 72
3			3c 80
4			3d 81
5			3e 73
6			3f 74
7			3g 60
8			3h 51
9	Ph-CH ₂ -CN		3i 82
10			3j 80 (78) ^[c]
11			3k 85
12			3l 83
13			3m 82
14			3n 70

Table 2. (Continued)

Nitrile	Product	Yield [%] ^[b]
15 ^[d]	1o	3o 42
16 ^[d]	1p	3p 52
17 ^[d]	1q	3q 65

[a] Reaction conditions, unless otherwise noted: **1** (0.2 mmol), **2a** (0.4 mmol), Pd(OH)₂ (5 mol %), 1,10-phen (10 mol %), CF₃CO₂H (0.2 mL), 1,4-dioxane (0.3 mL), H₂O (0.2 mL), 110 °C, 20 h, under air. [b] Isolated yield based on **1**. [c] The yield in parenthesis was on a 5 mmol scale. [d] Pd(OH)₂ (10 mol %) and 2,2'-bipyridine (10 mol %) were used.

smoothly reacted with **2a** and gave the corresponding alkyl aryl ketones in good yields (Table 2, entries 9–14). Pentanenitrile, octanenitrile, and decanenitrile were also reacted with **2a**, although under modified reaction conditions, and gave the desired products in 42, 52, and 65 % yield, respectively (Table 2, entries 15–17).

Additionally, the results of the reaction between various aryl sulfinic acid sodium salts and 4-methylbenzonitrile (**1b**) are presented in Table 3. Benzenesulfinic acid sodium salt (**2b**) was successfully reacted with **1b** and gave the corresponding product in 76 % yield (Table 3, entry 1). Various functional groups, including methoxy, *tert*-butyl, fluoro, bromo, and trifluoromethyl, were tolerated, and the desired products were obtained in moderate to good yields (Table 3, entries 2–6). A more bulky substrate, 2-naphthylsulfinic acid sodium salt (**2h**), also efficiently reacted with **1b** and gave the product in 53 % yield (Table 3, entry 7).


A plausible mechanism to rationalize this transformation is illustrated in Scheme 2. Similar to the Pd^{II}-catalyzed decarboxylative addition of benzoic acids to nitriles,^[14] the proposed mechanism involves the following six steps: 1) coordination of the arenesulfinic acid (or its sodium salt) to palladium(II) to give complex **I**; 2) the desulfurative reaction of the sulfinic acid to generate aryl–palladium complex **II**; 3) coordination of the nitrile group to form complex **III**; 4) 1,2-carbopalladation of the nitrile to generate ketimine complex **IV**; 5) protonation of the ketimine by the acid to give the free ketimine and a palladium(II) species; and 6) hydrolysis of the ketimine under acidic conditions to afford the desired final aryl ketone.

In summary, we have developed a new Pd-catalyzed method that allows the direct synthesis of aryl ketones from aromatic sulfinic acid sodium salts and inexpensive nitrile compounds. Ligands and solvents played an important role in the reaction yield. A combination of Pd(OH)₂ and 1,10-phenanthroline in 1,4-dioxane/water (3:2) in air gave the desired product in the best yield. Aromatic and aliphatic nitriles were successfully reacted with arenesulfinic acid

Table 3. Desulfurative addition of **1b** with various sodium sulfonates.^[a]

Sodium sulfonate	Product	Yield [%] ^[b]
1	2b	3r 76
2	2c	3s 65
3	2d	3t 73
4	2e	3u 61
5 ^[c]	2f	3v 50
6	2g	3w 35
7	2h	3x 53

[a] Reaction conditions, unless otherwise noted: **1b** (0.2 mmol), **2** (0.4 mmol), Pd(OH)₂ (5 mol %), 1,10-phen (10 mol %), CF₃CO₂H (0.2 mL), 1,4-dioxane (0.3 mL), H₂O (0.2 mL), 110 °C, 20 h, under air. [b] Isolated yield based on **1b**. [c] Anisole was used instead of 1,4-dioxane.



Scheme 2. Proposed mechanism.

sodium salts to form aryl ketones in moderate to good yields. This procedure provides a rapid and convenient method for the small-scale synthesis of aryl ketones.^[22] Our current efforts are focused on further mechanistic studies to improve the catalytic efficiency and to overcome limitations in substrate scope.

Experimental Section

Representative experimental procedure (3a, CAS: 134-84-9): A reaction vessel (10 mL) was charged with a mixture of palladium hydroxide

(1.4 mg, 0.01 mmol), 1,10-phenanthroline (3.6 mg, 0.02 mmol), sodium *p*-toluenesulfonate (**2a**, 71.2 mg, 0.4 mmol), benzonitrile (**1a**, 20 μ L, 0.2 mmol), and trifluoroacetic acid (0.2 mL) in 1,4-dioxane/water (0.5 mL, 3:2). The reaction vessel was closed and the resulting solution was stirred at 110°C for 20 h. After cooling to room temperature, the volatiles were removed under vacuum and the residue was purified by column chromatography (silica gel, petroleum ether/ethyl acetate, 50:1) to give **3a** as a white solid (30.5 mg, 78%).

Acknowledgements

This work was supported by the National Natural Science Foundation of China (20902076), the "Academic Leader Program" of Xiangtan University and the "One Hundred Talent Project" of Hunan Province. We are also grateful to the Canada Research Chair (Tier I) foundation (to C.-J. Li).

Keywords: aryl ketones • desulfonation • ketones • nitriles • palladium • sulfinic acids

- [1] H. Surburg, J. Panten, *Common Fragrance and Flavor Materials*, 5th ed., Wiley-VCH, Weinheim, **2006**.
- [2] G. Sartori, R. Maggi, *Advances in Friedel–Crafts Acylation Reactions*, CRC Press, New York, **2010**.
- [3] a) G. Tojo, M. Fernández, *Oxidation of Alcohols to Aldehydes and Ketones*, Springer, Berlin, **2006**; b) J. E. Bäckvall, *Modern Oxidation Methods*, Wiley-VCH, Weinheim, **2010**; c) G. E. Dobereiner, R. H. Crabtree, *Chem. Rev.* **2010**, *110*, 681.
- [4] For selected recent examples, see: a) J. W. Ruan, O. Saidi, J. A. Iggo, J. L. Xiao, *J. Am. Chem. Soc.* **2008**, *130*, 10510; b) A. Takemiya, J. F. Hartwig, *J. Am. Chem. Soc.* **2006**, *128*, 14800; c) S. Ko, B. Kang, S. Chang, *Angew. Chem.* **2005**, *117*, 459; *Angew. Chem. Int. Ed.* **2005**, *44*, 455; d) L. V. Desai, D. Ren, T. Rosner, *Org. Lett.* **2010**, *12*, 1032; e) M. Pucheault, S. Darses, J. P. Genet, *J. Am. Chem. Soc.* **2004**, *126*, 15356; f) M. H. Savari, H. Sharghi, *J. Org. Chem.* **2004**, *69*, 6953; g) H. Firouzabadi, N. Iranpoor, F. Nowrouzi, *Tetrahedron* **2004**, *60*, 10843; h) A. Bensari, N. T. Zaveri, *Synthesis* **2003**, 267.
- [5] For recent reviews on hydroacylation, see: a) M. C. Willis, *Chem. Rev.* **2010**, *110*, 725, and references therein; b) D. C. Colby, R. G. Bergman, J. A. Ellman, *Chem. Rev.* **2010**, *110*, 624; c) G. Fu in *Modern Rhodium-Catalyzed Organic Reactions* (Ed.: P. Evans), Wiley-VCH, Weinheim, **2005**, pp. 79–91.
- [6] For reviews, see: a) B. I. Roman, N. De Kimpe, C. V. Stevens, *Chem. Rev.* **2010**, *110*, 5914; b) F. F. Fleming, Q. Z. Wang, *Chem. Rev.* **2003**, *103*, 2035.
- [7] a) C. X. Zhou, R. C. Larock, *J. Am. Chem. Soc.* **2004**, *126*, 2302; b) C. X. Zhou, R. C. Larock, *J. Org. Chem.* **2006**, *71*, 3551.
- [8] a) B. W. Zhao, X. Y. Lu, *Org. Lett.* **2006**, *8*, 5987; b) B. W. Zhao, X. Y. Lu, *Tetrahedron Lett.* **2006**, *47*, 6765.
- [9] K. Ueura, S. Miyamura, T. Satoh, M. Miura, *J. Organomet. Chem.* **2006**, *691*, 2821.
- [10] a) H. Shimizu, M. Murakami, *Chem. Commun.* **2007**, 2855; b) R. M. Ceder, G. Muller, M. Ordinas, J. I. Ordinas, *Dalton Trans.* **2007**, 83.
- [11] Y. Wong, K. Parthasarathy, C. Cheng, *Org. Lett.* **2010**, *12*, 1736.
- [12] a) D. G. Hall, *Boronic Acids: Preparation, Applications in Organic Synthesis and Medicine*, Wiley-VCH, Weinheim, **2005**; b) N. Miyaura, A. Suzuki, *Chem. Rev.* **1995**, *95*, 2457; c) N. Miyaura, *Top. Curr. Chem.* **2002**, *219*, 11.
- [13] For an excellent review on C–H boronation, see: I. A. I. Mkhaliid, J. H. Barnard, T. B. Marder, J. M. Murphy, J. F. Hartwig, *Chem. Rev.* **2010**, *110*, 890; for selected recent examples, see: a) D. W. Robbins, T. A. Boebel, J. F. Hartwig, *J. Am. Chem. Soc.* **2010**, *132*, 4068; b) C. S. Wei, C. A. Jiménez-Hyos, M. F. Videau, J. F. Hartwig, M. B. Hall, *J. Am. Chem. Soc.* **2010**, *132*, 3078; c) J. M. Murphy, J. D. Lawrence, K. Kawamura, C. Incarvito, J. F. Hartwig, *J. Am. Chem. Soc.* **2006**, *128*, 13684; d) H. Chen, S. Schlecht, T. C. Semple, J. F. Hartwig, *Science* **2000**, *287*, 1995; e) J. Y. Cho, M. K. Tse, D. Holmes, R. E. Maleczka, M. R. Smith, *Science* **2002**, *295*, 305; f) V. J. Olsson, K. J. Szabó, *Angew. Chem.* **2007**, *119*, 7015; *Angew. Chem. Int. Ed.* **2007**, *46*, 6891.
- [14] J. Lindh, P. Sjöberg, M. Larhed, *Angew. Chem.* **2010**, *122*, 7899; *Angew. Chem. Int. Ed.* **2010**, *49*, 7733.
- [15] For a review on the use of carboxylic acids as substrates, see: L. J. Goossen, N. Rodriguez, K. Goossen, *Angew. Chem.* **2008**, *120*, 3144; *Angew. Chem. Int. Ed.* **2008**, *47*, 3100.
- [16] a) P. Page, *Organosulfur Chemistry: Stereochemical Aspects*, Academic Press, San Diego, **1998**; b) S. Dubbaka, P. Vogel, *Angew. Chem.* **2005**, *117*, 7848; *Angew. Chem. Int. Ed.* **2005**, *44*, 7674; c) N. S. Simpkins, *Sulfones in Organic Synthesis*, Pergamon Press, Oxford, **1993**.
- [17] For desulfinative C–C bond formation by using sulfonyl chlorides as starting materials, see: a) A. Kasahara, T. Izumi, N. Kudou, H. Azami, S. Yamamoto, *Chem. Ind.* **1988**, 51; b) A. Kasahara, T. Izumi, K. Miyamoto, T. Sakai, *Chem. Ind.* **1989**, 192; c) M. Miura, H. Hashimoto, K. Itoh, M. Nomura, *Tetrahedron Lett.* **1989**, *30*, 975; d) M. Miura, H. Hashimoto, K. Itoh, M. Nomura, *J. Chem. Soc. Perkin Trans. 1* **1990**, 2207; e) S. Reddy Dubbaka, P. Vogel, *J. Am. Chem. Soc.* **2003**, *125*, 15292; f) S. Reddy Dubbaka, P. Vogel, *Org. Lett.* **2003**, *5*, 95; g) S. Reddy Dubbaka, P. Steunenberg, P. Vogel, *Synlett* **2004**, 1235; h) S. Reddy Dubbaka, P. Vogel, *Adv. Synth. Catal.* **2004**, *346*, 1793; i) S. R. Dubbaka, P. Vogel, *Chem. Eur. J.* **2005**, *11*, 2633; j) S. R. Dubbaka, D. B. Zhao, Z. F. Fei, C. M. R. Volla, P. J. Dyson, P. Vogel, *Synlett* **2006**, 3155; k) C. M. Rao Volla, P. Vogel, *Angew. Chem.* **2008**, *120*, 1325; *Angew. Chem. Int. Ed.* **2008**, *47*, 1305; l) C. M. Rao Volla, S. R. Dubbaka, P. Vogel, *Tetrahedron* **2009**, *65*, 504; m) C. M. Rao Volla, D. Markovic, S. R. Dubbaka, P. Vogel, *Eur. J. Org. Chem.* **2009**, 6281.
- [18] For recent examples of sulfonylation by using sodium sulfinates, see: a) F. Sandrinelli, S. Perrio, P. Beslin, *Org. Lett.* **1999**, *1*, 1177; b) W. Yang, C. Berthelette, *Tetrahedron Lett.* **2002**, *43*, 4537; c) S. Cacchi, G. Fabrizi, A. Goggiamani, L. Parisi, R. Berning, *J. Org. Chem.* **2004**, *69*, 5608; d) W. Zhu, D. Ma, *J. Org. Chem.* **2005**, *70*, 2696; e) C. Martin, F. Sandrinelli, C. Perrio, S. Perrio, M. Lasne, *J. Org. Chem.* **2006**, *71*, 210; f) Z. Guan, W. Zuo, L. Zhao, Z. Ren, Y. M. Liang, *Synthesis* **2007**, 1465; g) J. Drabowicz, M. Kaiatowska, P. Kielbasinski, *Synthesis* **2008**, 3563; h) B. Sreedhar, M. Reddy, P. Reddy, *Synlett* **2008**, 1949; i) H. Kim, A. Kasper, Y. Park, C. Wooten, J. Hong, E. Moon, M. Dewhirst, *Org. Lett.* **2009**, *11*, 89; j) A. Varela-Álvarez, D. Marković, P. Vogel, J. Sordo, *J. Am. Chem. Soc.* **2009**, *131*, 9547; k) C. Liu, M. Li, D. Cheng, C. Yang, S. Tian, *Org. Lett.* **2009**, *11*, 2543; l) M. Reddy, P. Reddy, B. Sreedhar, *Adv. Synth. Catal.* **2010**, *352*, 1861; m) Y. Li, K. Cheng, X. Lu, J. Sun, *Adv. Synth. Catal.* **2010**, *352*, 1876; n) M. Ueda, J. F. Hartwig, *Org. Lett.* **2010**, *12*, 92; o) K. Maloney, J. Kuethe, K. Linn, *Org. Lett.* **2011**, *13*, 102.
- [19] For early investigations on desulfinative reactions that use sulfinic acid (salt), see: a) K. Garves, *J. Org. Chem.* **1970**, *35*, 3273; b) R. Selke, W. Thiele, *J. Prakt. Chem.* **1971**, *313*, 875; c) E. Wenkert, T. W. Ferreira, E. L. Michelotti, *J. Chem. Soc. Chem. Commun.* **1979**, 637; d) for high efficiency biaryl synthesis by using sulfinic acid sodium salts and aromatic bromides, see: K. Sato, T. Okoshi, Patent, US5159082, **1992**.
- [20] X. Y. Zhou, J. Y. Luo, J. Liu, S. M. Peng, G. J. Deng, *Org. Lett.* **2011**, *13*, 1432.
- [21] A similar yield was observed if the reaction was carried out under an argon atmosphere and a much lower yield was obtained in oxygen.
- [22] The price of 4-methylbenzenesulfinic acid sodium salt (**2a**, CAS: 824-79-3) is about 0.2 \$ per gram, whereas the price of 4-methylbenzeneboronic acid (CAS: 5720-05-8) is about 20 \$ per gram (both from Alfa Aesar, 2011).

Received: April 23, 2011

Published online: May 30, 2011