

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/44619422>

ChemInform Abstract: An Unexpected Phosphine-Catalyzed Regio- and Diastereoselective [4 + 1] Annulation Reaction of Modified Allylic Compounds with Activated Enones.

ARTICLE *in* CHEMISTRY - AN ASIAN JOURNAL · JULY 2010

Impact Factor: 4.59 · DOI: 10.1002/asia.201000193 · Source: PubMed

CITATIONS

51

READS

24

2 AUTHORS, INCLUDING:



Zuliang Chen

East China Normal University

10 PUBLICATIONS 174 CITATIONS

SEE PROFILE

An Unexpected Phosphine-Catalyzed Regio- and Diastereoselective [4 + 1] Annulation Reaction of Modified Allylic Compounds with Activated Enones


Zuliang Chen^[a] and Junliang Zhang^{*[a, b]}

Nucleophilic phosphines are known to be useful mild catalysts for the synthesis of cyclic and heterocyclic compounds.^[1] Generally, tertiary-phosphine-mediated annulations are triggered by nucleophilic addition of phosphines to activated compounds. The resultant zwitterionic intermediates can react with various electrophiles such as aldehydes, imines, and activated polarized C=C bonds to furnish cyclic compounds.^[2] Herein, we report a phosphine-catalyzed highly diastereoselective synthesis of tetrasubstituted 2,3-dihydrofurans,^[3] which are subunits of a range of biologically active compounds (e.g., aflatoxin B₁ and clerodin),^[4] via [4+1] annulations of Morita–Baylis–Hillman carbonates with activated enones. In these annulations, Morita–Baylis–Hillman carbonates act as the one-carbon unit, which is distinguished from the modified allylic compounds reported by Lu et al. as the three-carbon units, such as in [3+2],^[5h,5i] [3+3],^[5n] [3+4],^[5m] and [3+6]^[5e] annulation reactions.

In recent years, Morita–Baylis–Hillman adducts have been illustrated as suitable starting materials for the synthesis of a variety of multifunctional compounds.^[5] During our study of the new chemistry of electron-deficient enynes,^[6] we became interested in the annulation between Morita–Baylis–Hillman carbonate **1a** and conjugated yne-enone **2a**. We were pleased to find that the reaction proceeded smoothly in toluene at room temperature under catalysis with 10 mol % PPh₃, leading to a [4+1] adduct, 2,3-dihydro-

furan **3a**, in 72 % yield of isolated product with 20:1 diastereoselectivity (Table 1, entry 1). The structure of **3a** was established by X-ray crystallography analysis of the analogous product **3f** (R²=*p*-BrC₆H₄; Figure 1).^[7] To improve the yield of **3a**, various solvents and tertiary phosphines were tested and the results are summarized in Table 1. A higher product yield was obtained when the reaction was performed in CH₂Cl₂, while the selectivity was slightly decreased (Table 1, entry 2). Good yields with high selectivity were also obtained in DCE, Et₂O, and 1,4-dioxane, albeit the reactions require a longer time to go to completion (Table 1, en-

Table 1. Screening the reaction conditions of annulation reaction of compound **1a** and **2a**.^[a]



| Entry | Solvent | <i>t</i> [h] | Yield [%] ^[b] | d.r. ^[c] |
|-------------------|---------------------------------|--------------|--------------------------|---------------------|
| 1 | toluene | 26 | 72 | 20:1 |
| 2 | CH ₂ Cl ₂ | 12 | 81 | 18:1 |
| 3 | DCE | 29 | 81 | 20:1 |
| 4 | Et ₂ O | 48 | 82 | 20:1 |
| 5 | 1,4-dioxane | 48 | 82 | 20:1 |
| 6 | hexane | 48 | 75 | 20:1 |
| 7 | THF | 24 | 75 | 20:1 |
| 8 | EtOAc | 48 | 74 | 20:1 |
| 9 | DMF | 13 | 72 | 20:1 |
| 10 | DMSO | 12 | 70 | 20:1 |
| 11 ^[d] | CH ₃ CN | 12 | 0 | – |
| 12 ^[e] | CH ₃ CN | 24 | 0 | – |
| 13 ^[f] | CH ₃ CN | 24 | 0 | – |
| 14 ^[g] | CH ₃ CN | 4 | 42 | 5:1 |
| 15 | CH ₃ CN | 11 | 82 | 20:1 |

[a] Reaction conditions: Under Ar, a mixture of **1a** (65.0 mg, 0.3 mmol), **2a** (62.0 mg, 0.2 mmol), and PPh₃ (5.3 mg, 0.02 mmol) in solvent (2 mL) was stirred at room temperature. [b] Yield of isolated product. [c] The diastereomeric ratio was determined by ¹H NMR spectroscopic analysis of the crude reaction mixture. [d] 10 mol % PCy₃ was used. [e] 10 mol % trifuran-2-ylphosphine was used. [f] 10 mol % tri-*o*-tolylphosphine was used. [g] The reaction was carried out at reflux.

[a] Z. Chen, Prof. Dr. J. Zhang
Shanghai Key Laboratory of Green Chemistry and Chemical Processes
Department of Chemistry
East China Normal University
3663 N. Zhongshan Road, Shanghai 200062 (P.R. China)
Fax: (+86) 21-6223-3213
E-mail: jlzhang@chem.ecnu.edu.cn

[b] Prof. Dr. J. Zhang
State Key Laboratory of Organometallic Chemistry,
Shanghai Institute of Organic Chemistry
Chinese Academy of Sciences
354 Fenglin Road, Shanghai 200032, (P.R. China)

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/asia.201000193>.




Figure 1. ORTEP representation of compound **3f**. Thermal ellipsoids for non-hydrogen atoms are shown at the 30% probability level.

tries 3–5). No reaction occurs for other tested phosphines such as PCy_3 , trifuran-2-ylphosphine, and tri-*o*-tolylphosphine (Table 1, entries 11–13). The product yield and the diastereoselectivity decreased dramatically when the reaction was carried out at reflux in CH_3CN with 10 mol % PPh_3 (Table 1, entry 14). Finally, it was found that CH_3CN was the best solvent, providing **3a** in 82% yield of isolated product with a 20:1 d.r. in 11 h at room temperature (Table 1, entry 15).

With the optimal reaction conditions in hand, we next examined the scope of this phosphine-catalyzed [4+1] annulation reaction, and the results are summarized in Table 2. These results showed that all three substituents (R^{1-3}) could be not only aromatic but also aliphatic groups to afford the corresponding tetrasubstituted 2,3-dihydrofurans. The annulations of **1a** with conjugated yne-enones **2b–d** bearing a 4-


Table 2. Synthesis of highly functionalized 2,3-dihydrofurans.^[a]

| Entry | Enyne 2 $\text{R}^1/\text{R}^2/\text{R}^3$ | <i>t</i> [h] | 3 Yield [%] ^[b] | d.r. ^[c] |
|-------------------|--|--------------|--------------------------------------|---------------------|
| 1 | Ph/Ph/Ph (2a) | 11 | 3a (82) | 20:1 |
| 2 | Ph/Ph/4- $\text{NO}_2\text{C}_6\text{H}_4$ (2b) | 10 | 3b (63) | 14:1 |
| 3 | Ph/Ph/4- MeOC_6H_4 (2c) | 18 | 3c (70) | 30:1 |
| 4 ^[d] | Ph/Ph/1-cyclohexenyl (2d) | 13 | 3d (60) | 20:1 |
| 5 | Ph/4- ClC_6H_4 /Ph (2e) | 8 | 3e (81) | 16:1 |
| 6 | Ph/4- BrC_6H_4 /Ph (2f) | 8 | 3f (76) | 13:1 |
| 7 | 2- BrC_6H_4 /Ph/Ph (2g) | 10 | 3g (80) | 10:1 |
| 8 ^[d] | Ph/4- MeOC_6H_4 /Ph (2h) | 12 | 3h (59) | 11:1 |
| 9 | 4- ClC_6H_4 /Ph/Ph (2i) | 8 | 3i (74) | 14:1 |
| 10 ^[d] | 4- MeOC_6H_4 /Ph/Ph (2j) | 16 | 3j (59) | 10:1 |
| 11 ^[d] | Ph/Me/Ph (2k) | 12 | 3k (72) | 20:1 |
| 12 ^[d] | Ph/Me/1-naphthyl (2l) | 22 | 3l (67) | 16:1 |
| 13 ^[d] | <i>n</i> Butyl/Me/Ph (2m) | 20 | 3m (47) | 20:1 |

[a] Reaction conditions: Under Ar, a mixture of **1a** (97.0 mg, 0.45 mmol), **2** (0.3 mmol), and PPh_3 (7.9 mg, 0.03 mmol) in solvent (3 mL) was stirred at room temperature. [b] Yield of isolated product. [c] The diastereomeric ratio was determined by ^1H NMR spectroscopic analysis of the crude reaction mixture. [d] 20 mol % PPh_3 was used.

nitrophenyl, 4-methoxyphenyl, or 1-cyclohexenyl substituent at the alkyne (R^3) worked well to give the expected products in 60–70% yields (Table 2, entries 2–4). Furthermore, substituents such as chloro, bromo, and methoxyl groups at R^1 or the R^2 phenyl ring are also tolerant (Table 2, entries 5–10). Finally, the substituents R^1 and R^2 could also be an aliphatic group. For example, annulations of **2k–m** with **1a** could give 2,3-dihydrofurans **3k–m**, albeit with a higher catalyst loading (20 mol % PPh_3) (Table 2, entries 11–13). It is noteworthy that all these annulations are highly diastereoselective (d.r. = 10:1–30:1).


Since the alkyne moiety of yne-enones **2** was not involved in the [4+1] annulations, we envisaged that other types of activated enones may undergo similar transformation. Indeed, this hypothesis was proved by further investigation [Eqs. (1)–(5)]. For example, the reaction of **1a** with chalcone



2n could give the desired product **3n** in 25% yield along with 63% yield of recovered **2n**, indicating that the alkyne moiety in the yne-enones acts as an activating group to make them more reactive by lowering the LUMO energy level.^[6h] Pleasingly, activated enones **2o–q** are appropriate substrates to give the corresponding cycloadducts **3o–q** in 60–67% yields. In these cases, THF was found to be a better solvent [Eqs. (2)–(4)]. Moreover, the methyl 2-((*tert*-butoxycarbonyl-oxy)(phenyl)methyl)acrylate **1b** could also react with **2a** in refluxing dioxane to give product **3r** in 60% yield as a single diastereomer [Eq. (5)].

The detailed mechanism of this unexpected annulation reaction has not been clarified, and the reason why it did not undergo [3+2] annulations^[5] is unclear. According to these

experimental results and some literature precedents^[2a,5h,j] for phosphine-catalyzed reactions, we propose a plausible mechanism for this unexpected phosphine-catalyzed regio- and diastereoselective [4+1] annulation reaction as follows (Scheme 1). The reaction might be triggered by the forma-



Scheme 1. A plausible mechanism.

tion of the phosphonium salt **A** via the addition–elimination mechanism.^[2a] Subsequent deprotonation by the in situ generated *tert*-butoxide anion would generate ylide **B**. Nucleophilic addition of the resultant intermediate **B** to conjugated yne-enone **2a** would yield intermediate **C** via favored γ -carbon addition. Intermediate **C** would isomerize into intermediate **D** under the conditions via 1,3-H shift. Subsequent intramolecular S_N2' nucleophilic substitution would give the final product (**3r**) and regenerate Ph_3P . The reaction pathway via addition of α -carbon of the ylide **B** to **2a** is unfavored owing to the steric hindrance of the bulky phosphine group and the substituent R^1 of **2a**.

In summary, we have discovered an unexpected phosphine-catalyzed regio- and diastereoselective [4+1] annulation reaction of readily available Morita–Baylis–Hillman carbonates and activated enones, which provided a simple, efficient, and practical method for the synthesis of highly functionalized 2,3-dihydrofurans. Further studies will focus on the development of related organocatalytic transformations of activated enones, as well as synthetic applications of the functionalized 2,3-dihydrofurans.

Experimental Section

Typical procedure for the synthesis of **3a** (Table 2, entry 1): Under Ar, a mixture of **1a** (97.0 mg, 0.45 mmol), **2a** (93.0 mg, 0.3 mmol), and PPh_3 (8.0 mg, 0.03 mmol) in CH_3CN (3 mL) was stirred at room temperature until the reaction was complete (monitored by TLC). Then the reaction mixture was diluted with water (3 mL) and extracted with CH_2Cl_2 . The organic layers were combined, washed with brine (10 mL), and dried over anhydrous MgSO_4 . After filtration and concentration under reduced pressure, the crude product was purified by flash column chromatography on silica gel (hexanes/AcOEt = 20:1) to afford 100 mg (82%) of **3a** as an oil. ^1H NMR (300 MHz, CDCl_3): δ = 8.15 (d, J = 7.2 Hz, 2H), 7.40–7.11 (m, 13H), 6.28 (s, 1H), 5.81 (s, 1H), 5.30 (d, J = 4.5 Hz, 1H), 4.07 (d, J =

4.5 Hz, 1H), 3.66 ppm (s, 3H); ^{13}C NMR (75 MHz, CDCl_3): δ = 165.68, 157.34, 142.07, 139.07, 131.00, 129.85, 129.58, 128.60, 128.32, 128.18, 127.70, 127.63, 127.16, 126.59, 124.75, 123.78, 96.39, 95.75, 85.59, 85.06, 59.50, 51.82 ppm; MS (EI) m/z (%): 406 [M^+] (54.17), 105 (100); HRMS calcd for $\text{C}_{28}\text{H}_{22}\text{O}_3$: 406.1569, found: 406.1564. For preparative procedures and spectroscopic data for all new compounds, see the Supporting Information.

Acknowledgements

Financial support from NSFC (20702015, 20972054) and STCSM (08dj1400101), Shanghai Shuguang Program (07SG27) are greatly appreciated.

Keywords: annulation reactions • diastereoselectivity • dihydrofurans • enynes • phosphines

- [1] For recent reviews, see: a) X. Lu, C. Zhang, Z. Xu, *Acc. Chem. Res.* **2001**, *34*, 535; b) D. H. Valentine, Jr., J. H. Hillhouse, *Synthesis* **2003**, 317; c) J. L. Methot, W. R. Roush, *Adv. Synth. Catal.* **2004**, *346*, 1035; d) X. Lu, Y. Du, C. Lu, *Pure Appl. Chem.* **2005**, *77*, 1985; e) V. Nair, R. S. Menon, A. R. Sreekanth, N. Abhilash, A. T. Biju, *Acc. Chem. Res.* **2006**, *39*, 520; f) S. E. Denmark, G. L. Beutner, *Angew. Chem.* **2008**, *120*, 1584; *Angew. Chem. Int. Ed.* **2008**, *47*, 1560; g) L. Ye, J. Zhou, Y. Tang, *Chem. Soc. Rev.* **2008**, *37*, 1140.
- [2] For recent reports on phosphine-catalyzed reactions, see: a) Y. Du, X. Lu, C. Zhang, *Angew. Chem.* **2003**, *115*, 1065; *Angew. Chem. Int. Ed.* **2003**, *42*, 1035; b) M. Shi, L. Chen, *Chem. Commun.* **2003**, 1310; c) R. P. Wurz, G. C. Fu, *J. Am. Chem. Soc.* **2005**, *127*, 12234; d) V. Nair, A. Biju, K. Mohanan, E. Suresh, *Org. Lett.* **2006**, *8*, 2213; e) X. Lu, Z. Lu, X. Zhang, *Tetrahedron* **2006**, *62*, 457; f) D. Virieux, A. Guillozic, H. Cristau, *Tetrahedron* **2006**, *62*, 3710; g) A. Scherer, J. Gladysz, *Tetrahedron Lett.* **2006**, *47*, 6335; h) S. Castellano, H. D. G. Fijj, S. S. Kinderman, M. Watanabe, P. de Leon, F. Tamanoi, O. Kwon, *J. Am. Chem. Soc.* **2007**, *129*, 5843; i) D. J. Wallace, R. L. Sidda, R. A. Reamer, *J. Org. Chem.* **2007**, *72*, 1051; j) C. E. Henry, O. Kwon, *Org. Lett.* **2007**, *9*, 3069; k) B. J. Cowen, S. J. Miller, *J. Am. Chem. Soc.* **2007**, *129*, 10988; l) G. S. Creech, O. Kwon, *Org. Lett.* **2008**, *10*, 429; m) Y. Fang, E. N. Jacobsen, *J. Am. Chem. Soc.* **2008**, *130*, 5660.
- [3] For selected reports, see: a) A. Funayama, T. Satoh, M. Miura, *J. Am. Chem. Soc.* **2005**, *127*, 15354; b) A. M. Bernard, A. Frongia, P. P. Piras, F. Secci, M. Spiga, *Org. Lett.* **2005**, *7*, 4565; c) S. Son, G. C. Fu, *J. Am. Chem. Soc.* **2007**, *129*, 1046; d) R. Shen, X. Huang, *Org. Lett.* **2008**, *10*, 3283; e) V. Singh, S. Batra, *Tetrahedron* **2008**, *64*, 4511; f) R. Shen, S. Zhu, X. Huang, *J. Org. Chem.* **2009**, *74*, 4118; g) Q. Wang, H. Hou, L. Hui, C. Yan, *J. Org. Chem.* **2009**, *74*, 7403.
- [4] For reviews, see: a) B. M. Fraga, *Nat. Prod. Rep.* **1992**, *9*, 217; b) A. T. Merritt, S. V. Ley, *Nat. Prod. Rep.* **1992**, *9*, 243; c) J. P. Michael, *Nat. Prod. Rep.* **2000**, *17*, 603; d) T. G. Kilroy, T. P. O'Sullivan, P. J. Guiry, *Eur. J. Org. Chem.* **2005**, 4929.
- [5] Selected leading examples, see: a) D. Basavaiah, A. Jaganmohan Rao, T. Satyanarayana, *Chem. Rev.* **2003**, *103*, 811; b) G. W. Kabalka, B. Venkataiah, G. Dong, *Org. Lett.* **2003**, *5*, 3803; c) C. Cho, M. J. Krische, *Angew. Chem.* **2004**, *116*, 6857; *Angew. Chem. Int. Ed.* **2004**, *43*, 6689; d) G. W. Kabalka, B. Venkataiah, G. Dong, *J. Org. Chem.* **2004**, *69*, 5807; e) Y. Du, J. Feng, X. Lu, *Org. Lett.* **2005**, *7*, 1987; f) L. Ye, X. Sun, Q. Wang, Y. Tang, *Angew. Chem.* **2007**, *119*, 6055; *Angew. Chem. Int. Ed.* **2007**, *46*, 5951; g) D. J. V. C. van Steenis, T. Marcelli, M. Lutz, A. L. Spek, J. H. van Maarseveen, H. Hiemstra, *Adv. Synth. Catal.* **2007**, *349*, 281; h) J. Feng, X. Lu, A. Kong, X. Han, *Tetrahedron* **2007**, *63*, 6035; i) Y.-Q. Jiang, Y.-L. Shi, M. Shi, *J. Am. Chem. Soc.* **2008**, *130*, 7202; j) S. Zheng, X. Lu, *Org. Lett.* **2008**, *10*, 4481; k) W. Zhong, Y. Zhao, W. Su, *Tetrahedron* **2008**, *64*, 5491; l) H. Cui, X.

- Feng, J. Peng, J. Lei, K. Jiang, Y. Chen, *Angew. Chem.* **2009**, *121*, 5847; *Angew. Chem. Int. Ed.* **2009**, *48*, 5737; m) S. Zheng, X. Lu, *Org. Lett.* **2009**, *11*, 3978; n) S. Zheng, X. Lu, *Tetrahedron Lett.* **2009**, *50*, 4532; o) V. Declerck, J. Martinez, F. Lamaty, *Chem. Rev.* **2009**, *109*, 1.
- [6] a) Y. Xiao, J. Zhang, *Angew. Chem.* **2008**, *120*, 1929; *Angew. Chem. Int. Ed.* **2008**, *47*, 1903; b) X. Yu, H. Ren, Y. Xiao, J. Zhang, *Chem. Eur. J.* **2008**, *14*, 8481; c) L. Fan, W. Zhao, W. Jiang, J. Zhang, *Chem. Eur. J.* **2008**, *14*, 9139; d) Y. Xiao, J. Zhang, *Adv. Synth. Catal.* **2009**, *351*, 617; e) Y. Xiao, J. Zhang, *Chem. Commun.* **2009**, 3594; f) F. Liu, J. Zhang, *Angew. Chem.* **2009**, *121*, 5613; *Angew. Chem. Int. Ed.* **2009**, *48*, 5505; g) Z. Chen, J. Zhang, *Chem. Asian J.* **2009**, *4*, 1527; h) W. Li, Y. Xiao, J. Zhang, *Adv. Synth. Catal.* **2009**, *351*, 3083.
- [7] CCDC 736940 (**3 f**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif

Received: March 12, 2010

Published online: May 21, 2010